DESY 02-144
PITHA 02/15
TTP02-24

Top quark pair production and decay at hadron colliders: Predictions at NLO QCD including spin correlations*

W. Bernreuther[a†], A. Brandenburg[b‡], Z.G. Si[a*], P. Uwer[c]

[a]Institut f. Theoretische Physik, RWTH Aachen, 52056 Aachen, Germany

[b]DESY-Theorie, 22603 Hamburg, Germany

[c]Institut f. Theoretische Teilchenphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany

We present results at NLO QCD for hadronic production and decay of top quark pairs, taking into account $t\bar{t}$ spin correlations.

arXiv:hep-ph/0209202 v1 18 Sep 2002

The interactions of the top quark have not been precisely tested so far. The expected data samples of Run II of the Tevatron collider (eventually about 10^4 $t\bar{t}$ pairs per annum) and the LHC ($\sim$ 10^7 $t\bar{t}$ pairs p.a.) will make possible detailed investigations of top quark production and decay. In particular, one may search for possible deviations from the QCD production mechanism and the V-A structure of the top decay by analysing energy and angular distributions of the $t\bar{t}$ decay products. Such studies will be feasible since top quarks do not hadronize due to their very short lifetime, and perturbation theory gives a reliable description of top quark production and decay. To search for new physics effects, precise theoretical predictions for top quark pair production and decay at hadron colliders within the Standard Model are needed. We consider here in particular the following processes:

$$h_1 h_2 \to t\bar{t} + X \to \begin{cases} \ell^+\ell'^- + X \\ \ell^+ j_{\bar{t}} + X \\ \ell^- j_t + X \\ j_t j_{\bar{t}} + X, \end{cases} \qquad (1)$$

where $h_{1,2} = p, \bar{p}$; $\ell = e, \mu, \tau$, and j_t ($j_{\bar{t}}$) denote jets originating from hadronic t ($\bar{t}$) decays. An observable that is very sensitive to the dynamics of top quark production and decay in the above reactions is the double differential angular distribution of the top decay products, e.g. for the dilepton channel:

$$\frac{1}{\sigma}\frac{d^2\sigma}{d\cos\theta_+ \, d\cos\theta_-} = \frac{1}{4}(1 - C\cos\theta_+\cos\theta_-). \qquad (2)$$

In (2), $\theta_\pm$ are the angles between the $\ell^\pm$ direction of flight in the $t(\bar{t})$ rest frame with respect to arbitrary axes $\hat{\mathbf{a}}$ ($\hat{\mathbf{b}}$) which will be specified below. (Terms linear in $\cos\theta_\pm$ are absent in (2) for our choices of $\hat{\mathbf{a}}$, $\hat{\mathbf{b}}$ due to parity invariance of QCD.)

The calculation of the distribution (2) at next-to-leading order (NLO) QCD simplifies enormously in the leading pole approximation (LPA), which amounts to expanding the full amplitudes for (1) around the complex poles of the t and $\bar{t}$ propagators. Only the leading pole terms are kept in this expansion, i.e. one neglects terms of order $\Gamma_t/m_t \approx 1\%$. Within the LPA, the radiative corrections can be classified into factorizable and non-factorizable [1] contributions. Here we consider only the factorizable corrections. In this case the coefficient in the distribution (2) factorizes:

$$C = \kappa_+\kappa_- D. \qquad (3)$$

In (3), D is the $t\bar{t}$ double spin asymmetry

$$D = \frac{N(\uparrow\uparrow) + N(\downarrow\downarrow) - N(\uparrow\downarrow) - N(\downarrow\uparrow)}{N(\uparrow\uparrow) + N(\downarrow\downarrow) + N(\uparrow\downarrow) + N(\downarrow\uparrow)}, \qquad (4)$$

*Report DESY 02-144, PITHA 02/15, TTP02-24
†supported by BMBF contract 05 HT1 PAA 4
‡Speaker at the conference. Supported by a Heisenberg fellowship of D.F.G.

where $N(\uparrow\uparrow)$ denotes the number of $t\bar{t}$ pairs with t ($\bar{t}$) spin parallel to the reference axis $\hat{\mathbf{a}}$ ($\hat{\mathbf{b}}$) etc. The directions $\hat{\mathbf{a}}$ and $\hat{\mathbf{b}}$ can thus be identified with the spin quantization axes for t and $\bar{t}$, and D reflects the strength of the correlation between the t and $\bar{t}$ spins for the chosen axes. Further, $\kappa_\pm$ is the spin analysing power of the charged lepton in the decay $t(\bar{t}) \to b(\bar{b})\ell^\pm\nu(\bar{\nu})$ defined by the decay distribution

$$\frac{1}{\Gamma}\frac{d\Gamma}{d\cos\vartheta_\pm} = \frac{1 \pm \kappa_\pm \cos\vartheta_\pm}{2}, \tag{5}$$

where $\vartheta_\pm$ is the angle between the t ($\bar{t}$) spin and the $\ell^\pm$ direction of flight. From [2] we obtain

$$\kappa_+ = \kappa_- = 1 - 0.015\alpha_s, \tag{6}$$

which means that the charged lepton serves as a perfect analyser of the top quark spin. For hadronic decays $t \to bq\bar{q}'$ one has a decay distribution analogous to (5), and the spin analysing power of jets can be defined. To order α_s [3] (and using $\alpha_s(m_t) = 0.108$),

$$\kappa_b = -0.408 \times (1 - 0.340\alpha_s) = -0.393, \tag{7}$$
$$\kappa_j = +0.510 \times (1 - 0.654\alpha_s) = +0.474, \tag{8}$$

where κ_j is the analysing power of the least energetic non-b-quark jet. The results for C given below are for the dilepton channel. The corresponding results for the single lepton channel are obtained by replacing κ_+ (or κ_-) in Eq. (3) by κ_b or κ_j. The loss of analysing power using hadronic final states is overcompensated by the gain in statistics and in efficiency to reconstruct the t ($\bar{t}$) rest frames.

To compute D at NLO QCD, the differential cross sections for the following parton processes are needed to order α_s^3, where the full information on the t and $\bar{t}$ spins has to be kept:

$$q\bar{q} \to t\bar{t},\ t\bar{t}g;\ gg \to t\bar{t},\ t\bar{t}g;\ q(\bar{q})g \to t\bar{t}q(\bar{q}). \tag{9}$$

Results at NLO QCD for the $\overline{\rm MS}$ subtracted parton cross sections $\hat{\sigma}$ for the above processes with $t\bar{t}$ spins summed over have been obtained in [4], [5], while

$$\hat{\sigma}D = \hat{\sigma}(\uparrow\uparrow) + \hat{\sigma}(\downarrow\downarrow) - \hat{\sigma}(\uparrow\downarrow) - \hat{\sigma}(\downarrow\uparrow) \tag{10}$$



Figure 1. Dimensionless scaling functions $g^{(0)}(\eta)$ (full), $g^{(1)}(\eta)$ (dotted), and $\tilde{g}^{(1)}(\eta)$ (dashed) that determine $\hat{\sigma}D$ in the helicity basis for the gg initial state.

has been computed for different spin quantization axes in [6], [7]. This combination of spin-dependent cross sections can be written as follows

$$\hat{\sigma}D = \frac{\alpha_s^2}{m_t^2}\left[g^{(0)}(\eta) + 4\pi\alpha_s\mathcal{G}^{(1)}\right], \tag{11}$$

with

$$\mathcal{G}^{(1)} = g^{(1)}(\eta) + \tilde{g}^{(1)}(\eta)\ln\left(\frac{\mu^2}{m_t^2}\right), \tag{12}$$

where $\eta = \hat{s}/(4m_t^2) - 1$, and we use a common scale μ for the renormalization and factorization scale. As an example, Fig. 1 shows, for the parton process $gg \to t\bar{t}(g)$, the three functions that determine $\hat{\sigma}D$ for the helicity basis, i.e. $\hat{\mathbf{a}}$ ($\hat{\mathbf{b}}$) is chosen to be the t ($\bar{t}$) direction of flight. Apart from the helicity basis we also consider the beam basis, where $\hat{\mathbf{a}}$ and $\hat{\mathbf{b}}$ are equal to the direction of one of the hadron beams in the laboratory frame, and the so-called off-diagonal basis [8], which is defined by the requirement that $\hat{\sigma}(\uparrow\downarrow) = \hat{\sigma}(\downarrow\uparrow) = 0$ for the process $q\bar{q} \to t\bar{t}$ at tree level.

Table 1 lists our results [9] for the coefficient C in the double differential distribution (2) for

Table 1
Coefficient C defined in Eq. (2) to leading and next-to-leading order in α_s for the parton distribution functions of [10] and $\mu = m_t = 175$ GeV.

	$p\bar{p}$, $\sqrt{s} = 2$ TeV		pp, $\sqrt{s} = 14$ TeV	
	LO	NLO	LO	NLO
$C_{\rm hel.}$	-0.456	-0.389	0.305	0.311
$C_{\rm beam}$	0.910	0.806	-0.005	-0.072
$C_{\rm off.}$	0.918	0.813	-0.027	-0.089

Table 2
Upper part: Dependence on μ of the correlation coefficients C, computed at NLO with the PDF of ref.[10]. Lower part: Correlation coefficients C at NLO for $\mu = m_t$ and different sets of parton distribution functions: GRV98[11], CTEQ5[10], and MRST98 (c-g)[12].

	Tevatron			LHC
μ	$C_{\rm hel.}$	$C_{\rm beam}$	$C_{\rm off.}$	$C_{\rm hel.}$
$m_t/2$	-0.364	0.774	0.779	0.278
m_t	-0.389	0.806	0.813	0.311
$2m_t$	-0.407	0.829	0.836	0.331
PDF	$C_{\rm hel.}$	$C_{\rm beam}$	$C_{\rm off.}$	$C_{\rm hel.}$
GRV98	-0.325	0.734	0.739	0.332
CTEQ5	-0.389	0.806	0.813	0.311
MRST98	-0.417	0.838	0.846	0.315

these different choices of spin quantization axes, using the parton distribution functions (PDFs) of [10]. One sees that at the Tevatron, the spin correlations are largest in the beam and off-diagonal basis. The QCD corrections reduce the LO results by about 10%. For the LHC, the best choice of the above three bases is the helicity basis, and the QCD corrections are small in that case.

Table 2 shows the dependence of the NLO results for C on the scale μ (upper part) and on the choice of the PDFs (lower part). At the Tevatron the spread of results for different PDFs is larger than the scale uncertainty. This rather strong dependence on the PDFs is largely due to the fact that the contributions from $q\bar{q}$ and gg initial states contribute to C with opposite signs. This may offer the possibility to constrain the quark and gluon content of the proton by a precise measurement of the double angular distribution (2).

In all results above we used $m_t = 175$ GeV. A variation of m_t from 170 to 180 GeV changes the results for the Tevatron, again for $\mu = m_t$ and PDFs of [10] as follows: $C_{\rm hel.}$ varies from -0.378 to -0.397, $C_{\rm beam}$ from 0.790 to 0.817, and $C_{\rm off.}$ from 0.797 to 0.822. For the LHC, $C_{\rm hel.}$ changes by less than a percent.

The results above have been obtained without imposing any kinematic cuts. Standard cuts on the top quark transverse momentum and rapidity only have a small effect on C: For the Tevatron, demanding $|\mathbf{k}^T_{t,\bar{t}}| > 15$ GeV and $|r_{t,\bar{t}}| < 2$ leads to the following results: $C_{\rm hel.} = -0.386$, $C_{\rm beam} = 0.815$, $C_{\rm off.} = 0.823$. For the LHC, when imposing the cuts $|\mathbf{k}^T_{t,\bar{t}}| > 20$ GeV and $|r_{t,\bar{t}}| < 3$, we find $C_{\rm hel.} = 0.295$.

In summary, $t\bar{t}$ spin correlations are large effects that can be studied at the Tevatron and the LHC by measuring double angular distributions both in the dilepton and single lepton decay channels. The QCD corrections to these distributions have been computed and are under control. Spin correlations are suited to analyse in detail top quark interactions, search for new effects, and may help to constrain the parton content of the proton.

REFERENCES

1. W. Beenakker, F.A. Berends, A.P. Chapovsky, Phys. Lett. B **454** (1999) 129.
2. A. Czarnecki, M. Jezabek, J.H. Kühn, Nucl. Phys. B **351** (1991) 70.
3. A. Brandenburg, Z.G. Si, P. Uwer, Phys. Lett. B **539** (2002) 235.
4. P. Nason, S. Dawson, R.K. Ellis, Nucl. Phys. B **303** (1988) 607; Nucl. Phys. B **327** (1989) 49.
5. W. Beenakker, H. Kuijf, W.L. van Neerven, J. Smith, Phys. Rev. D **40** (1989) 54; W. Beenakker *et al.*, Nucl. Phys. B **351** (1991) 507.
6. W. Bernreuther, A. Brandenburg, Z.G. Si, Phys. Lett. B **483** (2000) 99.
7. W. Bernreuther, A. Brandenburg, Z.G. Si, P. Uwer, Phys. Lett. B **509** (2001) 53.

8. G. Mahlon, S. Parke, Phys. Lett. B **411** (1997) 173.
9. W. Bernreuther, A. Brandenburg, Z.G. Si, P. Uwer, Phys. Rev. Lett. 87 (2001) 242002.
10. H.L. Lai *et al.* [CTEQ Collaboration], Eur. Phys. J. C **12** (2000) 375.
11. M. Glück, E. Reya, A. Vogt, Eur. Phys. J. C **5** (1998) 461.
12. A.D. Martin, R.G. Roberts, W.J. Stirling, R.S. Thorne, Eur. Phys. J. C **4** (1998) 463.