

On Multiple Zeta Values of Even Arguments

Michael E. Hoffman¹

*Dept. of Mathematics, U. S. Naval Academy
Annapolis, MD 21402 USA*

and

*Deutsches Elektronen-Synchrotron DESY
Platanenalle 6, D-15738 Zeuthen, Germany*

meh@usna.edu

June 8, 2012

Keywords: multiple zeta values, symmetric functions, Bernoulli numbers

MR Classifications: Primary 11M32; Secondary 05E05, 11B68

Abstract

For $k \leq n$, let $E(2n, k)$ be the sum of all multiple zeta values with even arguments whose weight is $2n$ and whose depth is k . Of course $E(2n, 1)$ is the value $\zeta(2n)$ of the Riemann zeta function at $2n$, and it is well known that $E(2n, 2) = \frac{3}{4}\zeta(2n)$. Recently Z. Shen and T. Cai gave formulas for $E(2n, 3)$ and $E(2n, 4)$ in terms $\zeta(2n)$ and $\zeta(2)\zeta(2n-2)$. We give two formulas for $E(2n, k)$, both valid for arbitrary $k \leq n$, one of which generalizes the Shen-Cai results; by comparing the two we obtain a Bernoulli-number identity. We also give an explicit generating function for the numbers $E(2n, k)$.

¹Supported by a grant from the German Academic Exchange Service (DAAD) during the preparation of this paper. The author also thanks DESY for providing facilities and financial support for travel.

1 Introduction and Statement of Results

For positive integers i_1, \dots, i_k with $i_1 > 1$, we define the multiple zeta value $\zeta(i_1, \dots, i_k)$ by

$$\zeta(i_1, \dots, i_k) = \sum_{n_1 > \dots > n_k \geq 1} \frac{1}{n_1^{i_1} \cdots n_k^{i_k}}. \quad (1)$$

The multiple zeta value (1) is said to have weight $i_1 + \dots + i_k$ and depth k . Many remarkable identities have been proved about these numbers, but in this note we will concentrate on the case where the i_j are even integers. Let $E(2n, k)$ be the sum of all the multiple zeta values of even-integer arguments having weight $2n$ and depth k , i.e.,

$$E(2n, k) = \sum_{\substack{i_1, \dots, i_k \text{ even} \\ i_1 + \dots + i_k = 2n}} \zeta(i_1, \dots, i_k).$$

Of course

$$E(2n, 1) = \zeta(2n) = \frac{(-1)^{n-1} B_{2n} (2\pi)^{2n}}{2(2n)!}, \quad (2)$$

where B_{2n} is the $2n$ th Bernoulli number, by the classical formula of Euler. Euler also studied double zeta values (i.e., multiple zeta values of depth 2) and in his paper [2] gave two identities which read

$$\begin{aligned} \sum_{i=2}^{2n-1} (-1)^i \zeta(i, 2n-i) &= \frac{1}{2} \zeta(2n) \\ \sum_{i=2}^{2n-1} \zeta(i, 2n-i) &= \zeta(2n) \end{aligned}$$

in modern notation. From these it follows that

$$E(2n, 2) = \frac{3}{4} \zeta(2n),$$

though Gangl, Kaneko and Zagier [3] seem to be the first to have pointed it out in print. Recently Shen and Cai [10] proved the formulas

$$E(2n, 3) = \frac{5}{8} \zeta(2n) - \frac{1}{4} \zeta(2) \zeta(2n-2), \quad n \geq 3 \quad (3)$$

$$E(2n, 4) = \frac{35}{64} \zeta(2n) - \frac{5}{16} \zeta(2) \zeta(2n-2), \quad n \geq 4. \quad (4)$$

Identity (3) was also proved by Machide [9] using a different method.

This begs the question whether there is a general formula of this type for $E(2n, k)$. The pattern

$$\frac{3}{4}, \quad \frac{3}{4} \cdot \frac{5}{6} = \frac{5}{8}, \quad \frac{3}{4} \cdot \frac{5}{6} \cdot \frac{7}{8} = \frac{35}{64}$$

of the leading coefficients makes one curious. In fact, the general result is as follows.

Theorem 1. *For $k \leq n$,*

$$E(2n, k) = \frac{1}{2^{2(k-1)}} \binom{2k-1}{k} \zeta(2n) - \sum_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} \frac{1}{2^{2k-3}(2j+1)B_{2j}} \binom{2k-2j-1}{k} \zeta(2j) \zeta(2n-2j).$$

The next two cases after (4) are

$$E(2n, 5) = \frac{63}{128} \zeta(2n) - \frac{21}{64} \zeta(2) \zeta(2n-2) + \frac{3}{64} \zeta(4) \zeta(2n-4)$$

$$E(2n, 6) = \frac{231}{512} \zeta(2n) - \frac{21}{64} \zeta(2) \zeta(2n-2) + \frac{21}{256} \zeta(4) \zeta(2n-4).$$

We prove Theorem 1 in §3 below, using the generating function

$$F(t, s) = 1 + \sum_{n \geq k \geq 1} E(2n, k) t^n s^k.$$

In §2 we establish the following explicit formula.

Theorem 2.

$$F(t, s) = \frac{\sin(\pi \sqrt{1-s} \sqrt{t})}{\sqrt{1-s} \sin(\pi \sqrt{t})}.$$

Our proof uses symmetric functions. We define a homomorphism $\mathfrak{Z} : \text{Sym} \rightarrow \mathbf{R}$, where Sym is the algebra of symmetric functions, and a family $N_{n,k} \in \text{Sym}$ such that \mathfrak{Z} sends $N_{n,k}$ to $E(2n, k)$. We then obtain a formula for the generating functions

$$\mathcal{F}(t, s) = 1 + \sum_{n \geq k \geq 1} N_{n,k} t^n s^k \in \text{Sym}[[t, s]]$$

and apply **3** to get Theorem 2.

From the form of $\mathcal{F}(t, s)$ we show that it satisfies a partial differential equation (Proposition 1 below), which is equivalent to a recurrence for the $N_{n,k}$. From the latter we obtain a formula for the $N_{n,k}$ in terms of complete and elementary symmetric functions, to which **3** can be applied to give the following alternative formula for $E(2n, k)$.

Theorem 3. *For $k \leq n$,*

$$E(2n, k) = \frac{(-1)^{n-k-1} \pi^{2n}}{(2n+1)!} \sum_{i=0}^{n-k} \binom{n-i}{k} \binom{2n+1}{2i} 2(2^{2i-1} - 1) B_{2i}.$$

Note that the sum given by Theorem 3 has $n-k+1$ terms, while that given by Theorem 1 has $\lfloor \frac{k-1}{2} \rfloor + 1$ terms. Yet another explicit formula for $E(2n, k)$ can be obtained by setting $d = 1$ in Theorem 7.1 of Komori, Matsumoto and Tsumura [7]. That formula expresses $E(2n, k)$ as a sum over partitions of k , and it is not immediately clear how it relates to our two formulas.

Comparison of Theorems 1 and 3 establishes the following Bernoulli-number identity.

Theorem 4. *For $k \leq n$,*

$$\begin{aligned} \sum_{i=0}^{\lfloor \frac{k-1}{2} \rfloor} \binom{2k-2i-1}{k} \binom{2n+1}{2i+1} B_{2n-2i} = \\ (-1)^k 2^{2k-2n} \sum_{i=0}^{n-k} \binom{n-i}{k} \binom{2n+1}{2i} (2^{2i-1} - 1) B_{2i}. \end{aligned}$$

It is interesting to contrast this result with the Gessel-Viennot identity (see [1, Theorem 4.2]) valid on the complementary range:

$$\sum_{i=0}^{\lfloor \frac{k-1}{2} \rfloor} \binom{2k-2i-1}{k} \binom{2n+1}{2i+1} B_{2n-2i} = \frac{2n+1}{2} \binom{2k-2n}{k}, \quad k > n. \quad (5)$$

Note that the right-hand side of equation (5) is zero unless $k \geq 2n$.

2 Symmetric Functions

We think of Sym as the subring of $\mathbf{Q}[[x_1, x_2, \dots]]$ consisting of those formal power series of bounded degree that are invariant under permutations of the x_i . A useful reference is the first chapter of Macdonald [8]. We denote the elementary, complete, and power-sum symmetric functions of degree i by e_i , h_i , and p_i respectively. They have associated generating functions

$$\begin{aligned} E(t) &= \sum_{j=0}^{\infty} e_j t^j = \prod_{i=1}^{\infty} (1 + tx_i) \\ H(t) &= \sum_{j=0}^{\infty} h_j t^j = \prod_{i=1}^{\infty} \frac{1}{1 - tx_i} = E(-t)^{-1} \\ P(t) &= \sum_{j=1}^{\infty} p_j t^{j-1} = \sum_{i=1}^{\infty} \frac{x_i}{1 - tx_i} = \frac{H'(t)}{H(t)}. \end{aligned}$$

As explained in [5] and in greater detail in [6], there is a homomorphism $\zeta : \text{Sym}^0 \rightarrow \mathbf{R}$, where Sym^0 is the subalgebra of Sym generated by p_2, p_3, p_4, \dots , such that $\zeta(p_i)$ is the value $\zeta(i)$ of the Riemann zeta function at i , for $i \geq 2$ (in [5, 6] this homomorphism is extended to all of Sym , but we do not need the extension here). Let $\mathcal{D} : \text{Sym} \rightarrow \text{Sym}$ be the degree-doubling map that sends x_i to x_i^2 . Then $\mathcal{D}(\text{Sym}) \subset \text{Sym}^0$, so the composition $\mathfrak{Z} = \zeta \mathcal{D}$ is defined on all of Sym . (Alternatively, we can simply think of \mathfrak{Z} as sending x_i to $1/i^2$: see [8, Ch. I, §2, ex. 21].) Note that $\mathfrak{Z}(p_i) = \zeta(2i) \in \mathbf{R}$. Further, \mathfrak{Z} sends the monomial symmetric function m_{i_1, \dots, i_k} to the symmetrized sum of multiple zeta values

$$\frac{1}{|\text{Iso}(i_1, \dots, i_k)|} \sum_{\sigma \in S_k} \zeta(2i_{\sigma(1)}, 2i_{\sigma(2)}, \dots, 2i_{\sigma(k)}),$$

where S_k is the symmetric group on k letters and $\text{Iso}(i_1, \dots, i_k)$ is the subgroup of S_k that fixes (i_1, \dots, i_k) under the obvious action.

Now let $N_{n,k}$ be the sum of all the monomial symmetric functions corresponding to partitions of n having length k . Of course $N_{n,k} = 0$ unless $k \leq n$, and $N_{k,k} = e_k$. Then \mathfrak{Z} sends $N_{n,k}$ to $E(2n, k)$. Also, if we define (as in the introduction)

$$\mathcal{F}(t, s) = 1 + \sum_{n \geq k \geq 1} N_{n,k} t^n s^k,$$

then \mathfrak{Z} sends $\mathcal{F}(t, s)$ to the generating function $F(t, s)$. We have the following simple description of $\mathcal{F}(t, s)$.

Lemma 1. $\mathcal{F}(t, s) = E((s - 1)t)H(t)$.

Proof. Evidently $\mathcal{F}(t, s)$ has the formal factorization

$$\prod_{i=1}^{\infty} (1 + stx_i + st^2x_i^2 + \dots) = \prod_{i=1}^{\infty} \frac{1 + (s - 1)tx_i}{1 - tx_i} = E((s - 1)t)H(t).$$

□

Proof of Theorem 2. Using the well-known formula for $\zeta(2, 2, \dots, 2)$ [4, Cor. 2.3],

$$\mathfrak{Z}(e_n) = \zeta(\underbrace{2, 2, \dots, 2}_n) = \frac{\pi^{2n}}{(2n + 1)!}. \quad (6)$$

Hence

$$\mathfrak{Z}(E(t)) = \frac{\sinh(\pi\sqrt{t})}{\pi\sqrt{t}},$$

and the image of $H(t) = E(-t)^{-1}$ is

$$\mathfrak{Z}(H(t)) = \frac{\pi\sqrt{-t}}{\sinh(\pi\sqrt{-t})} = \frac{\pi\sqrt{t}}{\sin(\pi\sqrt{t})}.$$

Thus from Lemma 1 $F(t, s) = \mathfrak{Z}(\mathcal{F}(t, s))$ is

$$\mathfrak{Z}(E((s - 1)t)H(t)) = \frac{\sinh(\pi\sqrt{(s - 1)t})}{\pi\sqrt{(s - 1)t}} \frac{\pi\sqrt{t}}{\sin(\pi\sqrt{t})} = \frac{\sin(\pi\sqrt{(1 - s)t})}{\sqrt{1 - s}\sin(\pi\sqrt{t})}.$$

□

Taking limits as $s \rightarrow 1$ in Theorem 2, we obtain

$$F(t, 1) = \frac{\pi\sqrt{t}}{\sin \pi\sqrt{t}}$$

and so, taking the coefficient of t^n , the following result.

Corollary 1. For all $n \geq 1$,

$$\sum_{k=1}^n E(2n, k) = \frac{2(2^{2n-1} - 1)(-1)^{n-1} B_{2n} \pi^{2n}}{(2n)!}.$$

Another consequence of Lemma 1 is the following partial differential equation.

Proposition 1.

$$t \frac{\partial \mathcal{F}}{\partial t}(t, s) + (1 - s) \frac{\partial \mathcal{F}}{\partial s}(t, s) = tP(t)\mathcal{F}(t, s).$$

Proof. From Lemma 1 we have

$$\begin{aligned} \frac{\partial \mathcal{F}}{\partial t}(t, s) &= (s - 1)E'((s - 1)t)H(t) + E((s - 1)t)H'(t) \\ \frac{\partial \mathcal{F}}{\partial s}(t, s) &= tE'((s - 1)t)H(t) \end{aligned}$$

from which the conclusion follows. \square

Now examine the coefficient of $t^n s^k$ in Proposition 1 to get the following.

Proposition 2. For $n \geq k + 1$,

$$p_1 N_{n-1, k} + p_2 N_{n-2, k} + \cdots + p_{n-k} N_{k, k} = (n - k)N_{n, k} + (k + 1)N_{n, k+1}.$$

It is also possible to prove this result directly via a counting argument like that used to prove the lemma of [6, p. 16].

The preceding result allows us to write $N_{n, k}$ explicitly in terms of complete and elementary symmetric functions as follows.

Lemma 2. For $r \geq 0$,

$$N_{k+r, k} = \sum_{i=0}^r (-1)^i \binom{k+i}{i} h_{r-i} e_{k+i}.$$

Proof. We use induction on r , the result being evident for $r = 0$. Proposition 2 gives

$$\sum_{i=1}^{r+1} p_i N_{k+r+1-i, k} = (r + 1)N_{k+r+1, k} + (k + 1)N_{k+r+1, k+1},$$

which after application of the induction hypothesis becomes

$$\begin{aligned} \sum_{i=1}^{r+1} \sum_{j=0}^{r+1-j} (-1)^j p_i \binom{k+j}{j} h_{r+1-i-j} N_{k+j, k+j} = \\ (r+1)N_{k+r+1, k} + (k+1) \sum_{j=0}^r \binom{k+1+j}{j} h_{r-j} N_{k+1+j, k+1+j}. \end{aligned}$$

The latter equation can be rewritten

$$\begin{aligned} \sum_{j=0}^r (-1)^j \binom{k+j}{j} N_{k+j, k+j} \sum_{i=1}^{r+1-j} p_i h_{r+1-i-j} = \\ (r+1)N_{k+r+1, k} - (k+1) \sum_{j=1}^{r+1} (-1)^j \binom{k+j}{j-1} h_{r+1-j} N_{k+j, k+j}. \end{aligned}$$

Now the inner sum on the left-hand side is $(r+1-j)h_{r+1-j}$ by the recurrence relating the complete and power-sum symmetric functions, so we have

$$\begin{aligned} (r+1)N_{k+r+1, k} - (r+1)N_{k, k} h_{r+1} = \\ \sum_{j=1}^{r+1} (-1)^j h_{r+1-j} N_{k+j, k+j} \left((r+1-j) \binom{k+j}{j} + (k+1) \binom{k+j}{j-1} \right), \end{aligned}$$

and the conclusion follows after the observation that $(k+1) \binom{k+j}{j-1} = j \binom{k+j}{j}$. \square

Proof of Theorem 3. Rewrite Lemma 2 in the form

$$N_{n, k} = \sum_{i=0}^{n-k} \binom{n-i}{k} (-1)^{n-k-i} h_i e_{n-i}$$

and apply the homomorphism \mathfrak{Z} , using equation (6) and

$$\mathfrak{Z}(h_i) = \frac{2(2^{2i-1} - 1)(-1)^{i-1} B_{2i} \pi^{2i}}{(2i)!}.$$

\square

3 Proof of Theorems 1 and 4

From the introduction we recall the statement of Theorem 1:

$$E(2n, k) = \frac{1}{2^{2(k-1)}} \binom{2k-1}{k} \zeta(2n) - \sum_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} \frac{1}{2^{2k-3}(2j+1)B_{2j}} \binom{2k-2j-1}{k} \zeta(2j) \zeta(2n-2j).$$

We note that Euler's formula (2) can be used to write the result in the alternative form

$$E(2n, k) = \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-1)^j \pi^{2j} \zeta(2n-2j)}{2^{2k-2j-2}(2j+1)!} \binom{2k-2j-1}{k} \quad (7)$$

which avoids mention of Bernoulli numbers.

We now expand out the generating function $F(t, s)$. We have

$$F(t, s) = \frac{1}{\sqrt{1-s} \sin \pi \sqrt{t}} \sin(\pi \sqrt{t} \sqrt{1-s}) = \frac{\pi \sqrt{t}}{\sin \pi \sqrt{t}} \sum_{j=0}^{\infty} \frac{(-1)^j \pi^{2j} t^j (1-s)^j}{(2j+1)!} = \sum_{k=0}^{\infty} s^k G_k(t),$$

where

$$G_k(t) = (-1)^k \frac{\pi \sqrt{t}}{\sin \pi \sqrt{t}} \sum_{j \geq k} \frac{(-1)^j \pi^{2j} t^j}{(2j+1)!} \binom{j}{k}. \quad (8)$$

Then Theorem 1 is equivalent to the statement that

$$G_k(t) = \sum_{n \geq k} t^n \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-1)^j \pi^{2j} \zeta(2n-2j)}{2^{2k-2j-2}(2j+1)!} \binom{2k-2j-1}{k}$$

for all k . We can write the latter sum as

$$\begin{aligned}
& \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-4\pi^2 t)^j}{2^{2k-2}(2j+1)!} \binom{2k-2j-1}{k} \sum_{n \geq j+1} \zeta(2n-2j)t^{n-j} - \\
& \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-4\pi^2 t)^j}{2^{2k-2}(2j+1)!} \binom{2k-2j-1}{k} \sum_{n=j+1}^{k-1} \zeta(2n-2j)t^{n-j} = \\
& \frac{1}{2}(1 - \pi\sqrt{t} \cot \pi\sqrt{t}) \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-4\pi^2 t)^j}{2^{2k-2}(2j+1)!} \binom{2k-2j-1}{k} - \\
& \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-4\pi^2 t)^j}{2^{2k-2}(2j+1)!} \binom{2k-2j-1}{k} \sum_{n=j+1}^{k-1} \zeta(2n-2j)t^{n-j}, \quad (9)
\end{aligned}$$

where we have used the generating function

$$\frac{1}{2}(1 - \pi\sqrt{t} \cot \pi\sqrt{t}) = \sum_{i=1}^{\infty} \zeta(2i)t^i.$$

Note that the last sum in (9) is a polynomial that cancels exactly those terms in

$$\frac{1}{2}(1 - \pi\sqrt{t} \cot \pi\sqrt{t}) \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-4\pi^2 t)^j}{2^{2k-2}(2j+1)!} \binom{2k-2j-1}{k} \quad (10)$$

of degree less than k . Thus, to prove Theorem 1 it suffices to show that

$$G_k(t) = \text{terms of degree } \geq k \text{ in expression (10)}.$$

From equation (8) it is evident that

$$G_k(t) = \frac{\pi\sqrt{t}}{\sin \pi\sqrt{t}} \cdot \frac{(-t)^k}{k!} \cdot \frac{d^k}{dt^k} \left(\frac{\sin \pi\sqrt{t}}{\pi\sqrt{t}} \right). \quad (11)$$

We use this to obtain an explicit formula for $G_k(t)$.

Lemma 3. For $k \geq 0$,

$$G_k(t) = P_k(\pi^2 t)\pi\sqrt{t} \cot \pi\sqrt{t} + Q_k(\pi^2 t),$$

where P_k, Q_k are polynomials defined by

$$P_k(x) = - \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-4x)^j}{2^{2k-1}(2j+1)!} \binom{2k-2j-1}{k}$$

$$Q_k(x) = \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \frac{(-4x)^j}{2^{2k}(2j)!} \binom{2k-2j}{k}.$$

Proof. In view of equation (11), the conclusion is equivalent to

$$f^{(k)}(t) = (-1)^k k! t^{-k} P_k(\pi^2 t) \cos \pi \sqrt{t} + (-1)^k k! t^{-k} Q_k(\pi^2 t) f(t),$$

where $f(t) = \sin \pi \sqrt{t} / \pi \sqrt{t}$. Differentiating, one sees that the polynomials P_k and Q_k are determined by the recurrence

$$(k+1)P_{k+1}(x) = kP_k(x) - xP'_k(x) - \frac{1}{2}Q_k(x)$$

$$(k+1)Q_{k+1}(x) = \frac{2k+1}{2}Q_k(x) - xQ'_k(x) + \frac{x}{2}P_k(x)$$

together with the initial conditions $P_0(x) = 0$, $Q_0(x) = 1$. The recurrence and initial conditions are satisfied by the explicit formulas above. \square

Proof of Theorem 1. Using Lemma 3, we have

$$G_k(t) = - \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-4\pi^2 t)^j}{2^{2k-1}(2j+1)!} \binom{2k-2j-1}{k} \pi \sqrt{t} \cot \pi \sqrt{t}$$

$$+ \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \frac{(-4\pi^2 t)^j}{2^{2k}(2j)!} \binom{2k-2j}{k} =$$

$$\frac{1}{2}(1 - \pi \sqrt{t} \cot \pi \sqrt{t}) \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-4\pi^2 t)^j}{2^{2k-2}(2j+1)!} \binom{2k-2j-1}{k}$$

+ terms of degree $< k$,

and this completes the proof. \square

Proof of Theorem 4. Using Theorem 1 in the form of equation (7), eliminate $\zeta(2n - 2j)$ using Euler's formula (2) and then compare with Theorem 3 to get

$$\sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{(-1)^{n-1} \pi^{2n} B_{2n-2j}}{2^{2k-2n-1} (2n-2j)! (2j+1)!} \binom{2k-2j-1}{k} = \frac{(-1)^{n-k-1} \pi^{2n}}{(2n+1)!} \sum_{i=0}^{n-k} \binom{n-i}{k} \binom{2n+1}{2i} 2(2^{2i-1} - 1) B_{2i}.$$

Now multiply both sides by $(-1)^{n-1} 2^{2k-2n-1} \pi^{-2n} (2n+1)!$ and rewrite the factorials on the left-hand side as a binomial coefficient. \square

References

- [1] W. Y. C. Chen and L. H. Sun, Extended Zeilberger's algorithm for identities on Bernoulli and Euler polynomials, *J. Number Theory* **129** (2009), 2111-2132.
- [2] L. Euler, Meditationes circa singulare serierum genus, *Novi Comm. Acad. Sci. Petropol.* **20** (1775), 140-186; reprinted in *Opera Omnia*, ser. I, vol. 15, B. G. Teubner, Berlin, 1927, pp. 217-267.
- [3] H. Gangl, M. Kaneko, and D. Zagier, Double zeta values and modular forms, in *Automorphic Forms and Zeta Functions*, S. Böcherer *et. al.* (eds.), World Scientific, Singapore, 2006, pp. 71-106.
- [4] M. E. Hoffman, Multiple harmonic series, *Pacific J. Math.* **152** (1992), 275-290.
- [5] M. E. Hoffman, The algebra of multiple harmonic series, *J. Algebra* **194** (1997), 477-495.
- [6] M. E. Hoffman, A character on the quasi-symmetric functions coming from multiple zeta values, *Electron. J. Combin.* **15** (2008), res. art. 97.
- [7] Y. Komori, K. Matsumoto and H. Tsumura, A study on multiple zeta values from the viewpoint of zeta-functions of root systems, preprint [arXiv:1205.0182](https://arxiv.org/abs/1205.0182).

- [8] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, 2nd ed., Oxford Univ. Press, New York, 1995.
- [9] T. Machide, Extended double shuffle relations and the generating function of triple zeta values of any fixed weight, preprint [arXiv:1204.4085](https://arxiv.org/abs/1204.4085).
- [10] Z. Shen and T. Cai, Some formulas for multiple zeta values, *J. Number Theory* **132** (2012), 314-323.