

Little Higgs Model Limits from LHC – Input for Snowmass 2013

JÜRGEN REUTER^{1a}, MARCO TONINI^{2a}, MAIKEL DE VRIES^{3a}

^aDESY Theory Group, D-22603 Hamburg, Germany

ABSTRACT

The status of the most prominent model implementations of the Little Higgs paradigm, the Littlest Higgs with and without discrete T parity as well as the Simplest Little Higgs are reviewed. For this, we are taking into account a fit to 21 electroweak precision observables from LEP, SLC, Tevatron together with the full 25 fb^{-1} of Higgs data reported from ATLAS and CMS at Moriond 2013. We also – focusing on the Littlest Higgs with T parity – include an outlook on corresponding direct searches at the 8 TeV LHC and their competitiveness with the EW and Higgs data regarding their exclusion potential. This contribution to the Snowmass procedure contains preliminary results of [1] and serves as a guideline which regions in parameter space of Little Higgs models are still compatible for the upcoming LHC runs and future experiments at the energy frontier. For this we propose two different benchmark scenarios for the Littlest Higgs with T parity, one with heavy mirror quarks, one with light ones.

¹juergen.reuter@desy.de

²marco.tonini@desy.de

³maikel.devries@desy.de

1 Introduction: the Little Higgs Paradigm

The first run of LHC at 2 TeV, 7 TeV, and 8 TeV has brought as main results the discovery of a particle compatible with the properties of the Standard Model (SM) Higgs boson as well as with electroweak precision tests (EWPT), and no significant excesses that could be traced to new particles or forces. All of these data have been highly discriminative in the sense of constraining the parameter space of models beyond the SM (BSM), whose focus has been to solve the problem of the quadratic sensitivity of the scalar Higgs mass term in the SM to any kind of UV physics. Scenarios that are weakly coupled at the TeV scale have already been favored over strongly coupled models by EWPT. There is however a class of models where new strong interactions are compatible with EWPT as a mechanism called collective symmetry breaking allows those models to stay weakly coupled at the TeV scale and moves their compositeness scale of strong dynamics up by an order of magnitude. The Higgs (and other scalar bosons) appear as pseudo Nambu-Goldstone bosons of a broken global symmetry very similar to QCD. The problem of quadratic divergences is solved as the Higgs itself is a composite state. Those models are known as Little Higgs models (for reviews cf. [2,3], and references therein). There exists a huge variety of different implementations of the Little Higgs collective symmetry breaking mechanism, but all these models share several common features: an extended scalar sector compared to the SM Higgs, new heavy gauge bosons and new heavy fermions. In this Snowmass white paper we will focus on the three most common models, the Littlest Higgs [4], the Simplest Little Higgs [5] and the Littlest Higgs model with T parity [6]. The Littlest Higgs is the prime example of the product group models, while the Simplest Little Higgs is the corresponding front runner of the simple group models. In the product group models the EW gauge group is embedded into a product group structure, while the scalar sector is in an irreducible representation of the global structure group; for simple group models the EW group is embedded into a larger simple group, while the scalar sector is a reducible representation of the global symmetry structure. Both classes can in principle be discriminated by means of the EW quantum numbers of their additional scalar degrees of freedom [7,8]. T (TeV-scale) parity is a discrete symmetry that can (at least in principle) be introduced for any Little Higgs model. All new particles (with few exceptions) are odd under this parity, which avoids tree-level contributions to the EW precision observables and improves the compatibility with EWPT by roughly an order of magnitude [9,10,11]. Furthermore, it offers the possibility of a Dark Matter candidate with the lightest T-odd particle (LTOP). However now, all new particles must be pair-produced which also reduces the reach for direct searches.

The most important parameter of Little Higgs models is the mass scale f of the new particles, which coincides with the equivalent of the pion decay constant of the underlying non-linear sigma model parameterizing the global symmetry breaking. For the sake of brevity we do not repeat the full formulae defining the three models under consideration, which can be found in [14,1], where also the full set of original references can be found. Other relevant model parameters will be specified in the corresponding sections.

In this Snowmass white paper we will present the limits from the EWPT as well as the Higgs data from the two LHC collaborations ATLAS and CMS from the full 2011/2012 data sets for the Littlest Higgs model (L2H), the Simplest Little Higgs (SLH) and the Littlest Higgs

model with T parity (LHT). We also discuss search results for new particles recasted to the specifications of the LHT model (we concentrate on the LHT as the other models are constrained by Higgs data and EWPT to scales where direct searches at LHC8 are not feasible). Note that some of the direct search results are still partially preliminary and will be presented in [1]. Sec. 2 discusses the bounds from EWPT, Sec. 3 the constraints from the LHC Higgs data.

2 Electroweak Precision Tests (EWPT)

For the EWPT, we calculate the EW precision observables within the three different implementations of the Little Higgs models. In the L2H they arise from the tree-level exchange of the heavy gauge bosons as well as from the vacuum expectation value (vev) v' of the Higgs triplet fields. Loops from heavy quarks to the EW gauge boson self energies are negligible. All of the observables can be expressed through the four variables f , $c \equiv \cos \theta$, $c' \equiv \cos \theta'$, and $x \equiv 4v'f/v^2$, i.e. the L2H scale, the mixing angles in the non-Abelian and Abelian gauge boson sector as well as the dimensionless ratio of the triplet and doublet vev. For the SLH model, the dominant contributions come from the Z' boson corrections as well as the $Z - Z'$ mixing, and the parameter besides f that goes in is the mixing angle between the two Goldstone boson multiplets, $t_\beta := \tan \beta$. Introducing T parity removes all tree-level contributions in the Littlest Higgs to the EW precision observables (except maybe the contribution of the T -even top quark to top quark operators). To the oblique parameters, only the T -even top contributes at 1-loop, as the T -odd top is an EW singlet. An additional contribution from the mirror fermions and a logarithmic correction to the oblique parameters from the modified Higgs couplings to the EW gauge bosons are included, too. There are negligible pieces from the heavy gauge bosons to the EW precision observables. The corrections in the flavor sector in all three models can actually be expressed by corrections to the EW charged and neutral current couplings from their SM values as an expansion in v/f . The explicit expressions of the corrections can be found in [14].

For the calculation of the 21 EW precision parameters (low-energy data and on the Z pole) we use the values given in [19] taking the observed Higgs mass of 125 GeV into account. The values of the EW precision parameters are summarized in [14]. For simplicity, we assume no correlations among the EW precision observables and include them in a χ^2 fit defined just by quadratic deviations from the measured values divided by the (squared) experimental uncertainties.

From the EWPT, one gets the following exclusion limits at 95% CL on the Little Higgs scale for the corresponding models: $f \lesssim 5100$ GeV for the L2H, $f \lesssim 3700$ GeV for the SLH, and $f \lesssim 405$ GeV for the LHT. Exclusion plots can be found in [1], we show them here only in combination with the Higgs data in the next section.

3 Constraints from Higgs data

After the LHC discovery of the Higgs particle compatible with EWPT, the precision of the data (besides systematic uncertainties) grows with the square root of the integrated luminosity and is starting being competitive with the discriminative power of the EWPT itself [12,13].




Figure 1: *Exclusion limits at 95% CL (light-blue [light]) and 99% CL (dark-blue [dark]) from EWPT and Higgs data combined as a function of the Little Higgs model scale f . The thick black lines represent contours of required fine-tuning. Left: Littlest Higgs model (L2H), showing the exclusion region for the mixing angle $c \equiv \cos \theta$ in the gauge sector; right: Simplest Little Higgs (SLH) exclusion region in $\tan \beta$, the mixing angle of the two non-linear sigma model fields. The red [very dark] region gives no viable EW symmetry breaking.*

The two collaborations, ATLAS and CMS, express their results for the Higgs channels in terms of the signal strength modifier μ , which for a given Higgs mass, m_h , is the ratio between the observed cross section times branching ratio folded with the experimental efficiency normalized to the corresponding values of the SM. For the χ^2 analysis of the compatibility of the Higgs data with Little Higgs models we take all published experimental data into account, coming from the vector boson channels $\gamma\gamma$ (all different experimental categories), ZZ , WW , as well as the two fermionic channels bb and $\tau\tau$. We do not repeat the collection of all the values here, the details can be found in [14,1]. The 95% and 99% CL regions are then defined by the cumulative distribution function for an appropriate number of degrees of freedom (DOF) within our χ^2 measure. For all channels we take the latest public 7 and 8 TeV data of the 2011 and 2012 LHC runs. Our results are shown in Fig. 1 for the L2H (left) and the SLH model (right) in the (f, c) and (f, t_β) plane, respectively as well as for the LHT in Fig. 2 in the (f, R) plane. The black lines are contour lines of the fine tuning variable $1/\Delta$, with $\Delta = |\delta\mu^2|/\mu_{\text{obs}}^2$, $\mu_{\text{obs}}^2 = \frac{m_h^2}{2}$. Here, μ is the Higgs mass parameter and $\delta\mu$ its leading radiative correction. This means, a worse fine-tuning results in a smaller value of $1/\Delta$, as a certain parameter value has to be tuned to a much smaller level to achieve the experimentally observed Higgs mass. The 95% and 99% CL exclusion regions combining the Higgs and EW are shown in light blue [light] and dark blue




Figure 2: *Exclusion regions for the LHT model from EWPT and Higgs data combined, for the ratio $R = \lambda_1/\lambda_2$ of two Yukawa couplings in the T -parity doubled fermion sector. Color code as in Fig. 1*

[dark], respectively. Due to the strong contributions to the EW precision observables, in the case of the L2H and the SLH the EWPT dominate the exclusion region, and the inclusion of Higgs data compatible with the SM (in the large- f region) just reduces the χ^2/DOF . This leads to a weaker exclusion from the Higgs+EWPT combination as from the EWPT data alone: the 95% CL exclusion for the L2H is $f \lesssim 3700$ GeV, while it is $f \lesssim 3300$ GeV for the SLH. For the LHT, the general limit (independent of R [14]) improves to a 95% CL exclusion of $f \lesssim 694$ GeV, while the constraint from Higgs data alone would be $f \lesssim 607$ GeV at 95% CL. This comes mainly from the fact that the Higgs exclusion for the LHT is independent from R and does not suffer from the dip around $R \approx 1$ coming from a cancellation of LHT contributions to the T parameter.

4 Limits from Direct Searches

In this section, we take into account different searches published by the LHC experiments that have been optimized especially for low-energy supersymmetry. As the bounds from EWPT (and also Higgs data) for the L2H and SLH already push up the Little Higgs scale into the multi-TeV region, this makes direct discoveries of new particles at 8 TeV impossible and even at 14 TeV very difficult. On the other hand, most of the direct searches for the L2H and SLH could be covered with the Z' , W' searches as well as the searches for heavy vector-like quarks. For these reasons, we focus on the LHT for the direct searches.




Figure 3: *LHT benchmark scenarios, fixing $R = 1.0$ in order to minimize EWPT: (1) heavy mirror quarks ($k = 1.5$) or rather light ones ($k = 0.4$).*

The LHT model can be parameterized in principle by two parameters beyond those of the SM and the Little Higgs scale f : the ratio of Yukawa couplings in the top sector, $R := \lambda_1/\lambda_2$, as well as a parameter from the sector of mirror fermions necessary to implement T parity: a Yukawa coupling k that sets the scale of the mirror fermions. Generically, the heavy photon A_H is the lightest T -odd particle (LTOP), while the heavy vectors W_H and Z_H are degenerate up to corrections of order v^2/f^2 as demanded by EWPT. For our studies, the scalar triplet can be ignored. The top partners T_+, T_- are always heavier than all bosons. The parameter k determines the mass of the quark partners: for $k \gtrsim 0.46$ the quark partners are heavier than all gauge bosons, opening up the decay $q_H \rightarrow V_H q$. For lower values of k there is only the decay channel $q_H \rightarrow A_H q$. In the region $0.1 \lesssim k \lesssim 0.2$ the q_H - A_H mass difference is rather small, making the spectrum compressed. In Fig. 3 we propose two benchmark scenarios for the LHT, fixing $R \equiv 1.0$, thereby minimizing the contributions to the EW precision observables. The two different cases of decay patterns for the quark partners are reflected in the two different choices for k , $k = 1.5$ or $k = 0.4$. Tables with the dominant decay modes for the different states in the two benchmark cases can be found in Ref. [1].

In order to discuss direct search limits, we briefly discuss the major production modes at the LHC. The T -even top partner T_+ is the only new state to be singly produced, while all other particles have to be pair produced, highly reducing the available phase space with increasing masses. Due to the PDF enhancement, the pair production of the quark partners q_H could be significant, especially if their masses are not too large. Their production is mainly dominated by QCD processes, but also EW processes with a heavy W_H in the t -channel significantly contribute. All details about the cross sections and listings of the relevant signatures can be found in [1].

Before we discuss direct searches at the LHC, there are constraints coming from integrating

out the T -odd quark generating effective four-fermion operators, $\mathcal{O}_{4-f} = -\frac{k^2}{128\pi^2 f^2} \bar{\psi}_L \gamma^\mu \psi_L \bar{\psi}'_L \gamma_\mu \psi'_L + \mathcal{O}\left(\frac{g}{k}\right)$, under the assumption that k is diagonal and flavor-independent. ψ and ψ' are different SM fermions. Experimental bounds on four-fermion interactions provide an upper bound on the T -odd fermion masses, translating into an upper bound on k . At LHC, angular distributions and rates in dijet events lead to bounds of the order $\Lambda = 15$ TeV for (for constructive interference with the SM). The most stringent bounds, however, are still from LEP, constraining the coefficient of the operator $eedd$ operator [11,19]. This leads to the upper bound $k^2 \lesssim 0.367 \pi^3 (f [\text{TeV}])^2$. This bound is shown in the total exclusion plot in Fig. 4 as the triangular-shaped purple [dark] region in the upper half. We do expect LHC to improve these bounds, since 8 TeV analyses for operator bounds of this form still need to be published. Moreover many promising results from the 14 TeV run can be expected in this area, probably driving limits up to $\mathcal{O}(30 \text{ TeV})$.

Next, we will consider all searches from the 8 TeV LHC run relevant to the Littlest Higgs with T -parity. For each analysis, we identify the final states and the contributing production modes in the LHT model. Typically, the production cross section times branching ratio ($\sigma \times \text{Br}$) depends on the scale f and either the parameter k or R . Therefore exclusion limits can either be presented as contours in the (f, k) or the (f, R) plane. For this Snowmass white paper, we summarize the exclusion limits in (f, k) plane in Fig. 4. This allows to set a lower limit on the scale f . In the following, we briefly discuss the different searches. More details can be found in [1].

To obtain the exclusion limits from recasting the LHC analyses (mostly for SUSY searches) we generated signal events for the LHT model with MadGraph [15] and WHIZARD [16,17]. The events are then processed with the fast detector simulation Delphes 3.0 [18] to simulate either the ATLAS or CMS detector in order to evaluate the cut efficiencies of the different analyses applied to the assumed LHT signal. With these events, we have been recasting three different classes of searches, (1) monojets plus missing transverse energy (MET), (2) jets plus MET, and (3) jets, leptons and MET.

1. Both ATLAS and CMS have presented monojet searches with 8 TeV data for final states containing a single energetic jet and missing transverse energy [20,21]. They are quite similar, requiring a single energetic jet and at most two further jets with $p_T > 30$ GeV. Furthermore, a lepton veto is imposed and the presence of significant amounts of missing transverse energy is required in both analyses. For more details on the signal regions, cf. [1]. Additional suppression of QCD dijet background is done differently: ATLAS requires the azimuthal separation between the direction of missing E_T and the second leading jet (if present) to be greater than 0.5, whereas CMS only keeps two jet events if the azimuthal separation between the jets is less than 2.5. In the absence of any deviation from the SM, both experiments quote 95% CL upper bounds on the signal cross section times efficiency for their corresponding signal regions. Both monojet searches are suitable for final state topologies containing one or two hard jets and missing transverse energy. Hence both LHT production modes $pp \rightarrow q_H q_H$ and $pp \rightarrow q_H A_H$ may contribute, provided the heavy quark partner decays to a quark and a heavy photon $q_H \rightarrow A_H q$. Therefore these searches are mostly sensitive in the parameter regions where quark partners q_H are lighter

than the gauge bosons W_H and Z_H , i.e. for lower values of k (fixed f), such that they entirely decay as $q_H \rightarrow A_H q$. Fig. 4 shows the excluded regions recasting both ATLAS and CMS monojet + MET analysis as a red band at the lower end.

2. In this second category, all searches with at least two jets, missing transverse energy and no leptons in the final state are investigated. Numerous searches interpreted in terms of SUSY have been presented for the 8 TeV data by ATLAS [22,23] and CMS [24]. The first ATLAS search is optimized for squarks and gluinos, the second one for stops, whereas the CMS search looks more generically at squarks, sbottoms and gluinos. In the LHT scenario these searches can be applied to pair production of heavy gauge bosons and heavy quarks or associated productions like $V_H q_H$. In those cases, the heavy quarks decay to heavy vectors (decaying to SM gauge bosons, all hadronic final states, and MET from the A_H) and quarks. Also covered are possible LHT production modes like pair production of the




Figure 4: *Exclusion limits from direct searches at LHC8 displayed in the (f, k) plane. The different categories comprise limits from operator bounds and searches from monojets, jets and leptons plus jets. The contour lines show the mass of the heavy quark partners.*

T-odd top t_H with subsequent stop-like decay, or two heavy quarks where at least one decays like $q_H \rightarrow Z_H q$, giving the required two b-jets via $Z_H \rightarrow H A_H \rightarrow bbA_H$. Again, each of the above searches provide 95% CL limits on the cross sections times efficiencies in the absence of any signal. Also, the CMS analysis [24] can be recast as a search for pair production of heavy gauge bosons and quarks as well as the associated productions $V_H q_H$ with completely hadronic final states. A detailed list of the experimental signal regions, the SM background suppression, the cut flows and the efficiencies and the recast procedure can be found in [1]. The results from recasting the ATLAS and CMS searches from jets + MET described above for the LHT model are shown as light blue exclusion region in Fig. 4.

3. Here, all searches involving tagged leptons, at least two jets and missing transverse energy are collected. Some of the jets may be b-tagged. For this purpose, we consider the following searches by ATLAS and CMS resulting in corresponding constraints for the LHT model: [25,26,27]. These searches match also LHT topologies like production of two heavy quarks, which then decay to heavy gauge bosons $q_H \rightarrow W_H q$ or $\rightarrow Z_H q$ with one or both W s decaying leptonically. b -jets may arise from the Higgs decay in the $Z_H \rightarrow H A_H$ decay chain, or from the decay chain of the T-odd top partner $t_H \rightarrow t A_H$. Another signature covered is pair production of same charge heavy quarks $pp \rightarrow q_H q_H$ with subsequent decays into gauge boson partners W_H with now leptonic decays for the W s. From the searches with leptons, jets and MET described in this paragraph, 95% CL exclusion limits in the (f, k) plane can be extracted. The results from the recast are presented in Fig. 4 as yellow region in the middle of the left part of the plot. From our recast, we conclude that searches for both a single and two leptons perform equally, as long as no b -jets in the final states are required.

Finally, we combine all direct searches in Fig. 4, derive a combined exclusion on the scale f from it, and compare with the bounds from EWPT and Higgs data. We get an exclusion at 95% CL from direct searches on the scale f of $f \lesssim 620$ GeV.

5 Summary and Conclusion

We summarized constraints from EW precision test (EWPT), the Higgs data from the ATLAS and CMS analyses for the three most economic implementations of Little Higgs models, the Littlest Higgs (L2H), the Simplest Little Higgs (SLH) and the Littlest Higgs with T parity. For the latter, we also recasted direct searches from the LHC experiments optimized for SUSY searches. While the constraints from EWPT as well as from the Higgs data for L2H and SLH are in the regime above 3-3.5 TeV, LHT is in much better shape: from EWPT we get a 95% CL exclusion of $f \lesssim 405$ GeV, from Higgs data of $f \lesssim 607$ GeV, from Higgs data combined with EWPT of $f \lesssim 694$ GeV. Direct searches are in the same ballpark but not completely competitive yet, resulting in $f \lesssim 620$ GeV from monojet + MET, jets + MET, leptons, jets + MET, as well as bounds from contact interaction searches.

As we have not yet completed our studies for the deadline of the Snowmass process, this study is just a proceedings-style summary. Specifically, we do not give any strategies for the optimization of the cut flows for the direct search analysis which will be found in [1]. This would enable the LHC experiments to improve on their limits for the Little Higgs parameter spaces and would also allow to give a more reliable estimate for the prospects for 14 TeV, the high-luminosity LHC upgrade (HL-LHC), or possible future energy upgrades to 33 or 100 TeV. Such estimates do rely anyhow on a possible calibration of the experiments. Furthermore, prospects for a future ILC are also not included in this note (however, cf. [29]).

Acknowledgements

The authors like to thank Diptimoy Gosh, Paolo Gunnellini, Maxim Perelstein, and Andreas Weiler for valuable remarks and discussions. M.T. and M.deV have been partially supported by the Deutsche Forschungsgemeinschaft within the Collaborative Research Center SFB 676 "Particles, Strings, Early Universe".

References

- [1] J. Reuter, M. Tonini, M. de Vries, *Status and Prospect of the Littlest Higgs with T parity*, DESY 13-123, in preparation.
- [2] M. Schmaltz and D. Tucker-Smith, *Little Higgs review*, Ann. Rev. Nucl. Part. Sci. **55**, 229 (2005) [hep-ph/0502182].
- [3] M. Perelstein, *Little Higgs models and their phenomenology*, Prog. Part. Nucl. Phys. **58**, 247 (2007) [hep-ph/0512128].
- [4] N. Arkani-Hamed, A. G. Cohen, E. Katz and A. E. Nelson, *The Littlest Higgs*, JHEP **0207**, 034 (2002) [hep-ph/0206021].
- [5] M. Schmaltz, *The Simplest little Higgs*, JHEP **0408**, 056 (2004) [hep-ph/0407143].
- [6] H. -C. Cheng and I. Low, *TeV symmetry and the little hierarchy problem*, JHEP **0309**, 051 (2003) [hep-ph/0308199].
- [7] W. Kilian, D. Rainwater and J. Reuter, *Pseudo-axions in little Higgs models*, Phys. Rev. D **71**, 015008 (2005) [hep-ph/0411213].
- [8] W. Kilian, D. Rainwater and J. Reuter, *Distinguishing little-Higgs product and simple group models at the LHC and ILC*, Phys. Rev. D **74**, 095003 (2006) [Erratum-ibid. D **74**, 099905 (2006)] [hep-ph/0609119].
- [9] C. Csaki, J. Hubisz, G. D. Kribs, P. Meade and J. Terning, *Big corrections from a little Higgs*, Phys. Rev. D **67**, 115002 (2003) [hep-ph/0211124].

- [10] W. Kilian and J. Reuter, *The Low-energy structure of little Higgs models*, Phys. Rev. D **70**, 015004 (2004) [hep-ph/0311095].
- [11] J. Hubisz, P. Meade, A. Noble and M. Perelstein, *Electroweak precision constraints on the lightest Higgs model with T parity*, JHEP **0601**, 135 (2006) [hep-ph/0506042].
- [12] [ATLAS Collaboration], *Combined coupling measurements of the Higgs-like boson with the ATLAS detector using up to 25 fb^{-1} of proton-proton collision data*, ATLAS-CONF-2013-034.
- [13] [CMS Collaboration], *Combination of standard model Higgs boson searches and measurements of the properties of the new boson with a mass near 125 GeV* , CMS-PAS-HIG-13-005.
- [14] J. Reuter and M. Tonini, *Can the 125 GeV Higgs be the Little Higgs?*, JHEP **1302**, 077 (2013) [arXiv:1212.5930 [hep-ph]].
- [15] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer and T. Stelzer, *MadGraph 5 : Going Beyond*, JHEP **1106**, 128 (2011) [arXiv:1106.0522 [hep-ph]].
- [16] W. Kilian, T. Ohl and J. Reuter, *WHIZARD: Simulating Multi-Particle Processes at LHC and ILC*, Eur. Phys. J. C **71**, 1742 (2011) [arXiv:0708.4233 [hep-ph]].
- [17] M. Moretti, T. Ohl and J. Reuter, *O'Mega: An Optimizing matrix element generator*, In *2nd ECFA/DESY Study 1998-2001* 1981-2009 [hep-ph/0102195].
- [18] S. Ovin, X. Rouby and V. Lemaître, *DELPHES, a framework for fast simulation of a generic collider experiment*, arXiv:0903.2225 [hep-ph].
- [19] J. Beringer *et al.* [Particle Data Group Collaboration], *Review of Particle Physics (RPP)*, Phys. Rev. D **86**, 010001 (2012).
- [20] [ATLAS Collaboration], *Search for New Phenomena in Monojet plus Missing Transverse Momentum Final States using 10 fb^{-1} of pp Collisions at $\sqrt{s}=8 \text{ TeV}$ with the ATLAS detector at the LHC*, ATLAS-CONF-2012-147.
- [21] [CMS Collaboration], *Search for new physics in monojet events in pp collisions at $\sqrt{s}=8 \text{ TeV}$* , CMS-PAS-EXO-12-048.
- [22] [ATLAS Collaboration], *Search for squarks and gluinos with the ATLAS detector using final states with jets and missing transverse momentum and 5.8 fb^{-1} of $\sqrt{s}=8 \text{ TeV}$ proton-proton collision data*, ATLAS-CONF-2012-109.
- [23] [ATLAS Collaboration], *Search for direct production of the top squark in the all-hadronic $t\bar{t}$ + $e\text{miss}$ final state in 21 fb^{-1} of p - p collisions at $\sqrt{s}=8 \text{ TeV}$ with the ATLAS detector*, ATLAS-CONF-2013-024.

- [24] S. Chatrchyan *et al.* [CMS Collaboration], *Search for supersymmetry in hadronic final states with missing transverse energy using the variables α_T and b-quark multiplicity in pp collisions at $\sqrt{s} = 8$ TeV*, arXiv:1303.2985 [hep-ex].
- [25] [ATLAS Collaboration], *Search for supersymmetry at $\sqrt{s} = 8$ TeV in final states with jets, missing transverse momentum and one isolated lepton*, ATLAS-CONF-2012-104.
- [26] [ATLAS Collaboration], *Search for direct top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 8, 13$ TeV pp collisions using 21 fb^{-1} of ATLAS data*, ATLAS-CONF-2013-037.
- [27] [ATLAS Collaboration], *Search for strongly produced superpartners in final states with two same sign leptons with the ATLAS detector using 21 fb^{-1} of proton-proton collisions at $\sqrt{s} = 8$ TeV*, ATLAS-CONF-2013-007.
- [28] [ATLAS Collaboration], *Search for supersymmetric top-quark partner in final states with two leptons in $\sqrt{s} = 8$ TeV pp collisions using 13 fb^{-1} of ATLAS data*, ATLAS-CONF-2012-167 .
- [29] H. Baer, T. Barklow, K. Fujii, Y. Gao, A. Hoang, S. Kanemura, J. List and H. E. Logan *et al.*, *The International Linear Collider Technical Design Report - Volume 2: Physics*, arXiv:1306.6352 [hep-ph].