

**ELECTROWEAK VACUUM STABILITY
AND
FINITE QUADRATIC RADIATIVE CORRECTIONS**

Isabella Masina^{a,b}, Germano Nardini^c and Mariano Quiros^d

^a *Dip. di Fisica e Scienze della Terra, Università di Ferrara and INFN Sez. di Ferrara,
Via Saragat 1, I-44100 Ferrara, Italy*

^b *CP³-Origins and DIAS, Southern Denmark University,
Campusvej 55, DK-5230 Odense M, Denmark*

^c *Deutsches Elektronen Synchrotron, Notkestrasse 85, D-22603 Hamburg, Germany*

^d *Institució Catalana de Recerca i Estudis Avançats (ICREA) and
IFAE-UAB 08193 Bellaterra, Barcelona, Spain*

Abstract

If the Standard Model (SM) is an effective theory, as currently believed, it is valid up to some energy scale Λ to which the Higgs vacuum expectation value is sensitive throughout radiative quadratic terms. The latter ones destabilize the electroweak vacuum and generate the SM hierarchy problem. For a given perturbative Ultra-violet (UV) completion, the SM cutoff can be computed in terms of fundamental parameters. If the UV mass spectrum involves several scales the cutoff is not unique and each SM sector has its own UV cutoff Λ_i . We have performed this calculation assuming the Minimal Supersymmetric Standard Model (MSSM) is the SM UV completion. As a result, from the SM point of view, the quadratic corrections to the Higgs mass are equivalent to finite threshold contributions. For the measured values of the top quark and Higgs masses, and depending on the values of the different cutoffs Λ_i , these contributions can cancel even at renormalization scales as low as multi-TeV, unlike the case of a single cutoff where the cancellation only occurs at Planckian energies, a result originally obtained by Veltman. From the MSSM point of view, the requirement of stability of the electroweak minimum under radiative corrections is incorporated into the matching conditions and provides an extra constraint on the Focus Point solution to the little hierarchy problem in the MSSM. These matching conditions can be employed for precise calculations of the Higgs sector in scenarios with heavy supersymmetric fields.

I. INTRODUCTION

In the Standard Model (SM) [1] as an effective field theory with a physical cutoff Λ , the Higgs mass parameter is corrected by (uncalculable) quadratic divergences that can destabilize the electroweak vacuum. This fact is usually associated with the SM hierarchy problem [2]. In the presence of a perturbative Ultraviolet (UV) completion (beyond the TeV scale) with heavy fields coupled to the Higgs sector, the quadratic divergences appear as finite threshold effects, which can therefore be reliably computed in perturbation theory, after the heavy states are integrated out at the matching scale between the Low-Energy (LE) and the UV High-Energy (HE) effective theories. So, if the UV completion of the SM is known and it is perturbative, the hierarchy problem is entirely due to calculable finite effects and can be fully quantified. If the UV completion is non-perturbative, as it happens in the case where the Higgs is composite, the calculation cannot rely on perturbation theory, but the presence of a new scale, even if it is dynamically generated, makes it possible to estimate the size of the threshold corrections to the Higgs mass [3]. Here we will consider the former case where the UV completion is perturbative.

The absence of any departure from the SM predictions in current experimental data at the LHC is pointing towards the existence of new physics at least in the multi-TeV region, by which the naturalness problem is becoming more acute. This in turn is hinting at less conventional solutions to the hierarchy problem, as e.g. hypothetical solutions provided by the theory which breaks supersymmetry at the (high) scale where supersymmetry breaking is transmitted from the hidden to the observable sector. It is therefore important to compute the large radiative contributions to the hierarchy problem in order to settle the required conditions at the high scale for the stability of the electroweak vacuum in the effective theory (the SM) below the matching scale.

In this paper we will consider the SM as the LE effective theory of the Minimal Supersymmetric Standard Model (MSSM), matching the two theories at the decoupling scale \mathcal{Q}_m where the supersymmetric partners are integrated out. We assume the MSSM is valid up to scales of the order of the Planck scale M_P , where it can be understood as e.g. the flat limit ($M_P \rightarrow \infty$) of $N = 1$ supergravity [4], which should eventually be in turn UV completed by some more fundamental (superstring) theory. The hope is that the fundamental theory could provide the requirements for solving the SM hierarchy problem, under the form of some HE parameter relations. For that reason, in this paper we are trying to fix the required conditions which could lead to stability of the electroweak minimum, but by no means are we trying to claim any solution to the hierarchy problem, nor even a precise quantification of the fine-tuning.

Threshold effects when matching the SM with the MSSM have been extensively studied for dimensionless parameters, as e.g. the SM Yukawa and quartic couplings, thus fixing the physical Higgs and fermion masses [5]. For dimensional parameters, as the Higgs

mass parameter, the thresholds have not been systematically considered ¹. However, the hierarchy problem precisely resides in those dimensional parameters, as solving the equations of minimum providing the electroweak vacuum expectation value (VEV) of the Higgs field requires a certain amount of fine-tuning which quantifies the hierarchy problem ².

It is thus worth improving our knowledge on the dimensional parameter thresholds. To this aim, in this paper we analyze the effects on the SM Higgs mass parameter from the decoupling of the heavy MSSM fields. This will result in precise relations to be satisfied at the matching scale \mathcal{Q}_m , in order to have the stability of the electroweak minimum. Technically, we perform the matching in the one-loop RG-improved approximation, as going beyond one-loop should not add qualitative complications or dominant contributions. Similarly we perform the matching procedure in the symmetric phase for the MSSM, so our results should be affected by tiny corrections of $\mathcal{O}(v^2/\mathcal{Q}_m^2)$.

The outline of the paper goes as follows. In section II we present some general ideas about the decoupling using the scale invariance of the effective potential in the one-loop RG-improved approximation. We show that in the considered approximation the decoupling scale \mathcal{Q}_m is arbitrary, although in view of minimizing (unconsidered) higher loop corrections it is convenient to take it of the order of magnitude of the masses of the decoupled fields. Simple toy models to illustrate the general matching procedure are presented in section III, where we also stress the role, for scale invariance, of the anomalous dimensions of scalar fields (included in the wave-function radiative corrections) which is an ingredient alien to the effective potential, constructed at zero external momentum. The case of the MSSM is reviewed in section IV and the detailed matching between SM and MSSM Higgs mass parameters is performed in section V. The threshold effects induced in the effective theory are computed in section VI. In particular we show that for the MSSM scenario with degenerate soft breaking masses the finite correction to the SM Higgs mass parameter precisely reproduces the result obtained by Veltman (in the context of dimensional regularization in two dimensions) [8] if the SM cutoff is identified with the common mass of the degenerate and heavy supersymmetric partners. Instead, for the more general scenario with non degenerate heavy masses, the effective theory can be often interpreted as a SM with different cutoffs for each (quarks, gauge bosons, ...) sector. In such a case the finite correction to the SM Higgs mass parameter consists in a generalized Veltman result with deformations that can be negligible depending on the hierarchy of the spectrum. In section VII we express the HE parameters evaluated at the decoupling scale \mathcal{Q}_m in terms of their values at the scale \mathcal{M} where the supersymmetry breaking is transmitted to the observable sector, and we constrain these values to be

¹ For a previous analysis in the MSSM in the broken electroweak symmetric phase, see Ref. [6].

² The instability under radiative corrections coming from new heavy physics, which is the subject of the present paper, should not be confused with the instability driven by the Renormalization Group (RG) running of the quartic Higgs coupling towards negative values [7].

compatible with sensible matching conditions. We then quantify the thresholds effects and their impact on the Higgs sector. We focus on the parameter region corresponding to the Focus Point (FP) solutions. The regions where the electroweak stability is not spoiled by the finite corrections in the matching conditions can be understood as a generalized FP solutions which include thresholds effects. Our conclusions are presented in section VIII, as well as a discussion on the scale, gauge and renormalization scheme dependence of our results. Finally, technical details of the calculation of the radiative corrections to the SM Higgs mass parameter stemming from the different supersymmetric sectors are presented in section A. A nice check of the consistency of our calculation is the explicit proof of the one-loop scale invariance of our results, and this is presented in section B.

II. GENERAL IDEAS ABOUT THE DECOUPLING

Before putting forward the explicit relation of the effective potential V in the LE and HE regions, let us review some general ideas about the effective potential [9]. The effective potential improved by the RG depends, on top of the background value of fields ϕ_i , on a number of running parameters λ_I (they include dimensionless couplings as well as dimensionful parameters) and on the renormalization scale \mathcal{Q} , in such a way that the equation

$$\left[\mathcal{Q} \frac{\partial}{\partial \mathcal{Q}} + \beta_I \frac{\partial}{\partial \lambda_I} + \gamma_i \phi_i \frac{\partial}{\partial \phi_i} \right] V = 0 \quad (1)$$

is fulfilled. This equation, where γ_i are the anomalous dimensions of the fields ϕ_i , and β_I the β -functions of the parameters λ_I , highlights the renormalization-scale independence of the effective potential. The general solution of Eq. (1) reads as

$$V = V(\mathcal{Q}(t), \lambda_I(t), \phi_i(t)) \quad (2)$$

where

$$\mathcal{Q}(t) = \mathcal{Q} \exp(t), \quad \beta_I(\lambda_I(t)) = \frac{d\lambda_I(t)}{dt}, \quad \gamma_i(\lambda_I(t)) = \frac{d \log(\phi_i(t)/\phi_i)}{dt}, \quad (3)$$

together with the boundary conditions $\lambda_I(0) \equiv \lambda_I$, $\phi_i(0) \equiv \phi_i$.

In practice the scale independence of the effective potential (2) holds up to the level of perturbation theory where the potential is computed. In particular if we make a loop expansion of the operator V as

$$V = \sum_{\ell \geq 0} V^{(\ell)}, \quad (4)$$

the RG-improved potential $V^{(0)}$ has a very strong scale dependence. This dependence is reduced by considering $V^{(0)} + V^{(1)}$, where $V^{(1)}$ includes the terms that correspond to the field redefinitions $\phi_i(t) \equiv (1 + Z_i^{(1)}(t))\phi_i$, and the one-loop RG-improved Coleman-Weinberg contribution $\Delta V^{(1)}$. In fact, in $V^{(0)} + V^{(1)}$ the whole one-loop scale dependence cancels out and different choices of Q only affect higher order corrections. Whereas the

explicit expression of $Z_i(t)$ depends on the specific model, the contribution $\Delta V^{(1)}$ can be generically written as [10]³

$$\Delta V^{(1)} = \frac{1}{64\pi^2} STr \mathcal{M}^4(\phi_i) \left[\log \frac{\mathcal{M}^2(\phi_i)}{\mathcal{Q}^2} - \mathcal{C} \right] \quad (5)$$

where STr includes the number of degrees of freedom of the different mass eigenstates as well as a negative sign for fermions.

The electroweak-breaking condition, described as the solution to the equations of minimum

$$\frac{\partial V}{\partial \phi_i} = 0, \quad (6)$$

is also scale independent. Such condition can thus be deduced from the potential V at any scale \mathcal{Q} and any loop order. However since one is only able to compute $\Delta V^{(\ell)}$ and $Z_i^{(\ell)}$ with $\ell \leq n$ (i.e. up to some perturbative order n), and the minimization condition should eventually be related to electroweak observables, in practical cases it is advisable to minimize $\sum_{\ell \leq n} V^{(\ell)}$ (with scale dependence at $n+1$ loop order) at the electroweak scale $\mathcal{Q} = \mathcal{Q}_{EW}$. Employing this choice of renormalization scale is subtle when also heavy fields are involved, as we describe now.

We consider a HE theory with light and heavy fields. Light fields have electroweak-breaking and/or invariant masses of order (or below) \mathcal{Q}_{EW} , whereas heavy fields have masses $M \gg \mathcal{Q}_{EW}$. To any fixed order in the loop expansion heavy fields can induce large logarithms in the minimization condition evaluated at $\mathcal{Q} = \mathcal{Q}_{EW}$ [11]. For this reason, the minimization should still be performed at $\mathcal{Q} = \mathcal{Q}_{EW}$, but in the LE effective theory where the heavy fields have been decoupled.

As we are considering mass-independent renormalization schemes the decoupling of heavy fields has to be performed at some scale \mathcal{Q}_m . In such a case the effective description at $\mathcal{Q} \ll \mathcal{Q}_m$ is obtained in two steps: *i*) Matching at $\mathcal{Q} = \mathcal{Q}_m$ of the HE Lagrangian to an effective Lagrangian (which has all light-fields interactions allowed by the HE symmetry), and *ii*) Running of the effective couplings from \mathcal{Q}_m to \mathcal{Q}_{EW} . The matching of the couplings of the light scalar sector can be obtained by exploiting the LE and HE effective potentials.

By construction, the HE and LE theories (in the presence of only light-field backgrounds) have the same RG-improved potentials at the decoupling scale, i.e. $V_{LE}(\mathcal{Q}_m) = V(\mathcal{Q}_m)$ (LE quantities carry a “LE” subscript; for HE quantities the subscript “HE” is suppressed). In the ideal case of perfect scale invariance this equivalence is true at any scale, and the choice of \mathcal{Q}_m at which one matches the two potentials is fully arbitrary.

³ As customary, in Eq. (5) the matrix \mathcal{M}^2 is the squared mass spectrum in the presence of the background fields ϕ_i , and the diagonal matrix \mathcal{C} depends on the renormalization-scheme. Note that in Eq. (5) the t -dependence is implicit. Concerning the radiative corrections $Z_i(t)$, we remind that they appear due to the canonical normalization of the kinetic terms and $Z_i(0)$ can contain finite contributions which depend on the renormalization scheme.

However, in realistic situations where the potentials are calculated at a given loop approximation, \mathcal{Q}_m has to be set at a value that presumably minimizes the unknown higher order corrections coming from heavy fields. This motivates the choice $\mathcal{Q}_m \sim M$.

The decoupling procedure involves some technical details when applied to realistic frameworks. The first issue arises when the HE theory contains several Higgses acquiring VEVs. This complication does not lead to any difficulties in our analysis as we are considering scenarios with the SM as an effective description. In fact we are restricting ourselves to cases where all the multi-Higgs VEVs in the interaction-eigenstate basis can be aligned along a unique VEV in the mass-eigenstate basis (and this direction corresponds to the light Higgs of the SM). Clearly this is possible because we are assuming the mass of the extra-Higgses to be $M \gg \mathcal{Q}_{EW}$. A second issue arises when there are several heavy fields with masses M_I . If they differ by orders of magnitude, there exists no choice of \mathcal{Q}_m that avoids large logarithms in the matching conditions. In this case the decoupling procedure has to be repeated as many times as the number of hierarchically different heavy mass thresholds. Of course when all M_I are similar, the decoupling can be performed just once with \mathcal{Q}_m fixed at some intermediate value among M_I . The simplified concrete examples of the next sections will better clarify the details of the decoupling procedure.

III. DECOUPLING IN SOME TOY MODELS

In this section we illustrate the previous ideas about decoupling. We first analyze a toy model with only one heavy degree of freedom. Second we consider a case with several scalar heavy particles and light fermions which contribute to the light scalar wave function renormalization. The reader not interested in those technical details can jump straightforwardly to section IV.

A. First toy model: scalars

We consider a toy model consisting of a light scalar ϕ and a heavy scalar S with a HE Lagrangian

$$\mathcal{L} = -\Omega + \frac{1}{2}(\partial\phi)^2 + \frac{1}{2}(\partial S)^2 - \frac{1}{2}m^2\phi^2 - \frac{1}{4!}\lambda\phi^4 - \frac{1}{2}M^2S^2 - \frac{1}{2}h^2\phi^2S^2 \quad (7)$$

where for simplicity the quartic coupling of the S field has been set to zero, although it is not protected by any symmetry. After decoupling the S -field the theory is described by the effective Lagrangian

$$\mathcal{L}_{LE} = -\Omega_{LE} + \frac{1}{2}(\partial\phi_{LE})^2 - \frac{1}{2}m_{LE}^2\phi_{LE}^2 - \frac{1}{4!}\lambda_{LE}\phi_{LE}^4. \quad (8)$$

In both Lagrangians the parameters are running with the scale \mathcal{Q} .

Since S does not acquire a VEV, the electroweak breaking field ϕ is aligned to ϕ_{LE} . The tree-level (RG-improved) matching of the parameters at the scale \mathcal{Q}_m is then trivial:

$$\Omega_{LE}(\mathcal{Q}_m) = \Omega(\mathcal{Q}_m), \quad \phi_{LE}(\mathcal{Q}_m) = \phi(\mathcal{Q}_m), \quad m_{LE}(\mathcal{Q}_m) = m(\mathcal{Q}_m), \quad \lambda_{LE}(\mathcal{Q}_m) = \lambda(\mathcal{Q}_m). \quad (9)$$

At this point one could already run the LE parameters from \mathcal{Q}_m to \mathcal{Q}_{EW} and obtain the minimization condition. However the result would strongly depend on the choice of \mathcal{Q}_m . Indeed, since the LE and HE parameters run very differently, one would obtain different minimization conditions for different values of \mathcal{Q}_m in Eq. (9), even though the fundamental HE parameters would be kept fixed. This problem is alleviated by performing a one-loop matching. Hereafter we adopt the $\overline{\text{MS}}$ renormalization scheme to subtract the one-loop divergences.

In the present model there is no one-loop wave function renormalization and the tree level relation $\phi_{LE}(\mathcal{Q}_m) = \phi(\mathcal{Q}_m)$ is preserved at one loop. The one-loop matching of the other parameters can be obtained by matching the LE and the HE effective potentials where all parameters are at the scale \mathcal{Q}_m . We thus impose the relation

$$V(\phi) = V_{LE}(\phi) \quad (10)$$

with

$$V(\phi) = \Omega + \frac{1}{2}m^2\phi^2 + \frac{1}{4!}\lambda\phi^4 + \frac{1}{64\pi^2} \sum_{i=\phi,S} m_i^4 \left(\log \frac{m_i^2}{\mathcal{Q}_m^2} - \frac{3}{2} \right), \quad (11)$$

$$V_{LE}(\phi) = \Omega_{LE} + \frac{1}{2}m_{LE}^2\phi^2 + \frac{1}{4!}\lambda_{LE}\phi^4 + \frac{1}{64\pi^2}m_{\phi_{LE}}^4 \left(\log \frac{m_{\phi_{LE}}^2}{\mathcal{Q}_m^2} - \frac{3}{2} \right) + \mathcal{O}(\phi^6/M^2),$$

where $m_\phi^2 = m^2 + \frac{1}{2}\lambda\phi^2$, $m_S^2 = M^2 + h^2\phi^2$ and $m_{\phi_{LE}}^2 = m_{LE}^2 + \frac{1}{2}\lambda_{LE}\phi^2$. This matching leads to

$$\begin{aligned} \Omega_{LE}(\mathcal{Q}_m) &= \left[\Omega(\mathcal{Q}_m) + \frac{M^4}{64\pi^2} \log \frac{M^2}{\mathcal{Q}_m^2} \right] - \frac{3M^4}{128\pi^2}, \\ m_{LE}^2(\mathcal{Q}_m) &= \left[m^2(\mathcal{Q}_m) + \frac{h^2M^2}{16\pi^2} \log \frac{M^2}{\mathcal{Q}_m^2} \right] - \frac{h^2M^2}{16\pi^2}, \\ \lambda_{LE}(\mathcal{Q}_m) &= \left[\lambda(\mathcal{Q}_m) + \frac{12}{32\pi^2}h^4 \log \frac{M^2}{\mathcal{Q}_m^2} \right], \end{aligned} \quad (12)$$

which holds up to two-loop corrections⁴ and where all parameters are evaluated at the scale \mathcal{Q}_m . In particular due to the one-loop scale invariance of the HE and LE effective potentials, the arbitrariness of \mathcal{Q}_m in (12) is guaranteed at one-loop level. In fact, within

⁴ Notice that the difference between LE and HE parameters is at one-loop. Using the former or the latter in the one-loop contribution to the effective potential makes a difference only at the two-loop order.

each parenthesis in (12) the logarithms compensate the one-loop running of the parameters whose evolutions are given by the β -functions

$$\begin{aligned}\beta_{\Omega_{LE}} &= \frac{1}{32\pi^2} m_{LE}^4, & \beta_{\Omega} &= \frac{1}{32\pi^2} (m^4 + M^4), \\ \beta_{m_{LE}^2} &= \frac{1}{8\pi^2} \lambda_{LE} m_{LE}^2, & \beta_{m^2} &= \frac{1}{8\pi^2} (\lambda m^2 + h^2 M^2), \\ \beta_{\lambda_{LE}} &= \frac{6}{32\pi^2} \lambda_{LE}^2, & \beta_{\lambda} &= \frac{6}{32\pi^2} (\lambda^2 + 4h^4).\end{aligned}\tag{13}$$

This shows that in the one-loop approximation the running and matching procedures are commutative and the LE theory is independent of the decoupling scale.

The one-loop freedom in the choice of \mathcal{Q}_m can be used to minimize the higher order corrections to (12). To this aim it is sensible to choose $\mathcal{Q}_m = M$. In this particular case the matching for m_{LE}^2 turns out to be

$$m_{LE}^2(M) = m^2(M) - \frac{h^2}{16\pi^2} M^2.\tag{14}$$

The second term is the large threshold correction that, in a more realistic theory as the SM, would destabilize the electroweak vacuum and would introduce a hierarchy problem.

B. Second toy model: scalars and fermions

Further subtleties of the decoupling procedure can be described in a slightly more complicated scenario. We consider a HE theory consisting of a light real scalar ϕ , a light Dirac fermion Ψ , and a number of heavy scalars S_I with masses M_I of similar magnitude:

$$\mathcal{L} = -\Omega + \frac{1}{2}(\partial\phi)^2 - \frac{m^2}{2}\phi^2 - \frac{\lambda}{4!}\phi^4 + \frac{1}{2}\sum_I [(\partial S_I)^2 - (M_I^2 + h_I^2\phi^2)S_I^2] + i\bar{\Psi}\not{\partial}\Psi - Y\phi\bar{\Psi}\Psi.\tag{15}$$

This theory can be described at LE by

$$\mathcal{L}_{LE} = -\Omega_{LE} + \frac{1}{2}(\partial\phi_{LE})^2 - \frac{m_{LE}^2}{2}\phi_{LE}^2 - \frac{\lambda_{LE}}{4!}\phi_{LE}^4 + i\bar{\Psi}_{LE}\not{\partial}\Psi_{LE} - Y_{LE}\phi_{LE}\bar{\Psi}_{LE}\Psi_{LE}\tag{16}$$

and the dependence with respect to \mathcal{Q} is implicit in Eqs. (15) and (16).

Since the S_I fields do not acquire any VEV and their integration do not lead to any tree-level correction to the LE couplings, the tree-level matching is trivial. Instead the one-loop matching is less straightforward due, for instance, to the radiative corrections to the kinetic terms. As we are interested in the one-loop electroweak-breaking conditions, we will focus on the effective potential of the light scalar field. In particular we want to calculate the RG-improved one-loop matching at the scale \mathcal{Q}_m . To this aim we first run all parameters in (15) and (16) to \mathcal{Q}_m , and then we evaluate the LE and HE one-loop potentials. We

thus absorb the kinetic-term radiative corrections⁵ $(1 - 2Z_\phi(\mathcal{Q}_m))$ and $(1 - 2Z_{\phi_{LE}}(\mathcal{Q}_m))$ into field redefinitions of ϕ and ϕ_{LE} respectively, where $d[Z_i(\mathcal{Q})]/d\log \mathcal{Q} = \gamma_i(\mathcal{Q})$, the anomalous dimension at one-loop order. The expansion of the HE RG-improved one-loop potential at the scale \mathcal{Q}_m has to reproduce the LE one at the same scale. We hence impose the matching

$$V(\phi) = V_{LE}(\phi) \quad (17)$$

with

$$\begin{aligned} V(\phi) &= \Omega + \frac{1}{2}m^2(1 + 2Z_\phi)\phi^2 + \frac{1}{4!}\lambda(1 + 4Z_\phi)\phi^4 + \sum_{i=\phi, S_I, \Psi} n_i \frac{m_i^4}{64\pi^2} \left(\log \frac{m_i^2}{\mathcal{Q}_m^2} - \frac{3}{2} \right), \\ V_{LE}(\phi_{LE}) &= \Omega_{LE} + \frac{1}{2}m_{LE}^2(1 + 2Z_{\phi_{LE}})\phi_{LE}^2 + \frac{1}{4!}\lambda_{LE}(1 + 4Z_{\phi_{LE}})\phi_{LE}^4 \\ &+ \sum_{i=\phi_{LE}, \Psi_{LE}} n_i \frac{m_i^4}{64\pi^2} \left(\log \frac{m_i^2}{\mathcal{Q}_m^2} - \frac{3}{2} \right) + \mathcal{O}(\phi_{LE}^6/M^2), \end{aligned}$$

where $m_\phi^2 = m^2 + \frac{1}{2}\lambda\phi^2$, $m_{S_I}^2 = M_I^2 + h^2\phi^2$, $m_\Psi^2 = Y^2\phi^2$, $m_{\phi_{LE}}^2 = m_{LE}^2 + \frac{1}{2}\lambda_{LE}\phi_{LE}^2$, $m_{\Psi_{LE}}^2 = Y_{LE}^2\phi_{LE}^2$, and $n_i = 1(-4)$ for real bosons (Dirac fermions). This implies

$$\begin{aligned} \Omega_{LE}(\mathcal{Q}_m) &= \left[\Omega(\mathcal{Q}_m) + \sum_I \frac{M_I^4}{64\pi^2} \log \frac{M_I^2}{\mathcal{Q}_m^2} \right] - \sum_I \frac{3M_I^4}{128\pi^2}, \\ m_{LE}^2(\mathcal{Q}_m) &= \left[m^2(\mathcal{Q}_m)[1 + 2Z_\phi(\mathcal{Q}_m) - 2Z_{\phi_{LE}}(\mathcal{Q}_m)] + \sum_I \frac{h_I^2 M_I^2}{16\pi^2} \log \frac{M_I^2}{\mathcal{Q}_m^2} \right] - \sum_I \frac{h_I^2 M_I^2}{16\pi^2}, \\ \lambda_{LE}(\mathcal{Q}_m) &= \lambda(\mathcal{Q}_m)[1 + 4Z_\phi(\mathcal{Q}_m) - 4Z_{\phi_{LE}}(\mathcal{Q}_m)] + \sum_I \frac{12h_I^4}{32\pi^2} \log \frac{M_I^2}{\mathcal{Q}_m^2}. \end{aligned} \quad (18)$$

where we are using the tree-level matching condition $Y_{LE}(\mathcal{Q}_m) = Y(\mathcal{Q}_m)$ ⁶.

In this toy model only light degrees of freedom generate Z_ϕ and $Z_{\phi_{LE}}$ at one loop via couplings that have trivial tree-level matching: it thus follows that $Z_\phi = Z_{\phi_{LE}}$ ⁷. Moreover, in view of minimizing higher loop corrections in the matching of m_{LE}^2 , one can adopt the choice $\mathcal{Q}_m = \bar{M}$, where \bar{M} is defined as

$$\log \bar{M}^2 = \frac{\sum_I h_I^2 M_I^2 \log(M_I^2)}{\sum_I h_I^2 M_I^2}, \quad (19)$$

⁵ In the rest of the paper we will consider only the one-loop RG-improved Coleman-Weinberg potential and wave function corrections. We then simplify the notation by omitting the superscript “(1)” in the quantities $Z_i^{(1)}$ and $\Delta V^{(1)}$ introduced in section II.

⁶ The one-loop matching condition for the Yukawa coupling Y can be obtained diagrammatically. For the purposes of the present paper we do not need to make it explicit.

⁷ Note that in scenarios (as the MSSM) where also some heavy fields contribute to Z_ϕ , the $Z_\phi - Z_{\phi_{LE}}$ dependence on \mathcal{Q}_m is necessary to guarantee the scale independence of the one-loop matching conditions.

so that only the non-logarithmic one-loop contribution is left in the matching condition for m_{LE}^2 :

$$m_{LE}^2(\bar{M}) = m^2(\bar{M}) - \sum_I \frac{h_I^2}{16\pi^2} M_I^2. \quad (20)$$

Of course the choice (19) is expected to reduce the higher order correction in the m_{LE}^2 matching only if the relation $\log \bar{M}^2 \sim \log M_I^2$ occurs for all I 's. Otherwise, as already stressed in section II, it is worth reiterating the decoupling procedure at each mass scale M_I .

IV. THE SM/MSSM MATCHING

In this section we use the MSSM one-loop RG-improved effective potential to determine the radiative corrections that can destabilize the electroweak breaking condition in such a model. The Higgs sector contains two doublets H_1 and H_2 where in our convention H_2 gives the mass to the top quark and H_1 to the bottom quark and tau lepton. The MSSM Higgs Lagrangian, including only the one-loop wave function renormalization for the Higgs doublets Z_{h_i} , can be written as

$$\mathcal{L}_{MSSM} = \sum_i (1 - 2Z_{h_i}) |D_\mu H_i|^2 - V(H_1, H_2) \quad (21)$$

where $V(H_1, H_2)$ is the tree-level MSSM Higgs potential. Then the one-loop RG-improved MSSM potential of the neutral Higgs fields $h_i = \text{Re}H_i^0$ ($i = 1, 2$) is

$$\begin{aligned} V(h_1, h_2) = & m_1^2(1 + 2Z_{h_1})h_1^2 + m_2^2(1 + 2Z_{h_2})h_2^2 - 2m_3^2 h_1 h_2 (1 + Z_{h_1} + Z_{h_2}) \\ & + \frac{g_Z^2}{2} [h_1^2(1 + 2Z_{h_2}) - h_2^2(1 + 2Z_{h_1})]^2 + \Delta V_{MSSM}(h_1, h_2), \end{aligned} \quad (22)$$

where ΔV_{MSSM} is the Coleman-Weinberg contribution generated by all fields of the MSSM and ⁸

$$g_Z^2 \equiv (g_Y^2 + g_2^2)/4, \quad m_1^2 \equiv m_{H_1}^2 + \mu^2, \quad m_2^2 \equiv m_{H_2}^2 + \mu^2. \quad (23)$$

The above equation is understood at an arbitrary renormalization scale \mathcal{Q} .

In view of the strong LHC bounds on the masses of supersymmetric particles we match the MSSM with the SM at some high scale \mathcal{Q}_m , say (multi-)TeV. To this aim we employ the effective potential techniques adopted in the previous examples. For simplicity, we assume all parameters to be real although the extension to cases with complex parameters (and CP violation) is straightforward.

Contrarily to the previous examples, the MSSM has two fields, in the gauge eigenstate basis, that acquire VEVs. We then go to the mass eigenstate basis to work out the

⁸ We are using conventionally for the electroweak gauge couplings the notation: $g' \equiv g_Y$ and $g \equiv g_2$. g_Y is related to the $U(1)$ gauge coupling g_1 in the $SU(5)$ normalization by $g_Y = \sqrt{\frac{3}{5}}g_1$.

matching (first at the tree-level, then at one-loop). The field rotation can be performed by neglecting the $\mathcal{O}(v^2/\mathcal{Q}_m^2)$ electroweak-breaking contributions (i.e. we proceed in the electroweak symmetry unbroken phase) since the CP -odd Higgs mass m_A is assumed to be much heavier than the electroweak scale. The resulting potential can be matched to the SM potential whose one-loop RG-improved expression is given by

$$V_{LE}(h_{LE}) = -m_{LE}^2(1 + 2Z_{h_{LE}})h_{LE}^2 + \frac{\lambda_{LE}}{2}(1 + 4Z_{h_{LE}})h_{LE}^4 + \Delta V_{SM}(h_{LE}) , \quad (24)$$

where ΔV_{SM} is the SM one-loop RG-improved Coleman-Weinberg potential in the presence of the background field $h_{LE} = \text{Re } H_{LE}^0$ (with H_{LE} being the SM Higgs doublet). In Eq. (24) and hereafter the effective higher-order operators, which are small due the large hierarchy between heavy and light fields, are neglected.

A. Tree-level matching

In order to derive the RG-improved tree-level matching we will focus on the quadratic part of the tree-level MSSM potential, $V^{(0)}(h_1, h_2)$, which can be extracted from (22). At the matching scale \mathcal{Q}_m we thus obtain

$$V^{(0)}(h_1, h_2) = m_1^2 h_1^2 + m_2^2 h_2^2 - 2m_3^2 h_1 h_2 + \frac{g_Z^2}{2}(h_1^2 - h_2^2)^2 . \quad (25)$$

The potential $V^{(0)}(h_1, h_2)$ at \mathcal{Q}_m can be rewritten in the mass eigenstate basis as

$$V^{(0)}(h, H) = -m^2 h^2 + m_H^2 H^2 + \dots \quad (26)$$

with $m^2 \ll m_H^2 = m_1^2 + m_2^2 + m^2$. This field transformation is achieved by the rotation

$$\begin{pmatrix} h_1 \\ h_2 \end{pmatrix} = R_\beta \begin{pmatrix} h \\ H \end{pmatrix} , \quad R_\beta = \begin{pmatrix} \cos \beta & -\sin \beta \\ \sin \beta & \cos \beta \end{pmatrix} \quad (27)$$

such that

$$(h_1, h_2) \begin{pmatrix} m_1^2 & -m_3^2 \\ -m_3^2 & m_2^2 \end{pmatrix} \begin{pmatrix} h_1 \\ h_2 \end{pmatrix} = (h, H) \begin{pmatrix} -m^2 & 0 \\ 0 & m_H^2 \end{pmatrix} \begin{pmatrix} h \\ H \end{pmatrix} . \quad (28)$$

Note that Eqs. (27) and (28) are equivalent to require

$$m_3^4 = (m_1^2 + m^2)(m_2^2 + m^2) , \quad (29)$$

$$\tan 2\beta = \frac{2m_3^2}{m_2^2 - m_1^2} \quad (30)$$

or, alternatively

$$\begin{aligned} m_2^2 &= m_3^2 / \tan \beta - m^2 , \\ m_1^2 &= m_3^2 \tan \beta - m^2 , \end{aligned} \quad (31)$$

with $\tan \beta \geq 1$, as we are assuming $m_3^2 > 0$ and $m_1^2 \geq m_2^2$. In particular since we are in the decoupling limit $m^2 \ll m_H^2$, our definition of $\tan \beta$ coincides with the MSSM one, $\tan \beta = v_2/v_1$ where $v_i \equiv \langle h_i \rangle$. Combining the above relations we also have the explicit expression for the light mass eigenstate

$$m^2 = -m_1^2 \cos^2 \beta - m_2^2 \sin^2 \beta + m_3^2 \sin 2\beta . \quad (32)$$

In the mass eigenstate basis (h, H) it is easy to obtain the tree-level matching to the SM. If one extracts the $V_{LE}^{(0)}(h_{LE})$ part from the LE one-loop potential of h_{LE} , Eq. (24), and matches it to $V^{(0)}(h, H)$ at \mathcal{Q}_m , one obtains

$$h_{LE}(\mathcal{Q}_m) = h(\mathcal{Q}_m), \quad m_{LE}(\mathcal{Q}_m) = m(\mathcal{Q}_m), \quad \lambda_{LE}(\mathcal{Q}_m) = g_Z^2(\mathcal{Q}_m) \cos^2 2\beta \quad [\text{tree level}] . \quad (33)$$

Similarly the matching of the LE and HE Yukawa interactions at \mathcal{Q}_m yields

$$y_t(\mathcal{Q}_m) = Y_t(\mathcal{Q}_m) \sin \beta , \quad y_{b,\tau}(\mathcal{Q}_m) = Y_{b,\tau}(\mathcal{Q}_m) \cos \beta , \quad [\text{tree level}] \quad (34)$$

where $y_{t,\tau,b}$ and $Y_{t,\tau,b}$ are respectively the top quark, tau lepton and bottom quark Yukawa couplings in the SM and in the MSSM.

Of course there might already be a problem at tree level: the right hand side of Eq. (32) is a linear combination of potentially large masses squared while the left hand side is a mass squared which is required to be at the electroweak scale. This fine-tuning is essentially equivalent to that in Eq. (29). This is the main naturalness problem in the MSSM. This problem cannot be tackled unless we know the (fundamental) theory responsible for triggering supersymmetry breaking at the high scale \mathcal{M} in the hidden sector and dictating the size of the supersymmetry breaking parameters in the observable sector. The FP solution [12, 13] just uncovers the functional relationships between fundamental parameters at the high scale \mathcal{M} for which the naturalness problem is circumvented. However even if we accept that the fundamental theory might provide a solution to the tree-level stability we still have to worry for loop corrections, e.g. in the effective theory as those computed in Ref. [8]. The matching including one loop corrections will be done in the next section.

B. One-loop level matching

We now proceed with the one-loop matching. Again we want to work in the mass eigenstate basis. We then impose the tree-level matching conditions (31) in the one-loop term

ΔV_{MSSM} of (22), and expand ΔV_{MSSM} . Such expansion produces some new quadratic contributions that we absorb as

$$V(h_1, h_2) = \tilde{m}_1^2 h_1^2 + \tilde{m}_2^2 h_2^2 - 2\tilde{m}_3^2 h_1 h_2 + \dots \quad (35)$$

where

$$\begin{aligned} \tilde{m}_i^2 &= m_i^2 + 2Z_{h_i} m_i^2 + \Delta m_i^2, & \Delta m_i^2 &= \left. \frac{\partial \Delta V_{MSSM}}{\partial h_i^2} \right|_{h_i=0}, \\ \tilde{m}_3^2 &= m_3^2 + (Z_{h_1} + Z_{h_2}) m_3^2 + \Delta m_3^2, & \Delta m_3^2 &= -\frac{1}{2} \left. \frac{\partial \Delta V_{MSSM}}{\partial h_1 h_2} \right|_{h_i=0}, \end{aligned} \quad (36)$$

with $i = 1, 2$. As previously done for the tree-level matching, we diagonalize the quadratic potential (35) by a rotation R_β (whose angle differs from that of section IV A although for notational simplicity we are keeping the same notation for both) leading to a light mass eigenstate h with squared mass $-\tilde{m}^2$ and a heavy eigenstate H with squared mass $\tilde{m}_H^2 = \tilde{m}_1^2 + \tilde{m}_2^2 + \tilde{m}^2$, where

$$\begin{aligned} \tilde{m}^2 &= m^2 + 2m^2 Z_h + \Delta m^2, \\ m^2 &= -m_1^2 \cos^2 \beta - m_2^2 \sin^2 \beta + m_3^2 \sin 2\beta, \\ \Delta m^2 &= -\Delta m_1^2 \cos^2 \beta - \Delta m_2^2 \sin^2 \beta + \Delta m_3^2 \sin 2\beta, \end{aligned} \quad (37)$$

and $Z_h \equiv \cos^2 \beta Z_{h_1} + \sin^2 \beta Z_{h_2}$ is the wave function renormalization in the MSSM for the mass eigenstate h ⁹. This diagonalization requires

$$\begin{aligned} \tilde{m}_2^2 &= \tilde{m}_3^2 / \tan \beta - \tilde{m}^2, \\ \tilde{m}_1^2 &= \tilde{m}_3^2 \tan \beta - \tilde{m}^2, \end{aligned} \quad (38)$$

which can be used to express, as it is customary in the MSSM, $\tan \beta$ and the lightest eigenvalue $-\tilde{m}^2$ as functions of the fundamental parameters:

$$\begin{aligned} \tilde{m}^2 &= -\tilde{m}_1^2 \cos^2 \beta - \tilde{m}_2^2 \sin^2 \beta + \tilde{m}_3^2 \sin 2\beta, \\ \tan 2\beta &= \frac{2\tilde{m}_3^2}{\tilde{m}_2^2 - \tilde{m}_1^2}. \end{aligned} \quad (39)$$

In order to perform the complete one-loop matching, we would need to consider ΔV_{SM} . As the light (i.e. SM) fields provide the same contributions to ΔV_{SM} and ΔV_{MSSM} modulo the tree-level matching of Yukawa couplings in Eq. (34) (cf. also section V), we can proceed by taking into account only the heavy non-SM fields in Δm_1^2 , Δm_2^2 and Δm_3^2 . Then the

⁹ Notice that the identity $Z_h m^2 \equiv Z_{h_1} m_1^2 \cos^2 \beta + Z_{h_2} m_2^2 \sin^2 \beta - (Z_{h_1} + Z_{h_2}) m_3^2 \sin 2\beta$ is obtained after using the tree-level matching conditions, Eq. (31), on the masses and mixing angle, as required by the fact that the wave function renormalization is already a one-loop effect.

one-loop RG-improved matching of the quadratic term in the HE and LE theories turns out to be

$$m_{LE}^2(\mathcal{Q}_m) = m^2(\mathcal{Q}_m)(1 + 2\Delta Z_h(\mathcal{Q}_m)) + \Delta m^2(\mathcal{Q}_m), \quad (40)$$

where

$$\Delta Z_h(\mathcal{Q}_m) = Z_h(\mathcal{Q}_m) - Z_{h_{LE}}(\mathcal{Q}_m). \quad (41)$$

We hence stress that the requirement $m^2(\mathcal{Q}_m) \sim \mathcal{Q}_{EW}^2$ is not sufficient to guarantee sensible electroweak breaking conditions as these could be destabilized by Δm^2 and ΔZ_h .

V. Δm^2 IN THE MSSM WITH HIGH SUSY-BREAKING SCALE

We will now explain the main lines to determine Δm^2 . We remind that all the MSSM particles, except the SM ones, are assumed to be heavy. The one-loop potential $\Delta V_{MSSM}(h_1, h_2)$ can be split into two separated terms: one term contains contributions from the Higgs sector A, H, H^\pm and the SM fields, and the second one does it from the field superpartners \tilde{f} (with \tilde{f} representing the whole list of squarks, sleptons, charginos and neutralinos). Due to the triviality of the tree-level matching conditions (33) and (34), it is easy to see as we already noticed that the light fields provide the same contribution to $\Delta V_{MSSM}(h)$ and to $\Delta V_{SM}(h_{LE})$. Because of this property the correction Δm^2 can be calculated as in Eq. (37) with

$$\Delta m_I^2 = \sum_{r=\tilde{f}, A, H, H^\pm} \Delta m_{I,r}^2, \quad (42)$$

$$\Delta m_{i,r}^2 = \left. \frac{\partial \Delta V_{MSSM,r}}{\partial h_i^2} \right|_{h_i=0}, \quad \Delta m_{3,r}^2 = -\frac{1}{2} \left. \frac{\partial \Delta V_{MSSM,r}}{\partial h_1 h_2} \right|_{h_i=0}, \quad (43)$$

where $I = (i, 3)$, $i = 1, 2$ and $\Delta V_{MSSM,r}$ is the MSSM one-loop potential generated by the field r .

The explicit form of $\Delta m_{I,r}^2$ depends on the renormalization scheme. In the $\overline{\text{MS}}$ scheme (or equivalently at this level $\overline{\text{DR}}$), for which $\mathcal{C} = (3/2)\mathbf{1}$ in Eq. (5) for scalars and fermions, it results

$$\Delta m_{i,r}^2 = \frac{n_r}{32\pi^2} \left. \frac{\partial m_r^2}{\partial h_i^2} G(m_r^2) \right|_{h_i=0}, \quad \Delta m_{3,r}^2 = -\frac{n_r}{64\pi^2} \left. \frac{\partial m_r^2}{\partial h_1 h_2} G(m_r^2) \right|_{h_i=0}, \quad (44)$$

where n_r stands for the number of degrees of freedom of the particle r and is positive (negative) for bosons (fermions), while the function $G(x^2)$ is defined as

$$G(x^2) \equiv x^2 \left(\log \frac{x^2}{\mathcal{Q}_m^2} - 1 \right). \quad (45)$$

For simplicity we determine Δm^2 by neglecting the corrections coming from first and second generations of squarks and sleptons (the expressions are however fully general and

the first two generation sfermions could be easily included). Details of the calculation are furnished in appendix A. Here we only report on the final result:

$$\begin{aligned}
-\Delta m^2 = & \frac{1}{32\pi^2} \left\{ 6y_t^2 [G(m_Q^2) + G(m_U^2)] + 6y_b^2 [G(m_Q^2) + G(m_D^2)] + 2y_\tau^2 [G(m_L^2) + G(m_E^2)] \right. \\
& + 6y_t^2 X_t^2 \frac{G(m_Q^2) - G(m_U^2)}{m_Q^2 - m_U^2} + 6y_b^2 X_b^2 \frac{G(m_Q^2) - G(m_D^2)}{m_Q^2 - m_D^2} + 2y_\tau^2 X_\tau^2 \frac{G(m_L^2) - G(m_E^2)}{m_L^2 - m_E^2} \\
& - g_Y^2 (G(m_Q^2) - 2G(m_U^2) + G(m_D^2) - G(m_L^2) + G(m_E^2)) \cos 2\beta \\
& - 6g_2^2 \frac{M_2^2 G(M_2^2) - \mu^2 G(\mu^2)}{M_2^2 - \mu^2} - 2g_Y^2 \frac{M_1^2 G(M_1^2) - \mu^2 G(\mu^2)}{M_1^2 - \mu^2} \\
& - \left(12g_2^2 M_2 \mu \frac{G(M_2^2) - G(\mu^2)}{M_2^2 - \mu^2} + 4g_Y^2 M_1 \mu \frac{G(M_1^2) - G(\mu^2)}{M_1^2 - \mu^2} \right) \sin \beta \cos \beta \\
& \left. + G(m_H^2) (-6g_Z^2 \cos^2 2\beta + 2g_Z^2 + g_2^2) \right\}, \tag{46}
\end{aligned}$$

where the soft breaking terms $X_{t,b,\tau}$ are defined as

$$X_t = A_t - \frac{\mu}{\tan \beta}, \quad X_{b,\tau} = A_{b,\tau} - \mu \tan \beta. \tag{47}$$

In Eq. (46) the first two lines correspond to the contribution from sfermions, the third line the Fayet-Iliopoulos contribution from scalars, the fourth and fifth lines the contribution from charginos and neutralinos, and the last line the contribution from the heavy scalar, the pseudoscalar and charged Higgses. Notice that all supersymmetric parameters are defined at the scale \mathcal{Q}_m .

We therefore conclude that in the MSSM with heavy non-SM particles, the Higgs sector at LE appears like the one of the SM where the Higgs quadratic parameter at the scale \mathcal{Q}_m is given by the relation of Eq. (40) with m^2 and Δm^2 as in Eqs. (37) and (46) and $\tan \beta$ given by Eq. (39). Moreover the explicit expression of ΔZ_h ¹⁰ is not required in first approximation as we will see in section VI. Finally, it is worth noting that, by construction, in the considered heavy MSSM scenario the electroweak breaking condition at $\mathcal{Q} = \mathcal{Q}_{EW}$ can be evaluated in the LE theory (avoiding large logarithms) with no one-loop dependence on the choice of \mathcal{Q}_m (cf. appendix B).

VI. THE STABILITY OF THE SM EFFECTIVE THEORY

As reminded in section IV, even in the case that the fundamental theory naturally leads to m^2 of the order of the electroweak scale, we still have to worry about the destabilization and unnaturality due to radiative corrections. In this section we sketch some

¹⁰ In general ΔZ_h consists of two terms: one depending on the renormalization scale and proportional to the anomalous dimension difference $\gamma_h - \gamma_{h_{LE}}$; and a second one leading to a one-loop scale-independent difference between the LE and HE parameters (see e.g. [14]).

relationships that would help to not destabilize the vacuum. These can be deduced after evaluating the size of the radiative corrections at \mathcal{Q}_m and forcing them to be in the ballpark of $m^2(\mathcal{Q}_m)$.

Some preliminary observations are in order here:

- In general $m^2(\mathcal{Q})$ has a strong scale dependence and its value can span several order of magnitudes. Its (one-loop) radiative correction is given by $\Delta m^2(\mathcal{Q}) + 2m^2(\mathcal{Q})\Delta Z_h(\mathcal{Q})$, which is also strongly scale dependent. Hence claiming $m^2 = \mathcal{O}(100 \text{ GeV})^2$ makes sense only at a specific running scale and, concerning the hierarchy problem, the result is satisfactory only if at the same scale the radiative corrections contain no large logarithms (to keep perturbation theory trustable) and are of the order of the electroweak scale or below (to not destabilize the tree level result). In this section we assume that the supersymmetry breaking theory yields $m^2 = \mathcal{O}(100 \text{ GeV})^2$ at the specific scale \mathcal{Q}_m .
- The strong (one-loop) scale dependences of $m^2(\mathcal{Q})$ and its radiative correction have opposite signs and almost cancel out. Indeed the \mathcal{Q} -dependence of these two terms is equivalent to the one of $m_{LE}^2(\mathcal{Q})$ [cf. Eq. (40)], which amounts to the β -function of the SM (cf. appendix B) and is thus negligible for our purposes¹¹.
- Heavy particles have masses well above $m^2(\mathcal{Q}_m)$. This implies that the wave function correction $2m^2(\mathcal{Q}_m)\Delta Z_h(\mathcal{Q}_m)$ can be neglected in comparison to $\Delta m^2(\mathcal{Q}_m)$.

Therefore within the above approximations the stability of the electroweak breaking conditions under radiative corrections can be evaluated by means of the “stability parameter”

$$\mathcal{S} = \left| \frac{\Delta m^2(\mathcal{Q}_m)}{m_{LE}^2(\mathcal{Q}_m)} \right|, \quad (48)$$

where the running of m_{LE}^2 between \mathcal{Q}_{EW} and \mathcal{Q}_m can be neglected. We now proceed determining \mathcal{S} in some specific scenarios.

A. DEGENERATE CASE

We first consider the simplest MSSM scenario where all mass parameters are degenerate at some common value M :

$$m_Q = m_U = m_D = m_L = m_E = M_1 = M_2 = \mu = m_H \equiv M. \quad (49)$$

¹¹ We checked numerically that in the SM the quadratic term changes by about $\sim 10\%$ for a running from the electroweak to the Planck scale.

In this case all radiative corrections depend on the single logarithm $\log M^2/\mathcal{Q}_m^2$ and the simplest choice for the decoupling scale is obviously $\mathcal{Q}_m = M$. Eq. (46) thus yields

$$\Delta m^2(M) = \frac{M^2}{32\pi^2} \left\{ 12y_t^2 + 12y_b^2 + 4y_\tau^2 - 6\lambda - \frac{3}{2}g_Y^2 - \frac{9}{2}g_2^2 \right\}. \quad (50)$$

Eq. (50) reproduces the SM Higgs mass quadratic divergence obtained in the case that the SM has a cutoff $\Lambda \equiv M$ [8, 15]: the first three terms correspond to the quadratic divergences coming from exchange of top, bottom and tau fermions [with masses $m_{t,b,\tau}(h_{LE}) = y_{t,b,\tau}h_{LE}$], the fourth term corresponds to divergences coming from exchange of the SM Higgs [with mass $m_h^2(h_{LE}) = -m^2 + 3\lambda h_{LE}^2$] and neutral and charged Goldstone bosons [with masses $m_\chi^2(h_{LE}) = -m^2 + \lambda h_{LE}^2$], while the last two terms are equivalent to those due to exchange of W and Z gauge bosons [with masses $m_W^2(h_{LE}) = g_2^2 h_{LE}^2/2$ and $m_Z^2(h_{LE}) = (g_Y^2 + g_2^2)h_{LE}^2/2$]. In fact Eq. (50) can be written as a function of SM running masses as

$$\Delta m^2(M) = \frac{M^2}{32\pi^2 v^2} \left\{ \sum_f n_f m_f^2 - 3m_h^2 - 3m_Z^2 - 6m_W^2 \right\}, \quad (51)$$

where n_f is the number of degrees of freedom of the fermion f .

As it is well known, for experimental values of the SM masses the requirement $\Delta m^2(M) = 0$ in Eq. (51), usually dubbed Veltman condition [8], is not fulfilled at weak scales but at Planckian scales [16]. The value of this high scale is quantified in the left panel of Fig. 1 where the contour lines of $\Delta m^2(M) = 0$ (or equivalently $\mathcal{S} = 0$) are plotted in the plane $(\log_{10} M/\text{GeV}, M_t)$ (where M_t is the top quark pole mass), for different values of m_h and $\alpha_3(m_Z)$. The plot has been obtained by using the RG equations of the SM parameters appearing in Eq. (50) at the NNLO (as done e.g. in [17]).

B. A SIMPLE NON-DEGENERATE CASE

A simple non-degenerate case is the scenario where at the scale \mathcal{Q}_m the sfermion, electroweakino and Higgs sectors have each one a respective common mass M_0 , $M_{1/2}$ and M_H as:

$$m_Q = m_U = m_D = m_L = m_E \equiv M_0, \quad M_1 = M_2 = \mu \equiv M_{1/2}, \quad m_H \equiv M_H. \quad (52)$$

In this case we can express $\Delta m^2(\mathcal{Q}_m)$ as

$$\Delta m^2(\mathcal{Q}_m) = \Delta_f m^2(\mathcal{Q}_m) + \Delta_\ell m^2(\mathcal{Q}_m) \quad (53)$$




FIG. 1: *Left panel: Contour lines of $\mathcal{S} = 0$ for different values of $\alpha_3(m_Z)$ in the degenerate case of section VIA. The solid (dashed) lines are obtained taking $m_h = 126$ (125) GeV. Right panel: Contour lines of $\log_{10} M_0$ as a function of r_H and $r_{1/2}$ leading to $\mathcal{S} = 0$ in the non-degenerate case of section VIB. We choose $m_h = 125.1$ GeV, $m_t = 172.5$ GeV, $\alpha_3(m_Z) = 0.1196$ and $\tan \beta = 1$ for definiteness. Along the dotted straight line ($r_{1/2} = r_H$) the quadratic corrections can be understood as a generalization of the Veltman condition.*

with ¹²

$$\Delta_f m^2(\mathcal{Q}_m) = \frac{1}{32\pi^2} \left\{ (12y_t^2 + 12y_b^2 + 4y_\tau^2)M_0^2 - 4GM_{1/2}^2 - (6\lambda - G)M_H^2 \right\}, \quad (54)$$

$$\begin{aligned} \Delta_\ell m^2(\mathcal{Q}_m) = \frac{1}{32\pi^2} \left\{ - \left[(12y_t^2 + 12y_b^2 + 4y_\tau^2)M_0^2 + (6y_t^2 X_t^2 + 6y_b^2 X_b^2 + 2y_\tau^2 X_\tau^2) \right] \log \frac{M_0^2}{Q_m^2} \right. \\ \left. + 4GM_{1/2}^2 f_\beta \log \frac{M_{1/2}^2}{Q_m^2} + (6\lambda - G)M_H^2 \log \frac{M_H^2}{Q_m^2} \right\}, \quad (55) \end{aligned}$$

where $G = (g_Y^2 + 3g_2^2)/2$, $f_\beta = 2 + \sin 2\beta$ and all parameters are understood at the scale \mathcal{Q}_m . For definiteness we take $\mathcal{Q}_m = M_0$ hereafter. It follows

$$\Delta_f m^2(\mathcal{Q}_m) = \frac{M_0^2}{32\pi^2 v^2} \left\{ \sum_f n_f m_f^2 - 4(2m_W^2 + m_Z^2)r_{1/2}^2 - (3m_h^2 - 2m_W^2 - m_Z^2)r_H^2 \right\}, \quad (56)$$

$$\Delta_\ell m^2(\mathcal{Q}_m) = \frac{M_0^2}{32\pi^2 v^2} \left\{ 4f_\beta(2m_W^2 + m_Z^2)r_{1/2}^2 \log r_{1/2}^2 + (3m_h^2 - 2m_W^2 - m_Z^2)r_H^2 \log r_H^2 \right\}, \quad (57)$$

¹² The decomposition (53) separates the one-loop scale independent contribution from the dependent one, $\Delta_f m^2(\mathcal{Q}_m)$ and $\Delta_\ell m^2(\mathcal{Q}_m)$ respectively. This separation is defined up to an arbitrary one-loop scale independent quantity, but our results do not depend on the convention we are choosing.

where $r_H = M_H/M_0$, $r_{1/2} = M_{1/2}/M_0$ and all masses are running.

For $r_H = r_{1/2}$ the right hand side of Eq. (56) coincides with the quadratic divergences that one derives from the SM effective potential by using as regulators the cutoff $\Lambda_F = M_0$ for fermions and the cutoff $\Lambda_B = M_{1/2} = M_H$ for bosons¹³. In this case the requirement $\Delta_f m^2 = 0$ can be interpreted as a generalization of the Veltman condition expressing the Higgs quadratic divergence in terms of SM parameters and cutoffs¹⁴. However also the logarithmic corrections are sizeable and contribute to destabilizing the potential. With the constraint $r_H = r_{1/2}$, $\Delta_\ell m^2$ is negligible only when $r_H = r_{1/2} \approx 1$. Correspondingly the condition $\Delta m^2 = 0$ with cutoffs at the TeV scale is not possible anymore.

However, for $r_H \neq r_{1/2}$ the stability under perturbative corrections can still be fulfilled at scales of $\mathcal{O}(100 \text{ TeV})$ if one requires the cancellation of the total correction Δm^2 . This can be seen in the right panel of Fig. 1 where the constraints on r_H and $r_{1/2}$ leading to $\mathcal{S} = 0$ (solid curves) are plotted for several values of M_0 . The figure is obtained using Eqs. (56) and (57) with $\tan \beta = 5$, and running the SM parameter at NNLO with boundary conditions $m_h = 125.1 \text{ GeV}$, $M_t = 172 \text{ GeV}$ and $\alpha_3 = 0.1196$ at the electroweak scale as explained in Ref. [16]. The former case, $r_H = r_{1/2}$, is exhibited as the dotted straight line in the right panel of Fig. 1 where we can see that stability can be achieved for sub-Planckian scales.

C. GENERAL SOFT BREAKING TERMS

In principle one expects that all soft breaking parameters will be different at the scale \mathcal{Q}_m . In this general case the finite threshold contribution to the SM Higgs mass parameter is given by Eq. (46). The required condition for keeping Δm^2 at the order of the electroweak scale is then a hypersurface in the multidimensional space of supersymmetric parameters $(m_Q, m_U, m_D, m_L, m_E, A_t, A_b, A_\tau, m_1, m_2, \mu, M_1, M_2)$. In the next section some of these hypersurfaces are analyzed numerically for the case of negligible Fayet-Iliopoulos (FI) contribution¹⁵.

Before closing this section a couple of comments are in order. In this section we have assumed $m^2(\mathcal{Q}_m)$ at the electroweak scale without specifying the origin of such a value. The main ideas to naturally produce $m^2(\mathcal{Q}_m)$ at the electroweak scale are twofold:

¹³ A similar interpretation is not clear for $M_{1/2} \neq M_H$ due to the fact that both the heavy (neutral and charged) Higgs sector and charginos and neutralinos contribute to the generalized Veltman condition with terms proportional to squared gauge couplings.

¹⁴ In particular, for some choices of Λ_B and Λ_F at the TeV scale, the condition $\Delta_f m^2 = 0$ can be achieved at low energies.

¹⁵ We remind that the FI contribution is a RG invariant. Our assumption is thus valid only if the FI term is zero at the scale of supersymmetry breaking transmission. Otherwise it should be taken into account although its (tiny) contribution should not change the qualitative conclusions.

- If, *i*) the whole MSSM Higgs sector is at the electroweak scale (in which case the LE effective theory is not the SM but a two Higgs doublet model), and *ii*) the masses of the supersymmetric partners are in the low TeV region (i.e. $\mathcal{Q}_m \not\gg \mathcal{Q}_{EW}$), then $m^2(\mathcal{Q}_m)$ is at the electroweak scale. Moreover the requirement $|\Delta m^2(\mathcal{Q}_m)| \lesssim \mathcal{O}(100 \text{ GeV})^2$ is also automatically satisfied. This parameter configuration might be excluded soon by the LHC lower bounds on heavy Higgs and superpartner masses and we will not further discuss it.
- For $\tan \beta \gg 1$ the squared mass $m^2(\mathcal{Q}_m)$ is similar to $m_2^2(\mathcal{Q}_m)$ [cf. Eq. (37)]. The value of the latter is naturally small in the FP parameter region of the MSSM. This possibility has been broadly studied in the literature [12] although the modifications due to Δm^2 have been overlooked.

In the next section we will analyze the effects that the one-loop corrections have on the FP solution.

VII. NUMERICAL RESULTS: FOCUS POINT SOLUTIONS

In this section we concentrate on the FP solution including the one-loop radiative correction Δm^2 . We stress that till now we have expressed $m^2(\mathcal{Q}_m)$ and $\Delta m^2(\mathcal{Q}_m)$ as functions of supersymmetric parameters evaluated at the scale \mathcal{Q}_m . However, in view of a more fundamental supersymmetry-breaking description, m^2 and Δm^2 should be re-expressed in terms of the supersymmetric parameters evaluated at the messenger scale \mathcal{M} at which supersymmetry breaking is transmitted to the observable sector. As, depending on the supersymmetry breaking model, some of the parameters can unify at the messenger scale, the number of independent parameters at the scale \mathcal{M} can be smaller (than what would show up at the scale \mathcal{Q}_m). So, the required relation to keep Δm^2 and $m^2(\mathcal{Q}_m)$ of the order of the electroweak scale is simpler, and might appear more natural. This scenario is considered in this section assuming, at the scale \mathcal{M} , as a simple example the case ¹⁶

$$m_Q(\mathcal{M}) = m_U(\mathcal{M}) = M_0, \quad m_1(\mathcal{M}) = m_2(\mathcal{M}) = M_H, \quad M_a(\mathcal{M}) = M_{1/2}. \quad (58)$$

From Eqs. (39) and (40) we can write the matching conditions as

$$m_{LE}^2 \simeq m^2(\mathcal{Q}_m) + \Delta m^2(\mathcal{Q}_m) = \frac{1}{\tan^2 \beta - 1} \tilde{m}_1^2 - \frac{\tan^2 \beta}{\tan^2 \beta - 1} \tilde{m}_2^2 \quad (59)$$

where Δm^2 is given by Eq. (46), and the subleading radiative contribution $2m^2 \Delta Z_h$ from the wave function renormalization is neglected. For $\tan \beta \gg 1$, $\mu = \mathcal{O}(100 \text{ GeV})$ and small Δm^2 , the requirement $m^2(\mathcal{Q}_m) = \mathcal{O}(100 \text{ GeV})^2$ implies $|m_{H_2}^2| = \mathcal{O}(100 \text{ GeV})^2$,

¹⁶ Other cases can be obviously considered along similar lines. Here we just present the case of Eq. (58) dubbed as NUHM1 [18].

where $m_{H_{1,2}}^2$ are defined in Eq. (23). For heavy supersymmetric spectrum this size of $m_{H_2}^2$ can be generated in the neighborhood of the FP solution [13, 19] which is radiatively led to $m_{H_2}^2 = 0$. However in the FP parameter region there is no general reason why Δm^2 should be small. In fact the presence of Δm^2 can distort the standard FP solution usually considered in the literature. In this section we numerically analyze the parameter space of these modified FP solutions ¹⁷.

As it was mentioned in the previous section, it is useful to re-express the supersymmetric parameters of Eq. (59) in terms of their values at the high scale \mathcal{M} . On dimensional grounds we can rewrite them as

$$\begin{aligned}
m_X^2(\mathcal{Q}_m) &= m_X^2 + \eta_Q^X[\mathcal{Q}_m, \mathcal{M}](m_Q^2 + m_U^2 + m_2^2) + \sum_{a,b} \eta_{ab}^X[\mathcal{Q}_m, \mathcal{M}]M_a M_b \\
&+ \sum_a \eta_{aA}^X[\mathcal{Q}_m, \mathcal{M}]M_a A_0 + \eta_A^X[\mathcal{Q}_m, \mathcal{M}]A_0^2, \\
A_t(\mathcal{Q}_m) &= A_0 + f_A[\mathcal{Q}_m, \mathcal{M}]A_0 + \sum_a f_a[\mathcal{Q}_m, \mathcal{M}]M_a,
\end{aligned} \tag{60}$$

where A_0 , m_X^2 and M_a (with $X = H_i, Q_L, U_R, D_R$ and $a = 1, 2, 3$) are respectively the stop tri-linear mixing parameter, the sfermion masses and the Majorana gaugino masses at the scale \mathcal{M} .

Concerning the numerical procedure, we assume moderately large values of $\tan\beta$ (namely $\tan\beta \simeq 10$) which allows to approximate $X_t \simeq A_t$ and to safely neglect all Yukawa couplings except that of the top quark. Moreover we set A_t at the mass scale $\tilde{Q}^2 \equiv m_Q(\mathcal{Q}_m)m_U(\mathcal{Q}_m)$ in such a way that the SM RG evolution of the quartic coupling

$$\lambda(\tilde{Q}) = \frac{1}{4}(g_2^2(\tilde{Q}) + g_Y^2(\tilde{Q})) \cos^2 2\beta + \frac{3}{8\pi} y_t^4(\tilde{Q}) X_t^2 \left(1 - \frac{X_t^2}{12}\right) \tag{61}$$

from \tilde{Q} to \mathcal{Q}_{EW} reproduces the Higgs mass observation, namely $\lambda(\mathcal{Q}_{EW}) \approx (m_h/v)^2/2$. For the functions $\eta^X[\mathcal{Q}_m, \mathcal{M}]$ and $f[\mathcal{Q}_m, \mathcal{M}]$, which were obtained semi-analytically for $\mathcal{Q}_m = 2 \text{ TeV}$ and $\mu \sim 100 \text{ GeV}$ in Ref. [13], we use some simple generalized formulas where the effect of possible heavy Higgsinos is incorporated ¹⁸. Finally, as fundamental description of the supersymmetric parameters, we consider the relations in Eq. (58).

Figs. 2, 3, 4 and 5 display the values of fundamental parameters and their corresponding mass spectra leading to $m_{LE}^2 \sim (100 \text{ GeV})^2$. In the left panels of the figures we plot contour

¹⁷ We remind that, in the absence of Δm^2 , the FP solution is scale invariant with respect to a common multiplicative factor on the boundary conditions (at the scale \mathcal{M}) of the supersymmetric masses. This scale invariance is broken by Δm^2 which contains logarithms of the supersymmetry breaking masses over \mathcal{Q}_m . Still, as radiative corrections are small as compared to the tree level values, the scale invariance of the FP solutions is approximately preserved

¹⁸ For the case of heavy Higgsinos we determine the one-loop RG evolution of the MSSM parameters by neglecting the scale dependence of μ . This is justified by the fact that the variation of μ between \mathcal{Q}_m and \mathcal{M} is of the order of 1%.




FIG. 2: *Left panel: Contour lines of $\log_{10} \mathcal{M}/\text{GeV}$ (white labels) and stability parameter \mathcal{S} (orange labels) for $M_0 = 0$, and light Higgsinos, $\mu = 100$ GeV. The (yellow) external shadowed region corresponds to $m_U(\mathcal{Q}_m) < 750$ GeV and the (green) internal shadowed region corresponds to $m_U^2(\mathcal{Q}_m) < 0$. Middle panel: Corresponding contour lines of constant $m_Q(\mathcal{Q}_m)$ (solid black), $m_U(\mathcal{Q}_m)$ (dashed blue) and $m_H(\mathcal{Q}_m)$ (dash dotted red). Right panel: Corresponding contour lines of constant $M_3(\mathcal{Q}_m)$ (solid red), $M_2(\mathcal{Q}_m)$ (dashed blue) and $M_1(\mathcal{Q}_m)$ (dash dotted black). Labels in the middle and right panels are in TeV units.*




FIG. 3: *The same as in Fig. 2 but for $M_0 = 2$ TeV.*

(black dashed) lines for constant values of $\log_{10}(\mathcal{M}/\text{GeV})$ such that Eq. (59) at tree level is fulfilled, and contour (solid red) lines for the stability parameter \mathcal{S} of the one-loop radiative corrections (cf. Eqs. (46) and (48)). Similarly, the contour lines of $m_Q(\mathcal{Q}_m)$ (solid black), $m_U(\mathcal{Q}_m)$ (dashed blue) and $m_H(\mathcal{Q}_m)$ (dash-dotted red) are shown in the middle panels, whereas the contour lines of $M_3(\mathcal{Q}_m)$ (solid red), $M_2(\mathcal{Q}_m)$ (dashed blue) and $M_1(\mathcal{Q}_m)$ (dash-dotted black) are depicted in the right panels. We remind that the condition $m_{LE}^2 \sim (100 \text{ GeV})^2$ arises with no tuning between m^2 and Δm^2 when $\mathcal{S} = \mathcal{O}(1)$.

In Figs. 2 and 3 we consider the case $\mu \sim 100$ GeV, and $M_0 = 0$, $M_0 = 2$ TeV, respectively, and in Figs. 4 and 5 the case with $M_0 = 2$ TeV and heavy and superheavy




FIG. 4: The same as in Fig. 3 but for $\mu = 2$ TeV.




FIG. 5: The same as in Fig. 3 but for $\mu = M_{1/2}$. In the (magenta) upper left shadowed region the soft breaking terms $m_{H_U}^2 = m_{H_D}^2 < 0$ are tachyonic at the scale \mathcal{M} , although $m_1^2 = m_2^2 \equiv M_H^2 > 0$.

Higgsinos, with $\mu = 2$ TeV and $\mu = M_{1/2}$, respectively. The region with $m_U(\mathcal{Q}_m) < 750$ GeV [$m_U^2(\mathcal{Q}_m) < 0$] corresponds to the yellow [green] shadowed area. In fact the yellow band corresponds to the FP for the light stop scenario [20].

As we can see from the general expression of radiative corrections, Eq. (46), gauge coupling terms should tend to compensate top Yukawa coupling terms when $M_{1/2}, M_H > M_0$ ¹⁹. This is the case for the models analyzed in Figs. 2-4 where the stability condition is satisfied for $M_{1/2}, M_H \gtrsim 3$ TeV. Finally this trend is broken for superheavy Higgsinos,

¹⁹ This kind of spectra, where the boundary conditions for gauginos are heavier than sfermions, can be found e.g. in minimal gauge mediation (although the considered boundary conditions in our example of Eq. (58) do not match those of minimal gauge mediation) with a largish number of messengers or in some extra dimensional mechanisms of supersymmetry breaking as gaugino mediation [21, 22].

as the $\mu = M_{1/2}$ case shown in Fig. 5. Here for very heavy gauginos, $M_{1/2} \gg 1$ TeV, the value of $M_H^2 = m_{H_{u,d}}^2 + \mu^2$ becomes very large and the condition $\mathcal{S} = 1$ is overshoot. Therefore the stability condition requires lighter gauginos than scalars as we can see in Fig. 5.

VIII. OUTLOOK AND CONCLUSIONS

In view of the growing experimental evidence against the existence of sub-TeV new physics, and having in mind the naturalness problem of electroweak interactions, it is interesting to study the matching of the SM with possible UV completions solving the grand hierarchy problem. In general we expect that the presence of heavy mass states, coupled to the Higgs sector in the UV theory, would contribute at the matching scale \mathcal{Q}_m as finite threshold corrections to the SM Higgs mass term: these finite contributions trigger the hierarchy problem and destabilize the electroweak minimum. Determining these corrections is then of utmost importance to quantify the stability/naturalness problem and at least to find the parameter conditions that the underlying (final) theory at the unification, or Planck, scale should eventually provide to solve it.

In this paper we have pursued this task for the simplest perturbative UV completion of the SM solving the hierarchy problem: the MSSM. Although we have focused on this model, the qualitative features presented in this paper are expected to hold in any perturbative UV theory aiming to solve the grand hierarchy problem. The key ingredient of our analysis is the one-loop effective potential improved by the renormalization group equations, and the renormalization-scale invariance of such a potential. As a battleground we have considered the Landau gauge and dimensional regularization in the \overline{MS} renormalization scheme (at this level of the calculation, and for scalar and fermion fields integrated out, equivalent to the \overline{DR}). With these choices we have obtained the one-loop matching between the SM and MSSM Higgs sectors at the multi-TeV matching scale \mathcal{Q}_m , where the large leading logarithms between the high (unification) scale \mathcal{M} and \mathcal{Q}_m have been resummed.

Motivated by hints from experimental data, the matching has been performed assuming a large mass hierarchy between SM and non-SM fields, which allows to work in the unbroken electroweak symmetry phase. After having integrated the heavy fields out, the final one-loop identification between the SM mass term, m_{SM}^2 , and the MSSM mass terms, m_I^2 (with $I = 1, 2, 3$), is given by Eq. (40) which can be written as

$$m_{SM}^2 = [-m_1^2 c_\beta^2 - m_2^2 s_\beta^2 + 2m_3^2 s_\beta c_\beta] (1 + 2\Delta Z_h) + \Delta m^2, \quad (62)$$

where the radiative contributions ΔZ_h and Δm^2 are respectively coming from the matching between the wave functions and quadratic interactions of the SM and the MSSM lightest CP-even Higgs. The main features of the matching are as follows:

- We have integrated out only heavy states from the MSSM in the effective potential.

We have *not* integrated out the heavy modes from the light (SM) states, as in the Wilsonian action, which would have created a cutoff in the low energy theory. In this way we can still integrate momenta in the low energy theory up to infinity and then keep on using dimensional regularization for them.

- Using the scale independence of the effective potential the matching scale \mathcal{Q}_m is completely arbitrary in the considered one-loop approximation. If all heavy masses are of similar order of magnitude (as in the high-scale supersymmetry scenario considered here), \mathcal{Q}_m can be arbitrarily fixed at some intermediate value around them to avoid large logarithms and the breaking of perturbation theory at higher-loop orders. If very different heavy scales are present (as e.g. in split supersymmetry [23]) then different matching processes should be subsequently applied (cf. e.g. Ref. [14]).
- The size of the squared mass parameter of the lightest-Higgs in the high energy theory, $m^2 = -m_1^2 c_\beta^2 - m_2^2 s_\beta^2 + 2m_3^2 s_\beta c_\beta$, is not physically meaningful as it is strongly scale dependent. Its dependence is mostly compensated by the one of $2\Delta Z_h$ and Δm^2 , so that the final one-loop scale dependence of the right hand side of Eq. (62) is as weak as that of the SM mass parameter (in particular the dependence including large masses appears only at two loops). The natural scale of the SM electroweak sector should hence be deduced considering the sum of m^2 and Δm^2 (the ΔZ_h effect is in general subleading), possibly at the scale \mathcal{Q}_m to avoid perturbative problems. In particular, once the magnitude of $m^2(\mathcal{Q}_m)$ is enforced to be of the order of the electroweak scale, the stability of the electroweak breaking conditions is guaranteed by $\mathcal{S} \equiv |\Delta m^2(\mathcal{Q}_m)/m_{LE}^2(\mathcal{Q}_m)| \lesssim 1$.

We now present a short list of results obtained in the present paper:

- When the non-SM fields are heavy and degenerate, the expression for Δm^2 at the matching scale \mathcal{Q}_m reproduces the result obtained by Veltman in the SM using dimensional regularization and extracting the “quadratic” divergence as the residue of the pole in $d = 2$ dimensions. Veltman then interpreted this result as the coefficient of the cutoff Λ^2 , while we can express it as the coefficient of the common supersymmetric mass squared. Our result is thus consistent with Veltman’s and puts solid grounds in the understanding of the SM as an effective theory below the MSSM. As it is well known, the vanishing of the Veltman coefficient can only be achieved at (super)Planckian scales.
- When the MSSM fields are heavy, and of the same order of magnitude but not fully degenerate, the stability of the electroweak minimum provides a generalization of Veltman’s result, which amounts to introducing different cutoffs for the different SM sectors, plus a modification that is negligible only for a strictly non-hierarchical heavy spectrum. In particular by assuming the masses of the heavy Higgses and/or gauginos to be larger than those of sfermions, one can easily achieve the vanishing of Δm^2 at sub-Planckian scales.

- We have analyzed several cases where the MSSM mass parameters are generated at high scale, and transmitted to the observable sector by means of some mediators between the hidden and observable sectors, and we have determined the parameter regions where $m^2(\mathcal{Q}_m) \sim (100 \text{ GeV})^2$. However also the requirement $\Delta m^2(\mathcal{Q}_m) \lesssim \mathcal{O}(100 \text{ GeV})^2$ should be imposed in order to achieve satisfactory electroweak breaking conditions. The parameter regions where also this latter requirement is realized turn out to provide a sort of generalized focus point conditions which include threshold effects.

Let us finally make some considerations about the physical significance of the (stability) regions where both $m_{LE}^2(\mathcal{Q}_{EW}) = \mathcal{O}(100 \text{ GeV})^2$ and $\mathcal{S} \lesssim 1$ conditions are satisfied:

- As we have already mentioned, these regions are obtained by integrating out the heavy fields at one-loop. The procedure is performed at a renormalization scale nearby the energy scale of the heavy masses. The result is thus not jeopardized by large logarithms and it has no relevant one-loop scale dependence.
- The stability regions do depend on the threshold contributions we obtain from the effective potential in a particular (Landau) gauge. Therefore, in general, we could expect a gauge dependence in our results. However as we have performed the matching in the unbroken electroweak symmetry phase and upon integrating out only scalars and fermions (neither gauge nor Goldstone bosons), it turns out that our computation of the thresholds is *gauge independent* within the considered approximations.
- The effective potential depends on the particular renormalization scheme, and we have worked it out in the \overline{MS} scheme which amounts to subtracting (to define the counter-term) the infinite term proportional to $\frac{2}{\epsilon} - \gamma_E + \log(4\pi) - \delta$ with $\delta = 0$. In the \overline{MS} scheme the finite term in the effective potential contributions coming from heavy fermions and scalars is proportional to the constant $C = 3/2$ [cf. Eq. (5)]. Subtracting a different infinite counter term (with $\delta \neq 0$, as e.g. in the MS renormalization scheme or a variant thereof) would then lead to a shift in the constant C as $C \rightarrow C + \delta$. Consequently for the degenerate and almost degenerate cases, the stability condition $\mathcal{S} = 0$ leading respectively to the exact and generalized Veltman conditions (cf. sections VIA and VIB with $r_{1/2} \simeq r_H \simeq 1$) would receive additional contributions. Therefore, the Veltman-like conditions we obtain from the MSSM arise *only* in the \overline{DR} scheme and are relevant for the matching with the SM effective theory in the \overline{MS} renormalization scheme.

To conclude, given a scheme of supersymmetry breaking involving soft terms above the TeV scale, the threshold corrections to the Higgs mass parameters play a determinant role in the stability of the electroweak breaking conditions and the masses of the Higgs fields. In the present paper we have determined these threshold corrections in some MSSM scenarios with no relevant hierarchy in the heavy mass spectrum, and looked for

soft-parameter relations that could alleviate the little hierarchy problem. Of course it is clear that a similar analysis can also be performed for any perturbative theory that UV completes the SM. Moreover an equivalent analysis can be done even if the low energy theory is itself some extension of the SM, as one in which there is an aligned extended Higgs sector [24] giving rise to e.g. a two Higgs doublet model. If the UV completion of the SM is not perturbative, as in the case of a composite Higgs, the calculation cannot rely on perturbation theory and different methods to evaluate threshold effects should be used [3]. In general, whatever the final UV completion of the SM is, we expect it could provide an answer to the question on why our electroweak vacuum is stable and insensitive to high scale physics.

Acknowledgments

We are grateful to Pietro Slavich for reading the first version of the manuscript and providing useful comments. The work of I.M. is supported by the research grant ‘‘Theoretical Astroparticle Physics’’ number 2012CPPYP7 under the program PRIN 2012 funded by the Ministero dell’Istruzione, Universita e della Ricerca (MIUR). The work of G.N. is supported by the German Science Foundation (DFG) within the Collaborative Research Center (SFB) 676 ‘‘Particles, Strings and the Early Universe’’. The work of M.Q. is partly supported by the Spanish Consolider-Ingenio 2010 Programme CPAN (CSD2007-00042), by CICYT-FEDER-FPA2011-25948, by the Severo Ochoa excellence program of MINECO under the grant SO-2012-0234, by Secretaria d’Universitats i Recerca del Departament d’Economia i Coneixement de la Generalitat de Catalunya under Grant 2014 SGR 1450, and partly done at IFT and ICTP-SAIFR under CNPq grant 405559/2013-5. We also acknowledge support from CERN, where this work was partly done.

Appendix A: RADIATIVE CORRECTIONS TO MASS PARAMETERS

In this appendix we will present a detailed calculation of the radiative contributions to the mass parameters Δm_1^2 , Δm_2^2 and Δm_3^2 in the MSSM arising from the different sectors of sfermions, charginos, neutralinos and the Higgs scalar sector.

1. SQUARKS AND SLEPTONS

The mass squared matrices for top and bottom squarks and for tau-sneutrinos and staus can be written as

$$M_t^2 = \begin{pmatrix} m_Q^2 + Y_t^2 h_2^2 + \Pi_{\tilde{u}_L} & Y_t(A_t h_2 - \mu h_1) \\ Y_t(A_t h_2 - \mu h_1) & m_U^2 + Y_t^2 h_2^2 + \Pi_{\tilde{u}_R} \end{pmatrix} \quad (\text{A1})$$

$$M_{\tilde{b}}^2 = \begin{pmatrix} m_Q^2 + Y_b^2 h_1^2 + \Pi_{\tilde{d}_L} & Y_b(A_b h_1 - \mu h_2) \\ Y_b(A_b h_1 - \mu h_2) & m_D^2 + Y_b^2 h_1^2 + \Pi_{\tilde{d}_R} \end{pmatrix} \quad (\text{A2})$$

$$M_{\tilde{\nu}_\tau}^2 = \begin{pmatrix} m_L^2 + \Pi_{\tilde{\nu}_L} & 0 \\ 0 & m_N^2 \end{pmatrix} \quad (\text{A3})$$

$$M_{\tilde{\tau}}^2 = \begin{pmatrix} m_L^2 + Y_\tau^2 h_1^2 + \Pi_{\tilde{e}_L} & Y_\tau(A_\tau h_1 - \mu h_2) \\ Y_\tau(A_\tau h_1 - \mu h_2) & m_E^2 + Y_\tau^2 h_1^2 + \Pi_{\tilde{e}_R} \end{pmatrix} \quad (\text{A4})$$

where

$$\Pi_{\tilde{f}} = \frac{1}{2} \left[T_{3_{\tilde{f}}} (g_Y^2 + g_2^2) - Q_{\tilde{f}} g_Y^2 \right] (h_1^2 - h_2^2) . \quad (\text{A5})$$

Using the expressions in Eq. (44) it is straightforward to find the corresponding contributions to $\Delta m_{1,2,3}^2$ as

$$\begin{aligned} \Delta m_1^2 = & \frac{2}{32\pi^2} \left\{ 3Y_b^2 [G(m_Q^2) + G(m_D^2)] + Y_\tau^2 [G(m_L^2) + G(m_E^2)] + \frac{3Y_t^2 \mu^2}{m_Q^2 - m_U^2} [G(m_Q^2) - G(m_U^2)] \right. \\ & + \frac{3Y_b^2 A_b^2}{m_Q^2 - m_D^2} [G(m_Q^2) - G(m_D^2)] + \frac{Y_\tau^2 A_\tau^2}{m_L^2 - m_E^2} [G(m_L^2) - G(m_E^2)] \\ & \left. - \frac{g_Y^2}{2} [G(m_Q^2) - 2G(m_U^2) + G(m_D^2) - G(m_L^2) + G(m_E^2)] \right\} , \end{aligned}$$

$$\begin{aligned} \Delta m_2^2 = & \frac{2}{32\pi^2} \left\{ 3Y_t^2 [G(m_Q^2) + G(m_U^2)] + \frac{3Y_t^2 A_t^2}{m_Q^2 - m_U^2} [G(m_Q^2) - G(m_U^2)] \right. \\ & + \frac{3Y_b^2 \mu^2}{m_Q^2 - m_D^2} [G(m_Q^2) - G(m_D^2)] + \frac{Y_\tau^2 \mu^2}{m_L^2 - m_E^2} [G(m_L^2) - G(m_E^2)] \\ & \left. + \frac{g_Y^2}{2} [G(m_Q^2) - 2G(m_U^2) + G(m_D^2) - G(m_L^2) + G(m_E^2)] \right\} , \end{aligned}$$

$$\begin{aligned} \Delta m_3^2 = & \frac{2}{32\pi^2} \left\{ \frac{3Y_t^2 \mu A_t}{m_Q^2 - m_U^2} [G(m_Q^2) - G(m_U^2)] + \frac{3Y_b^2 \mu A_b}{m_Q^2 - m_D^2} [G(m_Q^2) - G(m_D^2)] \right. \\ & \left. + \frac{Y_\tau^2 \mu A_\tau}{m_L^2 - m_E^2} [G(m_L^2) - G(m_E^2)] \right\} , \end{aligned}$$

with

$$G(m_X^2) \equiv m_X^2 \left[\log \frac{m_X^2}{Q_m^2} - 1 - \delta \right] . \quad (\text{A6})$$

Here the function G has been generalized to the subtraction schemes having $\mathcal{C} = 3/2 + \delta$ in the scalar contributions of Eq. (5). Eq. (A6) recovers Eq. (45) for the \overline{MS} (or \overline{DR}) subtraction scheme where $\delta = 0$.

If some of the masses are degenerate the following limit turns useful:

$$\lim_{y \rightarrow x} \frac{G(x^2) - G(y^2)}{x^2 - y^2} = \log \frac{x^2}{Q_m^2} - \delta . \quad (\text{A7})$$

2. CHARGINOS

The squared mass matrix for the charginos can be written as

$$M_{ch}^2 = \begin{pmatrix} M_2^2 + g_2^2 h_2^2 & g_2(M_2 h_1 + \mu h_2) \\ g_2(M_2 h_1 + \mu^* h_2) & \mu^2 + g_2^2 h_1^2 \end{pmatrix}. \quad (\text{A8})$$

We can then compute the corresponding contributions to $\Delta m_{1,2,3}^2$ as

$$\Delta m_1^2 = \Delta m_2^2 = -\frac{4}{32\pi^2} \frac{g_2^2}{M_2^2 - \mu^2} [M_2^2 G(M_2^2) - \mu^2 G(\mu^2)], \quad (\text{A9})$$

$$\Delta m_3^2 = \frac{4}{32\pi^2} \frac{g_2^2 \mu M_2}{M_2^2 - \mu^2} [G(M_2^2) - G(\mu^2)], \quad (\text{A10})$$

with G defined in Eq. (A6). In case of degenerate spectra the following limit is useful:

$$\lim_{y \rightarrow x} \frac{x^2 G(x^2) - y^2 G(y^2)}{x^2 - y^2} = x^2 \left(-1 - 2\delta + 2 \log \frac{x^2}{Q_m^2} \right) \quad (\text{A11})$$

with $\delta = 0$ in the \overline{MS} or \overline{DR} schemes.

3. NEUTRALINOS

The squared mass matrix for the neutralinos can be written as

$$M_{ne}^2 = \begin{pmatrix} M_1^2 + \frac{g_Y^2}{2}(h_1^2 + h_2^2) & -\frac{1}{2}g_Y g_2(h_1^2 + h_2^2) & -\frac{g_Y}{\sqrt{2}}(M_1 h_1 + \mu h_2) & \frac{g_Y}{\sqrt{2}}(M_1 h_2 + \mu h_1) \\ -\frac{1}{2}g_Y g_2(h_1^2 + h_2^2) & M_2^2 + \frac{g_2^2}{2}(h_1^2 + h_2^2) & \frac{g_2}{\sqrt{2}}(M_2 h_1 + \mu h_2) & -\frac{g_2}{\sqrt{2}}(M_2 h_2 + \mu h_1) \\ -\frac{g_Y}{\sqrt{2}}(M_1 h_1 + \mu h_2) & \frac{g_2}{\sqrt{2}}(M_2 h_1 + \mu h_2) & \mu^2 + \frac{h_1^2}{2}(g_Y^2 + g_2^2) & -\frac{1}{2}h_1 h_2(g_Y^2 + g_2^2) \\ \frac{g_Y}{\sqrt{2}}(M_1 h_2 + \mu h_1) & -\frac{g_2}{\sqrt{2}}(M_2 h_2 + \mu h_1) & -\frac{1}{2}h_1 h_2(g_Y^2 + g_2^2) & \mu^2 + \frac{h_2^2}{2}(g_Y^2 + g_2^2) \end{pmatrix}. \quad (\text{A12})$$

The derivatives of the mass eigenvalues with respect to the backgrounds $\phi_j \equiv (h_1^2, h_2^2, h_1 h_2)$ can be easily computed following the techniques introduced in Ref. [25]. The squared mass eigenvalues are given by the solutions of the equation defined by the characteristic polynomial

$$\det(M^\dagger M - \lambda) \equiv \sum_n c^{(n)}(\phi_j) \lambda^n = 0 \quad (\text{A13})$$

where the coefficients of the characteristic polynomial are functions of h_1^2 , h_2^2 and $h_1 h_2$. Differentiating (A13) with respect to ϕ_j we obtain the required expressions

$$\left. \frac{\partial \lambda}{\partial \phi_j} \right|_{h_i=0} = - \left. \frac{\sum_n \frac{\partial c^{(n)}(\phi_j)}{\partial \phi_j} \lambda^n}{\sum_n n c^{(n)}(\phi_j) \lambda^{n-1}} \right|_{h_i=0} \quad (\text{A14})$$

where on the right hand side λ denotes the squared mass eigenvalues.

The calculation of $\Delta m_{1,2,3}^2$ is now straightforward and gives

$$\begin{aligned} \Delta m_1^2 = \Delta m_2^2 &= -\frac{2}{32\pi^2} \left\{ \frac{g_Y^2}{M_1^2 - \mu^2} [M_1^2 G(M_1^2) - \mu^2 G(\mu^2)] \right. \\ &\quad \left. + \frac{g_2^2}{M_2^2 - \mu^2} [M_2^2 G(M_2^2) - \mu^2 G(\mu^2)] \right\} \\ \Delta m_3^2 &= \frac{2}{32\pi^2} \left\{ \frac{g_Y^2 M_1 \mu}{M_1^2 - \mu^2} [G(M_1^2) - G(\mu^2)] + \frac{g_2^2 M_2 \mu}{M_2^2 - \mu^2} [G(M_2^2) - G(\mu^2)] \right\}. \end{aligned} \quad (\text{A15})$$

In the case of equal masses we can use the limiting behavior in Eqs. (A7) and (A11).

4. HIGGS SCALAR SECTOR

The general tree level potential for the scalar sector is given by

$$\begin{aligned} V &= m_1^2 |H_1|^2 + m_2^2 |H_2|^2 + m_3^2 (H_1 \cdot H_2 + h.c.) + \frac{g_Z^2}{2} (|H_2|^2 - |H_1|^2)^2 \\ &\quad + \frac{g_2^2}{2} |H_2^+ H_1^{0*} + H_1^{-*} H_2^0|^2 \end{aligned} \quad (\text{A16})$$

where $H_1 \cdot H_2 = H_1^a \varepsilon_{ab} H_2^b$ with $\varepsilon_{12} = -1$. The squared mass matrices for scalars [in the basis $(\text{Re } H_1^0, \text{Re } H_2^0)$], pseudoscalars [in the basis $(\text{Im } H_1^0, \text{Im } H_2^0)$] and charged scalars [in the basis (H_2^+, H_1^{-*})] are

$$M_S^2 = \begin{pmatrix} m_1^2 - g_Z^2 (h_2^2 - 3h_1^2) & -m_3^2 - 2g_Z^2 h_1 h_2 \\ -m_3^2 - 2g_Z^2 h_1 h_2 & m_2^2 + g_Z^2 (3h_2^2 - h_1^2) \end{pmatrix} \quad (\text{A17})$$

$$M_P^2 = \begin{pmatrix} m_1^2 - g_Z^2 (h_2^2 - h_1^2) & m_3^2 \\ m_3^2 & m_2^2 + g_Z^2 (h_2^2 - h_1^2) \end{pmatrix} \quad (\text{A18})$$

$$M_C^2 = \begin{pmatrix} m_2^2 + g_Z^2 (h_2^2 - h_1^2) + \frac{g_2^2}{2} h_1^2 & m_3^2 + \frac{g_2^2}{2} h_1 h_2 \\ m_3^2 + \frac{g_2^2}{2} h_1 h_2 & m_1^2 - g_Z^2 (h_2^2 - h_1^2) + \frac{g_2^2}{2} h_2^2 \end{pmatrix}. \quad (\text{A19})$$

For each of the previous matrices there are two eigenstates in the unbroken phase ($h_i = 0$), $H_{SM} = c_\beta H_1 + s_\beta \tilde{H}_2^*$ with mass eigenvalue $-m^2$, and $H_h = -s_\beta H_1 + c_\beta \tilde{H}_2^*$ with mass eigenvalue $m_H^2 = m_1^2 + m_2^2 + m^2$. The doublet H_{SM} is identified with the SM Higgs doublet, and we then exclude its contribution to Δm^2 . Using Eq. (A14) one obtains that

the heavy Higgs contributions are given by

$$\begin{aligned}\Delta m_1^2 &= \frac{1}{32\pi^2} \frac{G(m_H^2)}{m_1^2 + m_2^2 + 2m^2} [2g_Z^2(3m_1^2 - 2m_2^2 + m^2) + g_2^2(m_2^2 + m^2)] \\ \Delta m_2^2 &= \frac{1}{32\pi^2} \frac{G(m_H^2)}{m_1^2 + m_2^2 + 2m^2} [2g_Z^2(3m_2^2 - 2m_1^2 + m^2) + g_2^2(m_1^2 + m^2)] \\ \Delta m_3^2 &= \frac{1}{32\pi^2} \frac{G(m_H^2)}{m_1^2 + m_2^2 + 2m^2} [-2g_Z^2 - g_2^2] m_3^2 .\end{aligned}\tag{A20}$$

Using now the matching conditions (31), Eqs. (A20) can be also written as

$$\begin{aligned}\Delta m_1^2 &= \frac{1}{32\pi^2} G(m_H^2) [-4g_Z^2 \cos 2\beta + 2g_Z^2 \sin^2 \beta + g_2^2 \cos^2 \beta] \\ \Delta m_2^2 &= \frac{1}{32\pi^2} G(m_H^2) [4g_Z^2 \cos 2\beta + 2g_Z^2 \cos^2 \beta + g_2^2 \sin^2 \beta] \\ \Delta m_3^2 &= \frac{1}{32\pi^2} G(m_H^2) [-2g_Z^2 - g_2^2] \sin \beta \cos \beta .\end{aligned}\tag{A21}$$

Appendix B: ONE-LOOP SCALE INVARIANCE OF THE MATCHING

In section III we showed that by construction the one-loop matching conditions obtained via one-loop RG-improved effective potentials are independent of the choice of \mathcal{Q}_m . Here we use this property to check our finding (46).

The matching condition (40) led to (cf. section III B)

$$m_{LE}^2(\mathcal{Q}_m) = m^2(\mathcal{Q}_m)[1 + 2\Delta Z_h(\mathcal{Q}_m)] + \Delta m^2 ,\tag{B1}$$

where [cf. Eqs. (38) and (39)]

$$m^2 = -m_1^2 \cos^2 \beta - m_2^2 \sin^2 \beta + m_3^2 \sin 2\beta ,\tag{B2}$$

$$\Delta m^2 = -\Delta m_1^2 \cos^2 \beta - \Delta m_2^2 \sin^2 \beta + \Delta m_3^2 \sin 2\beta ,\tag{B3}$$

$$\tan 2\beta = 2\tilde{m}_3^2/(\tilde{m}_2^2 - \tilde{m}_1^2) .\tag{B4}$$

The one-loop scale invariance imposes that in Eq. (B1) the one-loop \mathcal{Q}_m dependence of the right hand side is the same of the SM. Here we check this issue.

The total derivative of the right hand side (RHS) in Eq. (B1) with respect to $\log \mathcal{Q}_m$ is given by ²⁰

$$\beta_{RHS} = -\beta_{\tilde{m}_1^2} \cos^2 \beta - \beta_{\tilde{m}_2^2} \sin^2 \beta + \beta_{\tilde{m}_3^2} \sin 2\beta + m^2(1 + 2\gamma_h - 2\gamma_{h_{LE}}) ,\tag{B5}$$

with

$$\tilde{m}_I^2 = m_I^2 + \Delta m_I^2 = m_I^2 + \sum_r \Delta m_{I,r}^2 , \quad \beta_{\tilde{m}_I^2} = \beta_{m_I^2} + \beta_{\Delta m_I^2} \equiv \beta_{m_I^2} - \sum_r \Delta_I^r ,\tag{B6}$$

²⁰ Note that the contribution to β_{RHS} coming from derivatives of the angle $\beta(\mathcal{Q}_m)$ cancels out after imposing Eq. (B4).

where the index r runs over the heavy fields as in Eq. (42). The (one-loop) contribution Δm_I^r can hence be deduced from the one-loop \mathcal{Q} dependence of $\Delta m_{I,r}^2$. Each quantity $\Delta m_{I,r}^2$ is provided in appendix A and can be decomposed into its logarithmic and non-logarithmic contributions as follows:

$$\Delta m_{I,r}^2 = \Delta_\ell^r m_I^2 + \Delta_f^r m_I^2 \equiv \frac{1}{32\pi^2} \ell_{I,r} + \frac{1}{32\pi^2} f_{I,r}. \quad (\text{B7})$$

In particular $\ell_{I,r} = 32\pi^2 \Delta m_{I,r}^2(G \rightarrow \tilde{G})$, with $\tilde{G}(x^2) \equiv x^2 \log(x^2/\mathcal{Q}^2)$, represents all terms proportional to logarithms of squared masses over \mathcal{Q}_m^2 , and $f_{I,r} = 32\pi^2 \Delta m_{I,r}^2(G \rightarrow \bar{G})$ with $\bar{G}(x^2) \equiv -x^2(1 + \delta)$ contains all non-logarithmic terms ($\delta = 0$ in the \overline{MS} and \overline{DR}). At one loop each $f_{I,r}$ is thus independent of \mathcal{Q}_m , and it results $\Delta_I^r = \partial \Delta m_{I,r}^2 / \partial \mathcal{Q}$.

The values of $\sum_r \Delta_I^r$ are determined in steps considering the contributions computed in appendix A for each sector. It comes out from sfermions

$$\begin{aligned} (16\pi^2) \Delta_1^{\tilde{f}} &= 6Y_b^2(m_Q^2 + m_D^2 + A_b^2) + 2Y_\tau^2(m_L^2 + m_E^2 + A_\tau^2) + 6Y_t^2 \mu^2 \\ &\quad - \frac{3}{5} g_1^2(m_Q^2 - 2m_U^2 + m_D^2 - m_L^2 + m_E^2), \\ (16\pi^2) \Delta_2^{\tilde{f}} &= 6Y_t^2(m_Q^2 + m_U^2 + A_t^2) + 6Y_b^2 \mu^2 + 2h_\tau^2 \mu^2 \\ &\quad + \frac{3}{5} g_1^2(m_Q^2 - 2m_U^2 + m_D^2 - m_L^2 + m_E^2), \\ (16\pi^2) \Delta_3^{\tilde{f}} &= 6Y_t^2 \mu A_t + 6Y_b^2 \mu A_b + 2Y_\tau^2 \mu A_\tau, \end{aligned} \quad (\text{B8})$$

from charginos

$$\begin{aligned} (16\pi^2) \Delta_1^{\tilde{\chi}^\pm} &= \Delta_2^{\tilde{\chi}^\pm} = -4g_2^2(M_2^2 + \mu^2), \\ (16\pi^2) \Delta_3^{\tilde{\chi}^\pm} &= 4g_2^2 \mu M_2, \end{aligned} \quad (\text{B9})$$

from neutralinos

$$\begin{aligned} (16\pi^2) \Delta_1^{\tilde{\chi}^0} &= \Delta_2^{\tilde{\chi}^0} = -2g_2^2(M_2^2 + \mu^2) - \frac{6}{5} g_1^2(M_1^2 + \mu^2), \\ (16\pi^2) \Delta_3^{\tilde{\chi}^0} &= 2g_2^2 M_2 \mu + \frac{6}{5} g_1^2 M_1 \mu, \end{aligned} \quad (\text{B10})$$

and from the scalar, pseudoscalar and charged Higgs sector

$$\begin{aligned} (16\pi^2) \Delta_1^H &= \frac{m_1^2 + m_2^2 + m^2}{m_1^2 + m_2^2 + 2m^2} [2g_Z^2(3m_1^2 - 2m_2^2 + m^2) + g_2^2(m_2^2 + m^2)], \\ (16\pi^2) \Delta_2^H &= \frac{m_1^2 + m_2^2 + m^2}{m_1^2 + m_2^2 + 2m^2} [2g_Z^2(3m_2^2 - 2m_1^2 + m^2) + g_2^2(m_1^2 + m^2)], \\ (16\pi^2) \Delta_3^H &= -\frac{m_1^2 + m_2^2 + m^2}{m_1^2 + m_2^2 + 2m^2} [2g_Z^2 + g_2^2] m_3^2. \end{aligned} \quad (\text{B11})$$

Now using Eq. (B6) and the MSSM one-loop β -functions [26]

$$\begin{aligned}
(16\pi^2)\beta_{m_1^2} &= 6Y_b^2(m_Q^2 + m_D^2 + m_1^2 + A_b^2) + 2Y_\tau^2(m_L^2 + m_E^2 + m_1^2 + A_\tau^2) \\
&\quad + (6Y_t^2 - 6g_2^2 - \frac{6}{5}g_1^2)\mu^2 - 6g_2^2M_2^2 - \frac{6}{5}g_1^2M_1^2 - \frac{3}{5}g_1^2S , \\
(16\pi^2)\beta_{m_2^2} &= 6Y_t^2(m_Q^2 + m_U^2 + m_2^2 + A_t^2) \\
&\quad + (6Y_b^2 + 2Y_\tau^2 - 6g_2^2 - \frac{6}{5}g_1^2)\mu^2 - 6g_2^2M_2^2 - \frac{6}{5}g_1^2M_1^2 + \frac{3}{5}g_1^2S , \\
(16\pi^2)\beta_{m_3^2} &= (3Y_t^2 + 3Y_b^2 + Y_\tau^2 - 3g_2^2 - \frac{3}{5}g_1^2)m_3^2 \\
&\quad + (6Y_t^2A_t^2 + 6Y_b^2A_b^2 + 2Y_\tau^2A_\tau^2 + 6g_2^2M_2 + \frac{6}{5}g_1^2M_1)\mu , \tag{B12}
\end{aligned}$$

where $S = m_Q^2 - 2m_U^2 + m_D^2 - m_L^2 + m_E^2 + m_2^2 - m_1^2$, we can write

$$\begin{aligned}
(16\pi^2)\beta_{\tilde{m}_1^2} &= (6Y_b^2 + 2Y_\tau^2 - 2g_Z^2 - g_2^2)m_1^2 - m^2[(2g_Z^2 + g_2^2)\sin^2\beta + 6g_Z^2\cos 2\beta] \\
(16\pi^2)\beta_{\tilde{m}_2^2} &= (6Y_t^2 - 2g_Z^2 - g_2^2)m_2^2 - m^2[(2g_Z^2 + g_2^2)\cos^2\beta - 6g_Z^2\cos 2\beta] \\
(16\pi^2)\beta_{\tilde{m}_3^2} &= (3Y_t^2 + 3Y_b^2 + Y_\tau^2 - 2g_Z^2 - g_2^2)m_3^2 - m^2(2g_Z^2 + g_2^2)\sin\beta\cos\beta . \tag{B13}
\end{aligned}$$

Plugging these expressions in Eq. (B5) yields

$$\begin{aligned}
\beta_{RHS} &= 6Y_t^2(\sin\beta\cos\beta m_3^2 - \sin^2\beta m_2^2) + (6Y_b^2 + 2Y_\tau^2)(\sin\beta\cos\beta m_3^2 - \cos^2\beta m_1^2) \\
&\quad - (2g_Z^2 + g_2^2)(-\cos^2\beta m_1^2 - \sin^2\beta m_2^2 + \sin 2\beta m_3^2) + (6g_Z^2c_{2\beta}^2 + 2\Delta\gamma)m^2 , \tag{B14}
\end{aligned}$$

where $\Delta\gamma = \gamma_h - \gamma_{LE} = \gamma_1\cos^2\beta + \gamma_2\sin^2\beta - \gamma_{hLE} = -\frac{3}{2}(\frac{1}{5}g_1^2 + g_2^2)$. (This difference between γ_h and γ_{LE} is caused by the heavy electroweakinos.) Finally one imposes the tree-level matching conditions of Eqs. (31) and (33), and obtains

$$\beta_{RHS} = \left(6y_t^2 + 6y_b^2 + 2y_\tau^2 + 6\lambda - \frac{9}{10}g_1^2 - \frac{9}{2}g_2^2 \right) m^2 . \tag{B15}$$

Eq. (B15) thus agrees with the β function of the quadratic term in the SM [27]. Note that in Eq. (B15) the use of tree-level relations within the one-loop β functions is justified by the fact that we are working at one-loop order.

Before concluding, notice that the splitting (B7) turns out to be useful also for Δm^2 . In such a case it reads as

$$\begin{aligned}
\Delta m^2 &= \Delta_\ell m^2 + \Delta_f m^2 , \\
-\Delta_\ell m^2 &= \Delta_\ell m_1^2 \cos^2\beta + \Delta_\ell m_2^2 \sin^2\beta - \Delta_\ell m_3^2 \sin 2\beta , \\
-\Delta_f m^2 &= \Delta_f m_1^2 \cos^2\beta + \Delta_f m_2^2 \sin^2\beta - \Delta_f m_3^2 \sin 2\beta \tag{B16}
\end{aligned}$$

although, as we already noticed, the splitting into a logarithmic (or one-loop scale dependent) and non-logarithmic (or one-loop scale independent) is not uniquely defined since

e.g. in the function $[G(x^2) - G(y^2)]/(x^2 - y^2)$ the splitting process does not commute with the limit $x \rightarrow y$ (cf. Eq. (A7)).

-
- [1] S. L. Glashow, Nucl. Phys. **22** (1961) 579; F. Englert and R. Brout, Phys. Rev. Lett. **13** (1964) 321; P. W. Higgs, Phys. Rev. Lett. **13** (1964) 508; S. Weinberg, Phys. Rev. Lett. **19** (1967) 1264; A. Salam (1968), N. Svartholm ed., "Elementary Particle Physics: Relativistic Groups and Analyticity", Eighth Nobel Symposium, p. 367.
- [2] E. Gildener and S. Weinberg, Phys. Rev. D **13** (1976) 3333; E. Gildener, Phys. Rev. D **14** (1976) 1667.
- [3] G. Marques Tavares, M. Schmaltz and W. Skiba, Phys. Rev. D **89** (2014) 1, 015009 [arXiv:1308.0025 [hep-ph]].
- [4] H. P. Nilles, Phys. Rept. **110** (1984) 1.
- [5] P. Draper, G. Lee and C. E. M. Wagner, Phys. Rev. D **89** (2014) 5, 055023 [arXiv:1312.5743 [hep-ph]]; E. Bagnaschi, G. F. Giudice, P. Slavich and A. Strumia, JHEP **1409** (2014) 092 [arXiv:1407.4081 [hep-ph]].
- [6] H. Baer, V. Barger, P. Huang, D. Mickelson, A. Mustafayev and X. Tata, Phys. Rev. D **87** (2013) 11, 115028 [arXiv:1212.2655 [hep-ph]].
- [7] M. Sher, Phys. Lett. B **317** (1993) 159 [Addendum-ibid. B **331** (1994) 448] [hep-ph/9307342]; G. Altarelli and G. Isidori, Phys. Lett. B **337** (1994) 141; J. A. Casas, J. R. Espinosa and M. Quiros, Phys. Lett. B **342** (1995) 171 [hep-ph/9409458]; J. R. Espinosa and M. Quiros, Phys. Lett. B **353** (1995) 257 [hep-ph/9504241]; J. A. Casas, J. R. Espinosa and M. Quiros, Phys. Lett. B **382** (1996) 374 [hep-ph/9603227]; G. Isidori, G. Ridolfi and A. Strumia, Nucl. Phys. B **609** (2001) 387 [hep-ph/0104016]; G. Degrandi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori and A. Strumia, JHEP **1208** (2012) 098 [arXiv:1205.6497 [hep-ph]]; D. Buttazzo, G. Degrandi, P. P. Giardino, G. F. Giudice, F. Sala, A. Salvio and A. Strumia, JHEP **1312** (2013) 089 [arXiv:1307.3536 [hep-ph]]; V. Branchina and E. Messina, Phys. Rev. Lett. **111** (2013) 241801 [arXiv:1307.5193 [hep-ph]]; V. Branchina, E. Messina and M. Sher, Phys. Rev. D **91** (2015) 013003 [arXiv:1408.5302 [hep-ph]]; J. R. Espinosa, G. F. Giudice, E. Morgante, A. Riotto, L. Senatore, A. Strumia and N. Tetradis, arXiv:1505.04825 [hep-ph].

- [8] M. J. G. Veltman, *Acta Phys. Polon. B* **12** (1981) 437.
- [9] C. Ford, D. R. T. Jones, P. W. Stephenson and M. B. Einhorn, *Nucl. Phys. B* **395** (1993) 17 [hep-lat/9210033].
- [10] S. R. Coleman and E. J. Weinberg, *Phys. Rev. D* **7** (1973) 1888.
- [11] M. Bando, T. Kugo, N. Maekawa and H. Nakano, *Prog. Theor. Phys.* **90** (1993) 405 [hep-ph/9210229]; H. Nakano and Y. Yoshida, *Phys. Rev. D* **49** (1994) 5393 [hep-ph/9309215]; C. Ford, hep-th/9609127; C. Ford and C. Wiesendanger, *Phys. Lett. B* **398** (1997) 342 [hep-th/9612193]; J. A. Casas, V. Di Clemente and M. Quiros, *Nucl. Phys. B* **553** (1999) 511 [hep-ph/9809275]; J. A. Casas, V. Di Clemente and M. Quiros, *Nucl. Phys. B* **581** (2000) 61 [hep-ph/0002205].
- [12] J. L. Feng, K. T. Matchev and T. Moroi, *Phys. Rev. D* **61** (2000) 075005 [hep-ph/9909334]; J. L. Feng, K. T. Matchev and D. Sanford, *Phys. Rev. D* **85** (2012) 075007 [arXiv:1112.3021 [hep-ph]].
- [13] C. E. M. Wagner, *Nucl. Phys. B* **528** (1998) 3 [hep-ph/9801376]; A. Delgado, M. Quiros and C. Wagner, *JHEP* **1404** (2014) 093 [arXiv:1402.1735 [hep-ph]].
- [14] M. Carena, G. Nardini, M. Quiros and C. E. M. Wagner, *JHEP* **0810** (2008) 062 [arXiv:0806.4297 [hep-ph]].
- [15] M. B. Einhorn and D. R. T. Jones, *Phys. Rev. D* **46** (1992) 5206.
- [16] I. Masina and M. Quiros, *Phys. Rev. D* **88** (2013) 093003 [arXiv:1308.1242 [hep-ph]].
- [17] I. Masina, *Phys. Rev. D* **87** (2013) 053001 [arXiv:1209.0393 [hep-ph]].
- [18] S. S. AbdusSalam, B. C. Allanach, H. K. Dreiner, J. Ellis, U. Ellwanger, J. Gunion, S. Heinemeyer and M. Kraemer *et al.*, *Eur. Phys. J. C* **71** (2011) 1835 [arXiv:1109.3859 [hep-ph]].
- [19] A. Delgado, M. Garcia and M. Quiros, *Phys. Rev. D* **90** (2014) 1, 015016 [arXiv:1312.3235 [hep-ph]].
- [20] A. Delgado, G. Nardini and M. Quiros, *JHEP* **1204** (2012) 137 [arXiv:1201.5164 [hep-ph]]; A. Delgado, M. Quiros and C. Wagner, *Phys. Rev. D* **90** (2014) 3, 035011 [arXiv:1406.2027 [hep-ph]].
- [21] G. F. Giudice and R. Rattazzi, *Phys. Rept.* **322** (1999) 419 [hep-ph/9801271].
- [22] I. Antoniadis, S. Dimopoulos, A. Pomarol and M. Quiros, *Nucl. Phys. B* **544** (1999) 503 [hep-ph/9810410]; A. Delgado, A. Pomarol and M. Quiros, *Phys. Rev. D* **60** (1999) 095008 [hep-ph/9812489]; R. Barbieri, L. J. Hall and Y. Nomura, *Phys. Rev. D* **63** (2001)

- 105007 [hep-ph/0011311]; A. Delgado and M. Quiros, Nucl. Phys. B **607** (2001) 99 [hep-ph/0103058]; A. Delgado, G. von Gersdorff and M. Quiros, Nucl. Phys. B **613** (2001) 49 [hep-ph/0107233]; M. Quiros, hep-ph/0302189; S. Dimopoulos, K. Howe and J. March-Russell, Phys. Rev. Lett. **113** (2014) 111802 [arXiv:1404.7554 [hep-ph]].
- [23] G. F. Giudice and A. Romanino, Nucl. Phys. B **699** (2004) 65 [Erratum-ibid. B **706** (2005) 65] [hep-ph/0406088]; N. Arkani-Hamed, S. Dimopoulos, G. F. Giudice and A. Romanino, Nucl. Phys. B **709** (2005) 3 [hep-ph/0409232].
- [24] J. F. Gunion and H. E. Haber, Phys. Rev. D **67** (2003) 075019 [hep-ph/0207010]; A. Delgado, G. Nardini and M. Quiros, JHEP **1307** (2013) 054 [arXiv:1303.0800 [hep-ph]]; M. Carena, I. Low, N. R. Shah and C. E. M. Wagner, JHEP **1404** (2014) 015 [arXiv:1310.2248 [hep-ph]].
- [25] T. Ibrahim and P. Nath, Phys. Rev. D **66** (2002) 015005 [hep-ph/0204092].
- [26] D. J. Castano, E. J. Piard and P. Ramond, Phys. Rev. D **49** (1994) 4882 [hep-ph/9308335]; S. P. Martin and M. T. Vaughn, Phys. Rev. D **50** (1994) 2282 [Erratum-ibid. D **78** (2008) 039903] [hep-ph/9311340].
- [27] M. x. Luo and Y. Xiao, Phys. Rev. Lett. **90** (2003) 011601 [hep-ph/0207271].