

## Recommendations for PDF usage in LHC predictions

A. Accardi<sup>a,b</sup>, S. Alekhin<sup>c,d</sup>, J. Blümlein<sup>e</sup>, M.V. Garzelli<sup>c</sup>, K. Lipka<sup>f</sup>,  
W. Melnitchouk<sup>b</sup>, S. Moch<sup>c</sup>, R. Plačákytė<sup>f</sup>, J.F. Owens<sup>g</sup>, E. Reya<sup>h</sup>, N. Sato<sup>b</sup>, A. Vogt<sup>i</sup>  
and O. Zenaiev<sup>f</sup>

<sup>a</sup> *Hampton University, Hampton, VA 23668, USA*

<sup>b</sup> *Jefferson Lab, Newport News, VA 23606, USA*

<sup>c</sup> *II. Institut für Theoretische Physik, Universität Hamburg  
Luruper Chaussee 149, D-22761 Hamburg, Germany*

<sup>d</sup> *Institute for High Energy Physics  
142281 Protvino, Moscow region, Russia*

<sup>e</sup> *Deutsches Elektronensynchrotron DESY  
Platanenallee 6, D-15738 Zeuthen, Germany*

<sup>f</sup> *Deutsches Elektronensynchrotron DESY  
Notkestraße 85, D-22607 Hamburg, Germany*

<sup>g</sup> *Florida State University, Tallahassee, FL 32306, USA*

<sup>h</sup> *Institut für Physik, Technische Universität Dortmund  
D-44221 Dortmund, Germany*

<sup>i</sup> *Department of Mathematical Sciences, University of Liverpool  
Liverpool L69 3BX, United Kingdom*

### Abstract:

We review the present status of the determination of parton distribution functions (PDFs) in the light of the precision requirements for the LHC in Run 2 and other future hadron colliders. We provide brief reviews of all currently available PDF sets and use them to compute cross sections for a number of benchmark processes, including Higgs boson production in gluon-gluon fusion at the LHC. We show that the differences in the predictions obtained with the various PDFs are due to particular theory assumptions made in the fits of those PDFs. We discuss PDF uncertainties in the kinematic region covered by the LHC and on averaging procedures for PDFs, such as advocated by the PDF4LHC15 sets, and provide recommendations for the usage of PDF sets for theory predictions at the LHC.

## Contents

<b>1. Introduction</b>	2
<b>2. Data sets and results for PDF fits</b>	3
2.1. Data sets used in PDF fits	3
2.2. Results for PDFs	5
<b>3. Theory for PDF fits</b>	13
3.1. Theory for analyses of DIS data	13
3.1.1. Massless PDFs	15
3.1.2. Heavy-quark structure functions	16
3.2. Heavy-flavor PDFs	17
3.3. Heavy-quarks schemes	18
3.3.1. Variable-flavor number schemes	18
3.3.2. Validation with DIS charm-quark production	22
3.3.3. Charm-quark mass	23
3.4. Light-flavor PDFs	25
3.4.1. Up- and down-quark distributions	25
3.4.2. Strange-quark distribution	28
3.5. Nuclear corrections	28
3.6. Software and tools	30
<b>4. Strong coupling constant</b>	31
<b>5. Cross section predictions for the LHC</b>	34
5.1. Higgs boson production	34
5.2. Hadro-production of heavy quarks	38
5.2.1. Top-quark hadro-production: inclusive cross section	38
5.2.2. Top-quark hadro-production: differential distributions	41
5.2.3. Bottom-quark hadro-production	42
5.2.4. Charm-quark hadro-production	43
5.3. $W'/Z'$ production	46
<b>6. Recommendations for PDF usage</b>	48
6.1. The 2015 PDF4LHC recommendations: A critical appraisal	49
6.2. New recommendations for PDF usage at the LHC	50
<b>7. Conclusion</b>	51
Acknowledgments	52
<b>References</b>	52

## 1. INTRODUCTION

In Run 2 of the Large Hadron Collider (LHC), the very details of the Standard Model (SM), including cross sections of different processes and Higgs bosons properties, are being measured with very high precision. At the same time, the new data at the highest center-of-mass collision energies ever achieved ( $\sqrt{s} = 13$  TeV) are used to search for physics phenomena beyond the SM (BSM). The experimental data used to perform those measurements are generally expected to have percent-level accuracy, depending on details such as the final states and the acceptance and efficiency of the detectors in particular kinematics ranges.

To further test the SM and to identify signals for new physics, measurements need to be compared to precise theoretical predictions, which need to incorporate higher order radiative corrections in Quantum Chromodynamics (QCD) and, possibly, the electroweak sector of the SM. In order to reach the benchmark precision set by the accuracy of the experimental data, next-to-next-to-leading order (NNLO) corrections in QCD are often required. At next-to-leading order (NLO) in QCD, the residual theoretical uncertainty from truncating the perturbative expansion commonly estimated by variations of the renormalization and factorization scales  $\mu_r$  and  $\mu_f$  are often too large compared to the experimental accuracy. Nonetheless, for observables with complex final states, and indeed for many BSM signals, one must still contend with NLO calculations, which will continue to require corresponding NLO fits.

Parton distribution functions (PDFs) in the proton serve as an essential input for any cross section prediction at hadron colliders and have been measured with increasing precision over the last three decades. Likewise, the strong coupling constant  $\alpha_s(M_Z)$  at the  $Z$  boson mass scale  $M_Z$  and the masses  $m_h$  of the heavy quarks  $h = c, b, t$  are well constrained by existing data and their determination is accurate at least to NNLO. However, despite steady improvements in the accuracy of PDF determinations over the years, the uncertainties associated with PDFs, the strong coupling  $\alpha_s(M_Z)$ , and quark masses still dominate many calculations of cross sections for SM processes at the LHC. A particularly prominent example is the cross section for the production of a SM Higgs boson in the gluon-gluon fusion channel.

The currently available PDF sets are CJ15 [1], accurate to NLO in QCD, as well as ABM12 [2], CT14 [3], HERAPDF2.0 [4], JR14 [5], MMHT14 [6], and NNPDF3.0 [7] to NNLO in QCD. These provide a detailed description of the parton content of the proton, which depends on the chosen sets of experimental data as well as on the theory assumptions and the underlying physics models used in the analyses. Both theoretical and experimental inputs have direct impact on the obtained nonperturbative parameters, namely, the fitted PDFs, the value of  $\alpha_s(M_Z)$  and the quark masses. Moreover, they can lead to large systematic shifts compared to the uncertainties of the experimental data used in the fit. For precision predictions in Run 2 of the LHC it is therefore very important to quantify those effects in detailed validations of the individual PDF sets in order to reduce the uncertainties in those nonperturbative input parameters. Moreover, this will allow one to pinpoint problems with the determination of certain PDFs. Any approach to determine the parton luminosities at the LHC which implies mixing or averaging of various PDFs or of their respective uncertainties, such as that advocated in the recent PDF4LHC recommendations [8], is therefore potentially dangerous in the context of precision measurements, in particular, or when studying processes at kinematic edges such as at large values of Bjorken  $x$  or small scales  $Q^2$ .

In this paper we briefly discuss the available world data used to constrain PDFs in Sec. 2 and stress the need to include only compatible data sets in any analysis. The data analysis relies on comparison with precise theoretical predictions, with many of these implemented in software tools. In this respect, we underline in Sec. 3 the importance of open-source code to provide

PDF sets	$\Delta\chi^2$ criterion	data sets used in analysis
ABM12 [2]	1	incl. DIS, DIS charm, DY
CJ15 [1]	100	incl. DIS, DY (incl. $p\bar{p} \rightarrow W^\pm X$ ), $p\bar{p}$ jets, $\gamma$ +jet
CT14 [3]	100	incl. DIS, DIS charm, DY, $p\bar{p}$ jets, $pp$ jets
HERAPDF2.0 [4]	1	incl. DIS, DIS charm, DIS jets
JR14 [5]	1	incl. DIS, DIS charm, DY, $p\bar{p}$ jets, DIS jets
MMHT14 [9]	2.3 ... 42.3 (dynamical)	incl. DIS, DIS charm, DY, $p\bar{p}$ jets, $pp$ jets, $t\bar{t}$
NNPDF3.0 [7]	1	incl. DIS, DIS charm, DY, $p\bar{p}$ jets, $pp$ jets, $t\bar{t}$ , $W$ + charm

TABLE 1: Summary of major data sets used in the various PDF analyses and the confidence level criteria employed. Detailed references to the data sets are given in Refs. [1–5, 7, 9].

benchmarks and to facilitate theory improvements through indication and reduction of possible errors. In addition, Sec. 3 is devoted to a discussion of a number of crucial theory aspects in PDF fits. These include the treatment of heavy quarks and their masses, QCD corrections for  $W^\pm$ - and  $Z$ -boson production applied in the fit of light-flavor PDFs, and the importance of nuclear corrections in scattering data off nuclei. The strong coupling constant is correlated with the PDFs and is therefore an important parameter to be determined simultaneously with the PDFs. The state of the art is reviewed in Sec. 4. The need to address PDF uncertainties for cross section predictions is illustrated in Sec. 5, with the Higgs boson cross section in the gluon-gluon fusion channel being the most prominent case. Other examples include the production of heavy quarks at the LHC in different kinematical regimes. Our observations illustrate important shortcomings of the recent PDF4LHC recommendations [8] which are addressed in Sec. 6, where alternative recommendations for the usage of sets of PDFs for theory predictions at the LHC are provided. Finally, we conclude in Sec. 7.

## 2. DATA SETS AND RESULTS FOR PDF FITS

We begin with an overview of the currently available data which can be used to determine PDFs and present the fit results of the various groups.

### 2.1. Data sets used in PDF fits

The data used in the various PDF fits overlap to a large extent, as indicated in Tab. 1. However, there are also substantial differences which are related to the accuracy required in the analysis, the feasibility of efficiently implementing the corresponding theoretical computations, or the subjective evaluation of the data quality, to name a few.

The core of all PDF fits comprises the deep-inelastic scattering (DIS) data obtained at the HERA electron-proton ( $ep$ ) collider and in fixed-target experiments. While the former has used

only a proton target, the latter have collected large amounts of data for the deuteron and heavier targets as well. The analysis of nuclear-target data requires an accurate account of nuclear effects. This is challenging already in the case of the loosely-bound deuteron (cf. Sec. 3), and even more so for heavier targets. Therefore, in general, data sets for DIS on targets heavier than deuteron are not used. Nonetheless, different combinations of data sets for the neutrino-induced DIS off iron and lead targets obtained by the CCFR/NuTeV, CDHSW, and CHORUS experiments are included in the CT14, MMHT14, and NNPDF3.0 analyses, but are not used by other groups to avoid any influence of nuclear correction uncertainties. One can also point out the abnormal dependence of the DIS structure functions on the beam energy in the NuTeV experiment [10] and the poor agreement of the CDHSW data with the QCD predictions on the  $Q^2$  slope of structure functions [11–13] as an additional motivations to exclude these data sets.

The kinematic cuts applied to the commonly used DIS data also differ in various analyses in order to minimize the influence of higher twist contributions. Another important feature of the DIS data analyses in PDF fits concerns the use of data for the DIS structure function  $F_2$  instead of the data for the measured cross sections. These aspects will be discussed in Sec. 3.

The inclusive DIS data are often supplemented by the semi-inclusive data on the neutral-current and charged-current DIS charm-quark production. The neutral-current sample collected by the HERA experiments provides a valuable tool to study the heavy-quark production mechanism. This is vital for pinning down PDFs, in particular the gluon PDF at small  $x$ , relevant for important phenomenological applications at the LHC (cf. Sec. 5). The charged-current charm production data help to constrain the strange sea PDF, which is strongly mixed with contributions from non-strange PDFs in other observables (cf. Sec. 3).

The Drell-Yan (DY) data are also a necessary ingredient of any PDF analysis since DIS data alone do not allow for a comprehensive disentangling of the quark and anti-quark distributions. Historically, for a long time only fixed-target DY data were available for PDF fits. In particular, this did not allow for a model-independent separation of the valence and sea quarks at small- $x$ . The high precision DY data obtained in proton-proton ( $pp$ ) and proton-anti-proton ( $p\bar{p}$ ) collisions from the LHC and the Tevatron open new possibilities to study the PDFs at small and large  $x$ . The LHC experiments are quickly accumulating statistics and are currently providing data samples at  $\sqrt{s}=7$  and 8 TeV for  $W$ - and  $Z$ -boson production with typical luminosities of  $\sim 1 \text{ fb}^{-1}$ . The rapid progress in experimental measurements causes a greatly non-uniform coverage of the recent DY data in various PDF fits (cf. Tab. 2) and leads to corresponding differences in the accuracy of the extracted PDFs. Another issue here is the theoretical accuracy achieved for the description of the DY data. This varies substantially and will be discussed in Sec. 3.

Often, jet production in  $pp$  and  $p\bar{p}$  collisions is used as an additional process to constrain the large- $x$  gluon PDF. Here, the QCD corrections are known to NLO and the calculation of the NNLO ones is in progress [14]. The incomplete knowledge of the latter is problematic in view of a consistent PDF analysis at NNLO when including those jet data. This will be discussed in Sec. 4 in connection with the determination of the value of the strong coupling constant  $\alpha_s$ .

In addition to these major categories of data commonly used to constrain PDFs, some complementary processes are also employed in some cases, as indicated in Tab. 1. These comprise the hadro-production of top-quark pairs from  $pp$  and  $p\bar{p}$  collisions and the associated production  $W$  bosons with charm quarks in  $pp$  collisions. Sometimes, also jet production in  $ep$  collisions and prompt photon ( $\gamma$ +jet) production from  $pp$  and  $p\bar{p}$  collisions is considered. Except for  $t\bar{t}$  production the necessary QCD corrections are known to NLO only, so that the same arguments as in the case of jet hadro-production data apply, if those data are included in a fit at NNLO accuracy. For  $t\bar{t}$  production, only the inclusive cross section is considered at the moment in the available PDFs and

there is a significant correlation with PDFs, especially of the gluon PDF with the top-quark mass.

Taken together, the set of these data has a number of data points (NDP) of the order of few thousand, and provides sufficient information to describe the PDFs with an ansatz of about  $O(30)$  free parameters. The parameters can include the strong coupling constant  $\alpha_s(M_Z)$  and the heavy-quark masses  $m_c$ ,  $m_b$  and  $m_t$ , which are correlated with the PDFs, as will be discussed in Secs. 3 and 4. This provides sufficient flexibility for all PDF groups and it is routinely checked that no additional terms are required to improve the quality of fit. The exception is the NNPDF group, which typically uses  $O(250)$  free parameters in the neural network.

Apart from those considerations there is the general problem of the quality of the experimental data, that is to say whether or not the PDFs are extracted from a consistent data set. The various groups have different approaches, which roughly fall into two classes according to the different confidence level (c.l.) criteria for the value of  $\chi^2$  in the goodness-of-fit test. One approach is to fit to a very wide (or even the widest possible) set of data, while the other one rejects inconsistent data sets. In the former case, a tolerance criterion for  $\Delta\chi^2$  is introduced (e.g.  $\Delta\chi^2 = 100$ ), while the latter approach maintains that  $\Delta\chi^2 = 1$ . For the various PDF groups this information is listed in Tab. 1.

For further reference, we quote here the definition of  $\chi^2$  used in data comparisons. It is expressed as follows:

$$\chi^2 = \sum_i \frac{[\mu_i - m_i(1 - \sum_j \gamma_j^i b_j)]^2}{\delta_{i,\text{unc}}^2 m_i^2 + \delta_{i,\text{stat}}^2 \mu_i m_i} + \sum_j b_j^2 + \sum_i \ln \frac{\delta_{i,\text{unc}}^2 m_i^2 + \delta_{i,\text{stat}}^2 \mu_i m_i}{\delta_{i,\text{unc}}^2 \mu_i^2 + \delta_{i,\text{stat}}^2 \mu_i^2}, \quad (1)$$

where  $\mu_i$  represents the measurement at the point  $i$ ,  $m_i$  is the corresponding theoretical prediction and  $\delta_{i,\text{stat}}$ ,  $\delta_{i,\text{unc}}$  are the relative statistical and uncorrelated systematic uncertainties, respectively.  $\gamma_j^i$  denotes the sensitivity of the measurement to the correlated systematic source  $j$  and  $b_j$  their shifts, with a penalty term  $\sum_j b_j^2$  added. In addition, a logarithmic term is introduced arising from the likelihood transition to  $\chi^2$  when scaling of the errors is applied [15].

## 2.2. Results for PDFs

Before we start a detailed discussion of the theoretical aspects of the PDF determinations we would like to illustrate the present status of PDF sets at NNLO in QCD and discuss briefly some differences, which are clearly visible. The currently available sets at NNLO in QCD are shown in Figs. 1–6. The light-quark ( $u$ ,  $d$ ) valence PDFs together with the gluon and the quark sea distributions ( $x\Sigma = 2x(\bar{u} + \bar{c} + \bar{d} + \bar{s})$  for four active flavors) with the respective uncertainty bands are displayed in Figs. 1, 3 and 5 at the scales  $Q^2 = 4 \text{ GeV}^2$ ,  $100 \text{ GeV}^2$  and  $M_Z^2$  in the range  $10^{-4} \leq x \leq 1$  for the sets ABM12 [2], HERAPDF2.0 [4] and JR14 [5]. Likewise, Figs. 2, 4 and 6 show the sets CT14 [3], MMHT14 [6] and NNPDF3.0 [7].

The main features of the present NNLO PDFs in Figs. 1–6 in the main kinematic region of  $x$  and  $Q^2$  relevant for hard scattering events at Tevatron and the LHC can be characterized as follows. The agreement in the distributions  $xu_v$ , and to a slightly lesser extent  $\Sigma$ , is very good for ABM12, JR14 and HERAPDF2.0, as shown in Fig. 1. For the valence PDF  $xd_v$  there is also an overall reasonable agreement, but the distribution deviates by more than  $1\sigma$  at  $x \gtrsim 0.1$  in the case of HERAPDF2.0. One should note that  $xd_v$  is more difficult to measure in  $e^\pm p$  DIS at HERA than  $xu_v$  and additional constraints from deuteron data are important to fix the details of this PDF, as discussed in Sec. 3 below.




FIG. 1: The  $u$ -valence,  $d$ -valence, gluon and sea quark ( $x\Sigma = 2x(\bar{u} + \bar{c} + \bar{d} + \bar{s})$ ) PDFs with their uncertainty bands of ABM12 [2], HERAPDF2.0 [4] and JR14 (set JR14NNLO08VF) [5] at NNLO at the scale  $Q^2 = 4 \text{ GeV}^2$ .

The results on the gluon momentum distribution  $xg$  are clearly different at low values of  $x$ . Here, JR14 obtains the largest values, followed by ABM12 and HERAPDF2.0, with the latter displaying a valence-like shape below  $x = 10^{-3}$ . For CT14, MMHT14 and NNPDF3.0 there is very good agreement for  $xu_v$ , cf. Fig. 2. Some differences are visible in case of  $xd_v$ , where CT14 reports larger values than NNPDF3.0. The spread in  $\Sigma$  for the sets in Fig. 2 is much greater than those by ABM12, JR14 and HERAPDF2.0. This is true as well for the gluon PDF  $xg$  with a particularly wide error band for CT14. The latter may be related to the error treatment of CT14, which tolerates an enhanced value in  $\Delta\chi^2$ . The CT14 uncertainty band for the gluon PDF also




FIG. 2: Same as Fig. 1 for the CT14 [3], MMHT14 [6] and NNPDF3.0 [7] PDF sets at NNLO.

covers the predictions for the distributions by JR14, ABM12, and HERAPDF2.0.

The disagreement in  $x d_v$  between HERAPDF2.0 and ABM12 or JR14 persists through the evolution from  $Q^2 = 4 \text{ GeV}^2$  to  $Q^2 = M_Z^2$ , cf. Fig. 3 and 5. Likewise, the spread in  $x d_v$  between CT14, MMHT14 and NNPDF3.0 becomes more pronounced, as shown in Fig. 4 and 6. On the other hand, differences in the singlet PDFs  $\Sigma$  and  $xg$ , while still somewhat visible at  $Q^2 = 100 \text{ GeV}^2$ , largely wash out at scales  $Q^2 = M_Z^2$  which govern the physics of central rapidity events at the LHC. Those remaining differences persist at large scales (as in the case of the gluon PDFs at large  $x > 0.1$ ) and will have a significant impact. The crucial test for all PDF sets comes through a detailed comparison of cross section predictions to data. This will be discussed in the remainder of the paper, in particular in Secs. 3 and 5.






FIG. 3: Same as Fig. 1 at the scale  $Q^2 = 100 \text{ GeV}^2$  with the sea  $x\Sigma = 2x(\bar{u} + \bar{c} + \bar{d} + \bar{s} + \bar{b})$ .

Experiment	ATLAS		CMS		DØ		LHCb		
	7	8	7	8	1.96		7	8	8
$\sqrt{s}$ (TeV)	7	8	7	8	1.96		7	8	8
Final states	$W^+ \rightarrow l^+ \nu$	$W^+ \rightarrow \mu^+ \nu$	$W^+ \rightarrow \mu^+ \nu$	$W^+ \rightarrow \mu^+ \nu$	$W^+ \rightarrow \mu^+ \nu$	$W^+ \rightarrow e^+ \nu$	$W^+ \rightarrow \mu^+ \nu$	$Z \rightarrow e^+ e^-$	$W^+ \rightarrow \mu^+ \nu$
	$W^- \rightarrow l^- \nu$	$W^- \rightarrow \mu^- \nu$	$W^- \rightarrow \mu^- \nu$	$W^- \rightarrow \mu^- \nu$	$W^- \rightarrow \mu^- \nu$	$W^- \rightarrow e^- \nu$	$W^- \rightarrow \mu^- \nu$		$W^- \rightarrow \mu^- \nu$
	$Z \rightarrow l^+ l^-$						$Z \rightarrow \mu^+ \mu^-$		$Z \rightarrow \mu^+ \mu^-$
Cut on the lepton $P_T$	$P_T^l > 20$ GeV	$P_T^\mu > 25$ GeV	$P_T^\mu > 25$ GeV	$P_T^\mu > 25$ GeV	$P_T^\mu > 25$ GeV	$P_T^e > 25$ GeV	$P_T^\mu > 20$ GeV	$P_T^e > 20$ GeV	$P_T^e > 20$ GeV
Reference	[16]	[17]	[18]	[19]	[20]	[21]	[22]	[23]	[23]
<i>NDP</i>	30	11	22	10	13	31	17	34	
ABMP15 [24] <sup>a</sup>	29.8	22.5	–	16.9	18.0	44.1	18.2	–	–
CJ15 [1] <sup>b</sup>	–	–	–	20	29	–	–	–	–
CT14 [3]	42	– <sup>c</sup>	–	–	34.7	–	–	–	–
JR14 [5]	–	–	–	–	–	–	–	–	–
HERAFitter [28]	–	–	–	13	19	–	–	–	–
MMHT14 [6]	39	–	–	21	–	–	–	–	–
NNPDF3.0 [7]	35.4	18.9	–	–	–	–	–	–	–

<sup>a</sup> The ABM12 [2] analysis has used data from ATLAS [16] and older data sets from CMS and LHCb which are superseded and not listed here.

<sup>b</sup> The CJ15 fit includes also  $l\nu$  data from CDF [25] ( $\chi^2/\text{DNP}=12/11$ ), as well as the reconstructed  $W^\pm$  asymmetry data from CDF [26] ( $\chi^2/\text{DNP}=16/13$ ) and DØ [27] ( $\chi^2/\text{DNP}=14/14$ ).

<sup>c</sup> Statistically less significant data with the cut of  $P_T^\mu > 35$  GeV are used.

TABLE 2: Compilation of precise data on  $W^-$  and  $Z$ -boson production in  $pp$  and  $p\bar{p}$  collisions and the  $\chi^2$  values per number of data points (NDP) obtained for these data sets in different PDF analyses. Missing table entries indicate that the respective data sets has not been used in the analysis. For data on the reconstructed  $W^\pm$ -boson charge asymmetry [27], see [1, 24]. Other data sets of lower accuracy, which have become obsolete and data sets superseded are not listed here.




FIG. 4: Same as Fig. 2 at the scale  $Q^2 = 100 \text{ GeV}^2$  with the sea  $x\Sigma = 2x(\bar{u} + \bar{c} + \bar{d} + \bar{s} + \bar{b})$ .




FIG. 5: Same as Fig. 3 at the scale  $Q^2 = M_Z^2$ .




FIG. 6: Same as Fig. 4 at the scale  $Q^2 = M_Z^2$ .

### 3. THEORY FOR PDF FITS

In the following we describe the basic theoretical issues for a consistent determination of the twist-two PDFs from DIS and other hard scattering data, on the basis of perturbative QCD at NNLO using the  $\overline{\text{MS}}$  scheme for renormalization and factorization.

#### 3.1. Theory for analyses of DIS data

The world DIS data are provided in terms of reduced cross sections by the different experiments. QED and electroweak radiative corrections [29, 30] are applied, which requires careful study of different kinematic variables [30–33]. In this way also the contributions from the exchange of more than one gauge boson to the partonic twist-2 terms are taken care of. In part, also the very small QED corrections to the hadronic tensor are already accounted for. These have a flat kinematic behavior and amount to  $\mathcal{O}(1\%)$  or less [34–37].

The reduced cross sections are differential in either two of the kinematic variables in the set  $\{x, y, Q^2\}$ . The virtuality  $Q^2 = -q^2$  of the process is given by the 4-momentum transfer  $q$  to the hadronic system. The Bjorken variable is defined as  $x = Q^2/(sy)$ , with  $y = 2p \cdot q/s$ , and  $s = (p+l)^2$  the squared center-of-mass energy, where  $p$  and  $l$  denote the 4-momenta of the nucleon and the lepton. At energies much greater than the nucleon mass  $M$ , in the nucleon rest frame  $y$  is the fractional energy of the lepton transferred to the nucleon. The double differential cross sections used in the QCD analyses are given by [30, 38, 39]

$$\frac{d^2\sigma_{\text{NC}}^{l\pm N}}{dxdy} = \frac{2\pi\alpha^2 s}{Q^4} \left\{ \left[ 2(1-y) - 2xy \frac{M^2}{s} \right] F_2^{NC}(x, Q^2) + Y_{-x} F_3^{NC}(x, Q^2) + y^2 \left( 1 - \frac{2m_l^2}{Q^2} \right) 2xF_1^{NC}(x, Q^2) \right\}, \quad (2)$$

$$\frac{d^2\sigma_{\text{NC}}^{v(\bar{v})N}}{dxdy} = \frac{G_F^2 s}{16\pi} \left[ \frac{M_Z^2}{Q^2 + M_Z^2} \right]^2 \left\{ Y_+ W_2^{\text{NC}}(x, Q^2) \pm Y_{-x} W_3^{\text{NC}}(x, Q^2) - y^2 W_L^{\text{NC}}(x, Q^2) \right\}, \quad (3)$$

$$\frac{d^2\sigma_{\text{CC}}}{dxdy} = \frac{G_F^2 s}{4\pi} \left[ \frac{M_W^2}{Q^2 + M_W^2} \right]^2 \left\{ Y_+ W_2^{\text{CC}}(x, Q^2) \pm Y_{-x} W_3^{\text{CC}}(x, Q^2) - y^2 W_L^{\text{CC}}(x, Q^2) \right\}, \quad (4)$$

where  $\alpha$  and  $G_F$  denote the fine-structure and Fermi constants,  $Y_{\pm} = 1 \pm (1-y)^2$  and we keep the dependence on the masses of the nucleon ( $M$ ), the  $W$  and  $Z$  boson ( $M_W$ ,  $M_Z$ ) and the lepton ( $m_l$ ).

The structure functions  $F_i^{NC}$  and  $W_i$  are nonperturbative quantities defining the hadronic tensor. They can be measured by varying  $y$  at fixed  $Q^2$  and  $x$  and form the input to the subsequent analysis. Note that in some previous experiments, assumptions were made about the longitudinal structure functions  $F_L^{NC}$  and  $W_L$ , where (in the massless limit)

$$F_L^{NC}(x, Q^2) = F_2^{NC}(x, Q^2) - 2xF_1^{NC}(x, Q^2), \quad (5)$$

since at the time of the data analysis the corresponding QCD corrections were still missing. Therefore, it is important to use the differential cross sections in Eqs. (2)–(4) and to add the correct longitudinal structure functions [40, 41], cf. also [42, 43]. The structure functions are measured for DIS off massive proton and deuteron targets and are, therefore, subject to target mass corrections,

which play an important role in the region of lower values of  $Q^2$  and larger values of  $x$ . They are available in Refs. [38, 44, 45].

The neutral- and charged-current structure functions  $F_i^{NC}$ ,  $W_i^{NC}$  and  $W_i^{CC}$  consist of a sum of several terms, each weighted by powers of the QED and electroweak couplings, and  $F_i^{NC}$  also include the  $\gamma - Z$  mixing, which has to be accounted for, cf. [30, 38, 39]. Then, considering one specific gauge boson exchange, one arrives at a representation for the individual structure functions  $F_i$ , which are only subject to QCD corrections. For example, for pure photon exchange, they are given by

$$F_i(x, Q^2) = F_i^{\tau=2}(x, Q^2) + \sum_{k=2}^{\infty} \frac{C_i^{\tau=2k}(x, Q^2)}{Q^{2(k-1)}}, \quad (6)$$

where  $F_i^{\tau=2}$  denotes the leading-twist term and the coefficients  $C_i^{\tau}$  parametrize the higher twist contributions. The latter terms are of relevance for all DIS data sets, see Sec. 2.

Present day QCD analyses are aimed at determining the leading-twist contributions to the structure functions. There are two ways to account for the higher twist terms:

- (i) One is fitting the higher twist terms in  $F_i$ . A rigorous approach requires the knowledge of their scaling violations (term by term) and of the various Wilson coefficients to higher orders in  $\alpha_s$ , see e.g. Sec. 16 in Ref. [38]. Since at present this is practically out of reach, such fits remains rather phenomenological. Moreover, the size of the (non-singlet) higher twist contributions to the structure function  $F_2$  vary strongly with the correction applied to the leading-twist term up to next-to-next-to-next-to-leading order (N<sup>3</sup>LO), as shown in Ref. [42, 46]. Also, the non-singlet and singlet higher twist contributions are different [47, 48].
- (ii) One has to find appropriate cuts to sufficiently reduce the higher twist terms. For instance, in the flavor non-singlet analysis of Ref. [42] the cuts are taken to be  $Q^2 \geq 4 \text{ GeV}^2$ ,  $W^2 = M^2 + Q^2(1-x)/x \geq 12.5 \text{ GeV}^2$ . In the combined singlet and non-singlet analysis of Ref. [48],  $Q^2 \geq 10 \text{ GeV}^2$ ,  $W^2 \geq 12.5 \text{ GeV}^2$  have been used. These bounds are found empirically by cutting on  $W^2$  and/or  $Q^2$  starting from larger values. Applying these cuts severely limits the amount of large- $x$  DIS data to be fitted, and usually leads to an increase of the errors of  $\alpha_s(M_Z)$  and other fitted fundamental parameters and distributions.

Both methods (i) and (ii) allow to access the leading-twist contributions to the DIS structure functions, with some qualifications, however.

The cuts suggested in (ii) remove the large- $x$  region potentially sensitive to the higher twist terms. However, they do not affect the data at  $x \lesssim 0.1$ , where higher twist terms still play an important role [48, 49]. To some extent, the influence of higher twist can be dampened by using the DIS data for the structure function  $F_2$  instead of the cross section, since in this case the contribution to the structure function  $F_L$  need not be considered. It should be kept in mind, though, that the experimental separation of the structure functions  $F_2$  and  $F_L$  in the full phase space of common DIS experiments is very difficult without dedicated longitudinal–transverse cross section separations. Therefore, the data on  $F_2$  and  $F_3$  are typically extracted from the cross section once a certain model for the structure function  $F_L$  is taken. This approach is justified only at large  $x$ , however, where the contribution of  $F_L$  is small and even large uncertainties in the modeling of  $F_L$  cannot affect the extracted values of  $F_2$  and  $F_3$ . The procedure is not applicable for HERA kinematics, on the other hand, and introduces a bias into the analysis of the data taken by the New

Muon Collaboration (NMC), in particular, a shift in the value of  $\alpha_s$  preferred by the fit [43, 50], cf. Sec. 4. Nonetheless, the MMHT14 analysis [6] is still based on the DIS structure function data, as are the CJ15 and CT14 analyses [1, 3]. The latter two use cross section data for HERA, and for HERA and NMC, respectively, and structure function data elsewhere.

The deep-inelastic structure functions are inclusive quantities and contain massless parton and heavy-quark contributions,

$$F_i^{\tau=2}(x, Q^2) = F_i^{\text{massless}}(x, Q^2) + F_i^{\text{massive}}(x, Q^2). \quad (7)$$

Here the massless terms are given by

$$F_i^{\text{massless}}(x, Q^2) = \sum_j C_{i,j} \left( x, \frac{Q^2}{\mu^2} \right) \otimes f_j(x, \mu^2), \quad (8)$$

where  $C_{i,j}$  denote the massless Wilson coefficients,  $f_j$  the massless PDFs and  $\mu^2$  is the factorization scale. The Mellin convolution is abbreviated by  $\otimes$  and the sum over  $j$  is over all contributing partons. The renormalization group equation for  $F_i^{\text{massless}}$  allows one to eliminate the dependence on  $\mu^2$  order-by-order in perturbation theory. This also applies to  $F_i^{\tau=2}$ . Through the massive contributions  $F_i^{\text{massive}}$  there is a dependence on the heavy-quark masses  $m_c$  and  $m_b$  in the present world DIS data. Note that  $F_i^{\text{massive}}$  is *not* the structure function of a tagged heavy-flavor sample, which would be infrared sensitive [51]. Rather,  $F_i^{\text{massive}}$  is just given as the difference of the complete structure function  $F_i^{\tau=2}$  and the massless one in Eq. (8).

### 3.1.1. Massless PDFs

For all QCD calculations we use perturbation theory. The factorized representation in terms of Wilson coefficients and PDFs is obtained using the light-cone expansion [52–55]. For a proper definition of the Wilson coefficients and the PDFs one has to use the LSZ formalism and refer to asymptotic states at large times  $t \rightarrow \pm\infty$ , given by massless partons. We first describe the massless contributions in Eqs. (7) and (8), and then discuss the contribution of heavy quarks. The Wilson coefficients in Eq. (8) have a perturbative expansion in the strong coupling constant. At one- [56], two- [57–64], and three-loop order [40, 41, 65–67] they have been calculated for the neutral-current structure functions  $F_i$ , with  $i = 1, 2, 3$ , except for the  $\gamma - Z$  mixing contribution at three loops.

The structure functions in general depend on the following three non-singlet and singlet combinations of parton densities:

$$q_{jk}^{\pm} = f_j \pm \bar{f}_j - (f_k \pm \bar{f}_k), \quad q^v = \sum_{l=1}^{n_f} (f_l - \bar{f}_l), \quad q^s = \sum_{l=1}^{n_f} (f_l + \bar{f}_l), \quad (9)$$

with the light-quark distributions  $f_i$  of flavor  $i$  and  $n_f$  the number of massless flavors. These combinations evolve in  $\mu^2$  from an initial scale  $\mu_0^2$  by the QCD evolution equations, where the singlet distribution  $q^s(x, \mu^2)$  mixes with the gluon distribution  $g(x, \mu^2)$ ,

$$\frac{d}{d \ln(\mu^2)} q^i(x, \mu^2) = P^i(x) \otimes q^i(x, \mu^2), \quad i = \pm, v, \quad (10)$$



$$\frac{d}{d\ln(\mu^2)} \begin{pmatrix} q^s(x, \mu^2) \\ g(x, \mu^2) \end{pmatrix} = \begin{pmatrix} P_{qq}(x) & P_{qg}(x) \\ P_{gq}(x) & P_{gg}(x) \end{pmatrix} \otimes \begin{pmatrix} q^s(x, \mu^2) \\ g(x, \mu^2) \end{pmatrix}. \quad (11)$$

The non-singlet splitting functions are given by

$$P^\pm(x) = P_{qq}(x) \pm P_{q\bar{q}}(x), \quad (12)$$

$$P^v(x) = P_{qq}(x) - P_{q\bar{q}}(x) + n_f (P_{qq}^s(x) - P_{q\bar{q}}^s(x)), \quad (13)$$

while the anomalous dimensions  $\gamma_{ij}$  corresponding to the splitting functions  $P_{ij}$  are obtained by a Mellin transform,

$$\gamma_{ij}(N) = - \int_0^1 dx x^N P_{ij}(x). \quad (14)$$

where we suppress for brevity the dependence of  $P_{ij}$  and  $\gamma_{ij}$  on the strong coupling  $a_s(\mu^2) = \alpha_s(\mu^2)/(4\pi)$ . The  $P_{ij}$  are known as well at one- [68–73], two- [64, 74–85] and at three-loop order [86, 87] (see also [88, 89] for checks of  $P_{ps}$  and  $P_{gg}$  at that order). The scale evolution of the strong coupling constant in the  $\overline{\text{MS}}$  scheme is given by

$$\frac{da_s(\mu^2)}{d\ln(\mu^2)} = - \sum_{k=0}^{\infty} \beta_k a_s^{k+2}(\mu^2), \quad (15)$$

where  $\beta_k$  denote the expansion coefficients of the QCD  $\beta$ -function [90–99].

The evolution equations (10), (11) can be either solved in  $x$ - or Mellin (or moment)  $N$ -space. In Mellin-space, defined by the transform Eq. (14), an analytic solution is possible [100–103] by arranging the solution systematically in powers of the coupling constants  $a_s(\mu^2)$  and  $a_s(\mu_0^2)$ , and even forming factorization-scheme invariant expressions. In case of the  $x$ -space solutions this is usually not done due to the necessary iterative solution. In the small- $x$  region the iterative solution usually leads to a pile-up of a few per cent [104]. This can be corrected for in  $x$ -space solutions by applying the method given in [105]. Likewise, the iterated solution can be obtained in Mellin  $N$ -space and is a standard option of the evolution program QCD-Pegasus [106].

### 3.1.2. Heavy-quark structure functions

Disregarding contributions from charm at the input scale (“intrinsic charm”), cf. [107–109], the heavy-flavor corrections to the DIS functions are described by Wilson coefficients. The leading order results are of  $\mathcal{O}(a_s)$ . Higher order corrections in the perturbative expansions are, therefore, of  $\mathcal{O}(a_s^2)$  at NLO, and of  $\mathcal{O}(a_s^3)$  at NNLO, similar to the case of the longitudinal structure function [40, 41]. The corrections in the neutral- and charged-current cases are available in one- [110–117] and two-loop order [118–121], where the latter corrections were given in semi-analytic form.

For the neutral-current exchange the heavy-flavor contributions to the structure functions  $F_i$  with  $i = 2, L$  are [122, 123]:

$$F_i^{\text{massive}}(x, n_f + 2, Q^2, m_c^2, m_b^2) = \sum_{i=c,b} x \left\{ \sum_{k=1}^{n_f} e_k^2 \left\{ L_{i,q}^{ns} \left( x, n_f + 1, \frac{Q^2}{\mu^2}, \frac{m_i^2}{\mu^2} \right) \otimes [f_k(x, \mu^2, n_f) + \bar{f}_k(x, \mu^2, n_f)] \right. \right.$$

$$\begin{aligned}
& + \frac{1}{n_f} L_{i,q}^{ps} \left( x, n_f + 1, \frac{Q^2}{\mu^2}, \frac{m_i^2}{\mu^2} \right) \otimes q^s(x, \mu^2, n_f) \\
& + \frac{1}{n_f} L_{i,g}^s \left( x, n_f + 1, \frac{Q^2}{\mu^2}, \frac{m_i^2}{\mu^2} \right) \otimes g(x, \mu^2, n_f) \Big\} \\
& + e_i^2 \left[ H_{i,q}^{ps} \left( x, n_f + 1, \frac{Q^2}{\mu^2}, \frac{m_i^2}{\mu^2} \right) \otimes q^s(x, \mu^2, n_f) \right. \\
& \quad \left. + H_{i,g}^s \left( x, n_f + 1, \frac{Q^2}{\mu^2}, \frac{m_i^2}{\mu^2} \right) \otimes g(x, \mu^2, n_f) \right] \Big\} \\
& + \delta_{i,2} F_2^{\text{massive},\{c,b\}}(x, n_f + 2, Q^2, m_c^2, m_b^2). \tag{16}
\end{aligned}$$

They are determined by five massive Wilson coefficients,  $L_{i,k}^{\{ns,ps,s\}}$  and  $H_{i,k}^{\{ps,s\}}$ , where the electroweak current couples either to a massless ( $L_{i,k}$ ) or the massive ( $H_{i,k}$ ) quark line. From three-loop order onwards there are contributions containing both heavy flavors  $c$  and  $b$  in a non-separable form, denoted by  $F_2^{\text{massive},\{c,b\}}$ , in Eq. (16). The PDFs and the coupling constant in Eq. (16) are defined in the  $\overline{\text{MS}}$  scheme, while the heavy quark masses are taken either in the on-shell or  $\overline{\text{MS}}$  schemes [123, 124]. The relations of the heavy quark masses between the pole mass (on-shell scheme) and the  $\overline{\text{MS}}$  scheme are available to four-loop order [125]. Due to its better perturbative stability, the  $\overline{\text{MS}}$  scheme for the definition of the heavy-quark mass is preferred.

For  $Q^2 \gg m_i^2$  the asymptotic corrections to  $F_L$  are available at three-loop order [122, 126]. For  $F_2$ , four out of the five massive Wilson coefficients,  $L_{2,q}^{ns}$ ,  $L_{2,q}^{ps}$ ,  $L_{2,g}^s$  and  $H_{2,q}^{ps}$  are known as well [88, 122, 127–129] at large scales  $Q^2$ . For the remaining coefficient,  $H_{2,g}^s$ , an estimate has been made in Ref. [130] based on the anticipated small- $x$  behavior [131], a series of moments calculated in [123], and two-loop operator matrix elements from Refs. [132, 133]. This provides a good approximation of the NNLO corrections.

### 3.2. Heavy-flavor PDFs

An important issue in PDF fits concerns the number of active quark flavors and the theoretical description of heavy quarks such as charm and bottom. Due to the large range of hard scales  $Q$  for the scattering processes considered, different effective theories may be applied. At low scales, when  $Q \simeq \mathcal{O}(\text{few})$  GeV, one typically works with  $n_f = 3$  massless quark flavors, setting  $n_f = 3$  in the hard scattering cross section, the evolution kernels and the anomalous dimensions. In this case, only the light-quark PDFs for up, down and strange are taken into account. At higher scales, e.g., for hadro-production of jets at high transverse momentum  $p_t$  or top quarks, additional dynamical degrees of freedom lead to theories with  $n_f > 3$ . By means of the renormalization group and matching these are related to the case with  $n_f = 3$  massless quarks. Technically, one has to apply decoupling relations [134] at some matching scale  $\mu$ , for instance in the transition of  $\alpha_s^{(n_f)} \rightarrow \alpha_s^{(n_f+1)}$ . This introduces some logarithmic dependence on the masses of the heavy quarks  $m_c$ ,  $m_b$  and  $m_t$  for charm, bottom and top. One should also note that the matching of the effective theories for  $n_f \rightarrow n_f + 1$  does not need to be smooth. In fact, it introduces discontinuities, such as for the running coupling as a solution of the QCD  $\beta$ -function at higher order in the perturbative expansion, where  $\alpha_s^{(n_f)}(\mu) \neq \alpha_s^{(n_f+1)}(\mu)$  in the  $\overline{\text{MS}}$  scheme at the matching scale  $\mu$ , see e.g., [135].

In a similar manner, PDFs in theories with a fixed number  $n_f > 3$  of quark flavors are related

to those for  $n_f = 3$  with the help of heavy-quark operator matrix elements (OMEs)  $A_{ij}$  at a chosen matching scale  $\mu$ . Starting with the PDFs in a so-called fixed-flavor number scheme (FFNS) with  $n_f$  fixed, one has  $f_i^{(n_f)} \rightarrow f_i^{(n_f+1)}$  for the light-quark distributions  $f_i$  and  $(q^{s,(n_f)}, g^{(n_f)}) \rightarrow (q^{s,(n_f+1)}, g^{(n_f+1)})$  for the gluon and the singlet quark distributions with operator mixing in the singlet sector. In particular, one has [123, 136]

$$\begin{aligned} f_k(n_f+1, \mu^2) + f_{\bar{k}}(n_f+1, \mu^2) &= A_{qq,h}^{ns}\left(n_f, \frac{\mu^2}{m^2}\right) \otimes [f_k(n_f, \mu^2) + f_{\bar{k}}(n_f, \mu^2)] \\ &\quad + \frac{1}{n_f} A_{qq,h}^{ps}\left(n_f, \frac{\mu^2}{m^2}\right) \otimes q^s(n_f, \mu^2) \\ &\quad + \frac{1}{n_f} A_{qg,h}^s\left(n_f, \frac{\mu^2}{m^2}\right) \otimes g(n_f, \mu^2), \end{aligned} \quad (17)$$

$$g(n_f+1, \mu^2) = A_{gq,h}^s\left(n_f, \frac{\mu^2}{m^2}\right) \otimes q^s(n_f, \mu^2) + A_{gg,h}^s\left(n_f, \frac{\mu^2}{m^2}\right) \otimes g(n_f, \mu^2), \quad (18)$$

$$\begin{aligned} q^s(n_f+1, \mu^2) &= \left[ A_{qq,h}^{ns}\left(n_f, \frac{\mu^2}{m^2}\right) + A_{qq,h}^{ps}\left(n_f, \frac{\mu^2}{m^2}\right) + A_{hq}^{ps}\left(n_f, \frac{\mu^2}{m^2}\right) \right] \otimes q^s(n_f, \mu^2) \\ &\quad + \left[ A_{qg,h}^s\left(n_f, \frac{\mu^2}{m^2}\right) + A_{hg}^s\left(n_f, \frac{\mu^2}{m^2}\right) \right] \otimes g(n_f, \mu^2). \end{aligned} \quad (19)$$

PDFs for charm and bottom ( $h = c, b$ ) are then constructed as

$$f_{h+\bar{h}}(n_f+1, \mu^2) = A_{hq}^{ps}\left(n_f, \frac{\mu^2}{m^2}\right) \otimes q^s(n_f, \mu^2) + A_{hg}^s\left(n_f, \frac{\mu^2}{m^2}\right) \otimes g(n_f, \mu^2) \quad (20)$$

at the matching scale  $\mu$  from the quark singlet and gluon PDFs with  $h = \bar{h}$ .

The matching conditions are typically imposed at the scale  $\mu = m_h$ , and  $f_{h+\bar{h}} = 0$  is assumed for scales  $\mu \leq m_h$ . The necessary heavy-quark OMEs  $A_{ij}$  depend logarithmically on the heavy-quark masses as  $\alpha_s^l \ln^k(\mu^2/m_h^2)$  with  $0 \leq k \leq l$  in the perturbative expansion. As discussed above, the OMEs are known to NLO analytically [132, 137] and at NNLO either exactly or to a good approximation [88, 123, 127, 130, 138]. Thus, charm and bottom PDFs can be consistently extracted in QCD with a fixed number  $n_f = 3, 4$  or  $5$ .

It should be stressed, however, that the decoupling relations for PDFs in Eqs. (17)–(20) assume the presence of one heavy quark at a time, with the bottom-quark contributions ignored when considering charm-quark PDFs, for instance. Beginning at three-loop order, however, there are graphs containing both charm- and bottom-quark lines, and charm quarks cannot be treated as massless at the scale of the bottom-quark due to  $(m_c/m_b)^2 \approx 1/10$ . Such terms cannot be attributed to either the charm- or bottom-quark PDFs, but rather one has to decouple charm and bottom quarks together at some large scale. The simultaneous decoupling of bottom and charm quarks in the strong coupling constant  $\alpha_s$  is discussed, for instance, in Ref. [139].

### 3.3. Heavy-quarks schemes

#### 3.3.1. Variable-flavor number schemes

The hard scattering cross sections also depend on the number of flavors  $n_f$  and additional parton channels may open up, which have to be included as well. In addition, processes involving

massive quarks depend logarithmically on the ratio  $Q^2/m_h^2$ , where  $Q$  is some hard scale associated with the scattering. For the heavy-flavor Wilson coefficients in Eq. (16) these logarithms are of the type  $\alpha_s^l \ln^k(Q^2/m_h^2)$  with  $1 \leq k \leq l$  in perturbation theory. These originate from collinear singularities screened by the heavy-quark mass due to the constrained phase space for gluon emission from massive quark lines, and as a prefactor of these logarithms one has the standard splitting functions. In addition to logarithmic terms, there are also power corrections  $(m_h^2/Q^2)^l$  in the heavy-flavor Wilson coefficients, usually appearing in form of higher transcendental functions. In the asymptotic regime of  $Q^2 \gg m_h^2$  the logarithms dominate and the kinematic dependence is measured experimentally, for instance in the tagged flavor case for charm-quark pairs in the structure function  $F_2^{c\bar{c}}$ . Logarithms of a similar kind are also experimentally observed in differential distributions, e.g. due to the QED corrections proportional to  $\ln^k(Q^2/m_l^2)$  with  $m_l$  being the charged lepton mass, cf. [29, 30].

The resummation of the logarithms  $\alpha_s^l \ln^k(Q^2/m_h^2)$  to all orders in perturbation theory is effectively carried out by the transition  $n_f \rightarrow n_f + 1$  along with the introduction of new heavy-quark PDFs as described in Eqs. (17)–(20). Whether such a transition is appropriate or not depends, of course, on the detailed kinematics. If the hard scale is closer to threshold,  $Q^2 \simeq m_h^2$ , a description with  $n_f$  light flavors is more suitable, while for  $Q^2 \gg m_h^2$  one switches to a theory with  $n_f + 1$  massless flavors. In order to achieve a unified description for hard scattering cross sections both at low scales  $Q^2 \simeq m_h^2$  and asymptotically for  $Q^2 \gg m_h^2$ , so-called variable-flavor number schemes (VFNS) have been constructed. Effectively, these aim at an interpolation between the asymptotic limits of the quarks being very light or very heavy relative to the other hard scales of the process. At the LHC such considerations apply to processes with bottom quarks in the initial state such as single top-quark production as well as bottom-quark initiated Higgs boson production (see Ref. [140] for more recent studies and Ref. [141] for the so-called *Santander matching* scheme for Higgs boson production in  $b\bar{b}$  annihilation).

Of particular interest for PDF fits is the reduced cross section to the pair-production of heavy quarks in DIS, which is parametrized in terms of the DIS heavy-quark structure functions  $F_i^h$  for  $i = 2, L$  in Eq. (16) and with heavy-flavor Wilson coefficients which are known exactly at NLO [118], and to a good approximation at NNLO [130] in QCD. For the interpolation  $n_f \rightarrow n_f + 1$  of the heavy-quark structure functions  $F_i^h$  a number of so-called general-mass VFNS (GM-VFNS) have been discussed in the literature, such as ACOT [142–144], BMSN [136], FONLL [145] or RT [146]. These keep  $m_h \neq 0$  and are to be distinguished from the zero-mass VFNS (ZM-VFNS), which describes essentially the massless case. Note that presently the GM-VFNS are applied only to one single heavy flavor at the time. That is the sequential transition  $n_f \rightarrow n_f + 1$ , so that the charm or bottom quarks are not considered simultaneously and charm-quark mass effects in the bottom-quark structure function  $F_i^b$  are neglected as discussed above.

The various GM-VFNS contain a number of additional assumptions, and some come in more than one variety. The GM-VFNS differ, for instance, in the way the low- $Q^2$  region is modeled. This modeling is a necessary undertaking to provide a reasonable behavior of the VFNS in the kinematical regime of present DIS data. Additional assumptions in the GM-VFNS are related to the matching scale  $\mu$  for the transition  $n_f \rightarrow n_f + 1$  as the adopted choice  $\mu = m_h$  is not unique, see [147] for an in-depth discussion.

Briefly, the problem can be illustrated with the heavy-quark velocity, the leading order formula [114] being

$$v = \sqrt{1 - \frac{4m_h^2}{s}} = \sqrt{1 - \frac{4m_h^2 x}{Q^2(1-x)}}, \quad x \leq \frac{1}{1 + 4m_h^2/Q^2}. \quad (21)$$




FIG. 7: Comparison of HERA data for the DIS pair-production of charm quarks [148] to the QCD predictions at NNLO in the FFNS using ABM12 [2] and JR14 [5] PDFs with a running charm-quark mass.

The transition  $n_f \rightarrow n_f + 1$  when the corresponding flavor is considered as nearly massless requires light-like velocities  $v \simeq 1$ . That implies the absence of all power corrections  $(m_h^2/Q^2)^l$  in the heavy-flavor Wilson coefficients at the matching scale  $\mu^2$ . In practice, the matching is often applied at the scale  $\mu^2 = m_h^2$  and for kinematics  $Q^2 \gg m_h^2$ , where this condition is not fulfilled, which implies restrictions on the range in  $x$  in Eq. (21).

Finally, the logarithmic accuracy of the resummation for large scales  $Q^2 \gg m_h^2$ , or the order of perturbation theory in current implementations of GM-VFNS, is often not consistent. For example, NNLO evolution [86, 87] of the massless PDFs is sometimes combined with the heavy-quark OMEs at NLO [132, 137], omitting NNLO results [88, 123, 127, 130].

Altogether, these facts introduce a significant model dependence in any GM-VFNS implementation. A sensitive parameter to test this model dependence is the extraction of the charm- or bottom-quark mass used in different versions of GM-VFNS and subsequent comparison with the Particle Data Group (PDG) results [39]. In addition, the quality of the various GM-VFNS can be quantified with the goodness-of-fit for the description of HERA data on DIS charm-quark production obtained from the combination of individual H1 and ZEUS results [148].

PDF sets	$m_c$ [GeV]	$m_c$ renorm. scheme	theory method ( $F_2^c$ scheme)	theory accuracy for heavy quark DIS Wilson coeff.	$\chi^2$ /NDP for HERA data [148] with xFitter [149, 150]	
					(with unc.)	(nominal)
ABM12 [2] <sup>a</sup>	$1.24^{+0.05}_{-0.03}$	$\overline{\text{MS}}$ $m_c(m_c)$	FFNS ( $n_f = 3$ )	NNLO <sub>approx</sub>	65/52	66/52
CJ15 [1]	1.3	$m_c^{\text{pole}}$	SACOT [143]	NLO	117/52	117/52
CT14 [3] <sup>b</sup>	(NLO)	$m_c^{\text{pole}}$	SACOT( $\chi$ ) [144]	NLO	51/47	70/47
	(NNLO)	$m_c^{\text{pole}}$	SACOT( $\chi$ ) [144]	NLO	64/47	130/47
HERAPDF2.0 [4]	(NLO)	$m_c^{\text{pole}}$	RT optimal [146]	NLO	67/52	67/52
	(NNLO)	$m_c^{\text{pole}}$	RT optimal [146]	NLO	62/52	62/52
JR14 [5] <sup>c</sup>	1.3	$\overline{\text{MS}}$ $m_c(m_c)$	FFNS ( $n_f = 3$ )	NNLO <sub>approx</sub>	62/52	62/52
MMHT14 [6]	(NLO)	$m_c^{\text{pole}}$	RT optimal [146]	NLO	72/52	78/52
	(NNLO)	$m_c^{\text{pole}}$	RT optimal [146]	NLO	71/52	83/52
NNPDF3.0 [7]	(NLO)	$m_c^{\text{pole}}$	FONLL-B [145]	NLO	58/52	60/52
	(NNLO)	$m_c^{\text{pole}}$	FONLL-C [145]	NLO	67/52	69/52
PDF4LHC15 [8] <sup>d</sup>	–	–	FONLL-B [145]	–	58/52	64/52
	–	–	RT optimal [146]	–	71/52	75/52
	–	–	SACOT( $\chi$ ) [144]	–	51/47	76/47

<sup>a</sup>The value of  $m_c$  in ABM12 is determined from a fit to HERA data [148].

<sup>b</sup> The data comparison always applies the SACOT( $\chi$ ) scheme at NLO as implemented in xFitter [149, 150]. The implementation of this scheme differs from the one used by CT14. Removing the  $Q^2$ -cut on the HERA data [148] one obtains  $\chi^2$ /NDP = 158/52 (582/52) with PDF uncertainties and 258/52 (648/52) for the central fit at NLO (NNLO).

<sup>c</sup> The  $\chi^2$ /NDP values are determined for the dynamical set JR14NNLO08VF.

<sup>d</sup> The data comparison uses the xFitter [149, 150] implementation of the schemes FONLL-B, RT optimal and SACOT( $\chi$ ) with the set PDF4LHC\_100 at NLO.

TABLE 3: Values of the charm quark mass and renormalization scheme used in the PDF fits together with a summary of schemes chosen for the description of the charm-quark structure function  $F_2^c$  and the theoretical accuracy for the massive quark DIS Wilson coefficients. The values of  $\chi^2$ /NDP for the DIS charm production cross section data [148] are given in the left column (with unc.) with the account of PDF uncertainties (CT14 PDF errors scaled from 90% c.l. to 68% c.l., i.e., reduced by a factor 1.645) and in the right column (nominal) for the central prediction of each PDF set. The values for CT14 and for PDF4LHC with the SACOT( $\chi$ ) scheme have been determined with a cut on  $Q^2 \geq 5 \text{ GeV}^2$  on the HERA data [148].




FIG. 8: Same as Fig. 7 with QCD predictions at NLO and NNLO in the RT optimal [146] VFNS using the HERAPDF2.0 [4] PDF sets at NLO and NNLO.

### 3.3.2. Validation with DIS charm-quark production

The H1 and ZEUS combined data are unique for tests of GM-VFNS and span the region of  $2.5 \leq Q^2 \leq 2000 \text{ GeV}^2$  and  $3 \times 10^{-5} \leq x \leq 0.05$ . Values for the charm-quark mass and  $\chi^2/\text{NDP}$  for the individual PDF sets ABM12, CJ15, CT14, HERAPDF2.0, JR14, MMHT14, NNPDF3.0 as well as the averaged set PDF4LHC15 are given in Tab. 3, along with the information on the scheme choice for the heavy-quark structure functions and on the theoretical accuracy for the massive quark DIS Wilson coefficients. Comparisons to data are shown in Figs. 7–10. Note that Tab. 3 adopts the standard definition of perturbative orders for the heavy-quark structure functions. This is not shared by CT14, MMHT14 and NNPDF3.0 in their GM-VFNS. There the Born contribution to the heavy-quark Wilson coefficients for  $ep \rightarrow c\bar{c}$ , which is proportional to  $\mathcal{O}(\alpha_s)$ , is referred to as being “NLO”. Analogously, the one-loop corrections of order  $\mathcal{O}(\alpha_s^2)$  are denoted by “NNLO”.

Table 3 and Fig. 7 show that the ABM12 [2] and JR14 [5] PDFs at NNLO, using charm-quark masses in the  $\overline{\text{MS}}$  scheme, provide a good description of the data. Both ABM12 and JR14 use the approximate massive three-loop Wilson coefficients as obtained in [130] by interpolating between existing soft-gluon threshold resummation results and asymptotic ( $Q^2 \gg m_c^2$ ) coefficients [123, 127]. This is referred to as  $\text{NNLO}_{\text{approx}}$  in Tab. 3. The HERAPDF2.0 fit [4] also obtains a good description of the data, cf. Fig. 8. On the other hand, the SACOT [143] GM-VFNS at NLO used by CJ15 [1] does not describe the data too well, although we should note that the HERA charm data were not included in the CJ15 fit itself.




FIG. 9: Same as Fig. 7 with QCD predictions at NLO and different versions of VFNS using the PDFs CT14 [3] (SACOT- $\chi$  [144]), MMHT14 [6] (RT optimal [146]), and NNPDF3.0 [7] (FONLL-B [145]).

The remarkable fact in Tab. 3 and Fig. 9 is, however, that the GM-VFNS SACOT( $\chi$ ) [144] of CT14 [3] and RT optimal [146] of MMHT14 [6] have difficulties in describing the data. Note that CT14 has applied a cut  $Q^2 \geq 5 \text{ GeV}^2$  on the HERA data [148], excluding the bin with data at  $Q^2 = 2.5 \text{ GeV}^2$  (cf. the upper left plot in Fig. 9) from the fit. We have checked that including the low  $Q^2$  bin leads to a dramatic deterioration of the fit quality.

In addition, the schemes SACOT( $\chi$ ) and RT optimal as well as FONLL-C [145] of NNPDF3.0 [7] do not improve the fit quality when comparing NLO and NNLO fits. We note in this context that those fits do not include essential NNLO results for the heavy-quark Wilson coefficients [88, 123, 127, 130] in their theory predictions. The averaged set PDF4LHC15 [8], shown in Fig. 10, mixes PDFs derived with different mass schemes (ACOT, FONLL and RT) and does not describe the data very well for virtualities up to  $Q^2 \lesssim 20 \text{ GeV}^2$ .

### 3.3.3. Charm-quark mass

Dedicated studies of the charm-quark mass dependence have been performed by several groups. In the  $\overline{\text{MS}}$  scheme, the value of  $m_c(m_c) = 1.24^{+0.04}_{-0.08} \text{ GeV}$  has been obtained in [151] together with  $\chi^2/\text{NPD}=61/52$  for the description of the HERA data [148] as a variant of the ABM11 fit [48]. Other groups, which keep a fixed value of  $m_c$  in the analyses, cf. Tab. 3, have studied the effects of varying  $m_c$  in predefined ranges. This has been done, for example, in the older NNPDF2.1 [152]






FIG. 10: Same as Fig. 7 with QCD predictions at NLO using the PDF4LHC\_100 PDF set and various VFNS: FONLL-B [145], RT optimal [146] and SACOT- $\chi$  [144].

and MSTW analyses [153] as well as for the MMHT PDFs [9]. The latter yields a pole mass of  $m_c^{\text{pole}} = 1.25$  GeV as the best fit with  $\chi^2/\text{NPD} = 75/52$ , while the nominal fit uses  $m_c^{\text{pole}} = 1.4$  GeV at the price of a deterioration in the value of  $\chi^2/\text{NPD} = 82/52$ . HERAPDF2.0 [4] has performed a scan of the values of  $\chi^2/\text{NPD}$  leading to  $m_c^{\text{pole}} = 1.43$  GeV at NNLO quoted in Tab. 3 as the best fit. NNPDF3.0 computes heavy quark structure functions with expressions for the pole mass definition, but adopts numerical values for the charm quark pole mass,  $m_c^{\text{pole}} = 1.275$  GeV, which corresponds to the current PDG value for the  $\overline{\text{MS}}$  mass. This value is different from the one used in NNPDF2.3, namely  $m_c^{\text{pole}} = \sqrt{2}$  GeV.

In this context, it is worth to point out that the running mass  $m_c(\mu)$  in the  $\overline{\text{MS}}$  scheme is free from renormalon ambiguities and can therefore be determined with high precision. The PDG [39] quotes  $m_c(m_c) = 1.275 \pm 0.025$  GeV based on the averaging different mass determination in various kinematics. DIS charm-quark production analyzed in the FFNS ( $n_f = 3$ ) leads to  $m_c(m_c) = 1.24 \pm 0.03 \pm_{-0.03}^{+0.03}$  GeV at NNLO [151], while measurements of the  $\overline{\text{MS}}$  mass in  $e^+e^-$  annihilation give, for instance,  $m_c(m_c) = 1.279 \pm 0.013$  GeV [154] and  $m_c(m_c) = 1.288 \pm 0.020$  GeV [155]. The determination from quarkonium 1S energy levels yields  $m_c(m_c) = 1.246 \pm 0.023$  GeV [156]. All these values are consistent with each other within the uncertainties.

In contrast, the accuracy of the pole mass  $m_c^{\text{pole}}$  is limited to be of the order of the QCD scale  $\Lambda_{\text{QCD}}$  and, moreover, the conversion from the  $\overline{\text{MS}}$  mass  $m_c(m_c)$  at low scales to the pole mass  $m_c^{\text{pole}}$  does not converge. Using  $\alpha_s(M_Z) = 0.1184$ , for example, the conversion yields for the central value




FIG. 11: The  $1\sigma$  band for the  $d/u$  ratio for the 4-flavor scheme and at the factorization scale  $\mu = 2$  GeV obtained in the PDF analyses including forward  $W^\pm$  data (CT14 [3]: red right-tilted hatch, ABMP15 [24]: gray shaded area, CJ15 [1]: black dotted lines) in comparison to those including the central  $W,Z$  data only and a cut of  $W^2 \gtrsim 13$  GeV<sup>2</sup> imposed on the deuteron DIS data (MMHT14 [6]: blue dashed lines, NNPDF3.0 [7]: green left-tilted hatch).

of the PDG  $m_c^{\text{pole}} = 1.47$  GeV at one loop,  $m_c^{\text{pole}} = 1.67$  GeV at two loops,  $m_c^{\text{pole}} = 1.93$  GeV at three loops, and  $m_c^{\text{pole}} = 2.39$  GeV at four-loops [125]. The PDG quotes  $m_c^{\text{pole}} = 1.67 \pm 0.07$  GeV for conversion at two loops.

The low values for the pole mass of the charm quark assumed or obtained in some PDF fits as shown in Tab. 3 are thus not compatible with other determinations and with the world average. The rigorous determination of the charm-quark mass discussed, for instance, in [151] provides a more controlled way of determining  $m_c$  from the world DIS data, taking also into account its correlation with  $\alpha_s(M_Z)$ .

### 3.4. Light-flavor PDFs

#### 3.4.1. Up- and down-quark distributions

The total quark contribution to nucleon matrix elements is known fairly well due to constraints from the available DIS data obtained in the fixed-target and collider experiments in the  $x$ -range  $10^{-4} \lesssim x \lesssim 0.8$ . However, a thorough disentangling of the quark flavor structure is still a challenging task in any PDF analysis. At moderate and large  $x$  values this has been routinely achieved by using a combination of the DIS data obtained on proton and deuteron targets. However, uncertainties in the modeling of nuclear corrections in the deuteron introduce a controllable source of theoretical uncertainty on the  $d$ -quark PDF obtained in this way, especially at large  $x$ , as discussed below.

An alternative way to resolve the  $u$ - and  $d$ -quark contributions is to use data on  $W$ - and  $Z$ -boson production obtained in  $pp$  and  $p\bar{p}$  collisions at the LHC and Tevatron, respectively. Those experiments probe the  $W$  and  $Z$  rapidity distributions up to rapidities of  $y = 3 - 4$ , depending




FIG. 12: Same as Fig. 11 for the SU(2) flavor asymmetry of the light-quark sea, or the “isospin” asymmetry,  $I(x) = [\bar{d}(x) - \bar{u}(x)]$ .

PDF sets	$\chi^2/\text{NDP}$ (ATLAS data [16])	theory accuracy	theory method
ABM12 [2]	34.5/30	NNLO	FEWZ3.1 [157], DYNNLO [158]
ABMP15 [24]	32.3/30	NNLO	FEWZ3.1 [157]
CT14 [3]	42/30	NNLL	ResBos [159]
MMHT14 [9]	39/30	(NNLO)	APPLGrid [160] (NNLO $K$ -factors constant)
NNPDF3.0 [7]	35.4/30	NNLO	APPLGrid [160], $C$ -factors [161] (kinematic dependence with FEWZ3.1 [157])

TABLE 4: Description of the ATLAS data at  $\sqrt{s} = 7$  TeV for  $W^\pm \rightarrow l^\pm \nu$ ,  $Z \rightarrow l^+ l^-$  (Ref. [16]) used in the PDF fits. The columns indicate the QCD accuracy of the theoretical predictions along with the tools used to obtain them.

on details of the experiments, with an integrated luminosity of  $O(1) \text{ fb}^{-1}$  achieved in each run. Such data samples are quite competitive in accuracy with the ones obtained in fixed-target DIS experiments, and provide simultaneously constraints on the quark and anti-quark PDFs at large and small  $x$ . Furthermore, the  $d$ -quark PDF extracted from a combination of the existing data on DIS off protons and  $W/Z$ -boson production in  $pp(p\bar{p})$  collisions are not sensitive to nuclear corrections. Moreover, if DIS data with large hadronic invariant masses  $W^2$  are not used in the analyses in order to reduce the sensitivity to higher twist contributions, the statistical potential of the deuteron data is reduced and they become less competitive as compared to the collider data, cf. Fig. 11.




FIG. 13: The  $1\sigma$  band for the strange sea suppression factor  $r_s = (s + \bar{s})/(2\bar{d})$  as a function of Bjorken  $x$  obtained in the variants of the ABM analysis [162] based on the combination of the data by NuTeV/CCFR [163], CHORUS [164] and NOMAD [165] (shaded area), and CHORUS [164], CMS [166] and ATLAS [167] (dashed lines), compared with the results obtained by the CMS analysis [17] (hatched area) and by the ATLAS  $epWZ$ -fit [167, 168] at different values of  $x$  (full circles). All quantities refer to the factorization scale  $\mu^2 = 1.9 \text{ GeV}^2$ .

As mentioned above, one can further constrain the  $u$ - and  $d$ -quark flavor separated distributions by utilizing fixed-target deuteron DIS data. However, nuclear effects need to be accounted for in cross sections and structure functions in order to access the underlying PDFs. The theoretical uncertainty inherent in this nuclear correction procedure should be added to the statistical PDF uncertainties. Nonetheless, the reduction of the uncertainties due to the increased number of fitted data points is even greater, leading to an overall smaller  $d$ -quark PDF uncertainty than in fits performed without deuterium data [24, 169, 170]. Furthermore, as shown in [1], and discussed in more detail below, it is possible to significantly reduce the nuclear correction uncertainty by exploiting the interplay of the deuteron DIS data and the recent high-statistics  $D\bar{O}$  data on the reconstructed  $W^\pm$  boson charge asymmetry at large rapidity, which is equally sensitive to the  $d/u$  ratio but is not affected by nuclear corrections.

The  $W/Z$ -boson collider data also provide a valuable constraint on the small- $x$  quark PDFs. In particular, the charge asymmetry of leptons originating from the  $W$  decays is sensitive to the  $SU(2)$  flavor asymmetry of the non-strange sea, also referred to as the “isospin” asymmetry  $I(x) = [\bar{d}(x) - \bar{u}(x)]$  at small  $x$ . This asymmetry is constrained by the DY data from fixed-targets with protons and deuterons collected by the Fermilab experiment E866 [171]. However, the E866 data are not sensitive to the value of  $I(x)$  at small  $x$  ( $x \lesssim 0.2$ ). Therefore,  $I(x)$  is sometimes parametrized in a Regge-like form as  $I(x) \sim x^{0.5}$  such that it vanishes at  $x = 0$  (cf. the MMHT results in Fig. 12).

The large-rapidity tail of the  $W/Z$ -boson production data allows for a model-independent check of  $I(x)$  at small  $x$ . The asymmetry preferred by the combination of the currently available LHC and Tevatron data turns out to be negative at  $x < 0.01$ , while the Regge-like limit with a vanishing  $I(x)$  can still be recovered at  $x \lesssim 10^{-5}$ , cf. the ABMP15 results in Fig.12. The CT14 analysis only includes the Tevatron forward DY data, but also confirms the negative trend in  $I(x)$  at small  $x$ , with errors in  $I(x)$  being substantially larger than those from ABMP15.

Finally, an important issue is the theoretical accuracy which is employed in the description of the DY data. There are significant differences as shown in Tab. 4 and these cause an additional spread in the fit quality and the results for the PDFs when comparing different NNLO PDF sets.

### 3.4.2. Strange-quark distribution

The main information on the strange sea distribution comes from charm-quark production in neutrino-induced charged-current DIS experiments. The publication of data from CHORUS and NOMAD has recently enlarged the statistics available for those experiments. As a net result, the uncertainty in the strange PDF is now reduced down to a few percent at  $x \sim 0.1$  (cf. Fig. 13). However, at small  $x$  the strange sea distribution is still poorly known in view of the restricted kinematics of the production of charm quarks from fixed targets. Supplementary information on the strange sea at small  $x$  can be obtained from the associated production of charm quarks and  $W$  bosons in the  $pp$  collisions at the LHC. A constraint from collider data on  $W + c$  is potentially less sensitive to the  $c$ -quark fragmentation model compared to the one from semi-leptonic decays of charm, which plays major role in the existing fixed-target DIS experiments. The  $W + c$  data collected by ATLAS and CMS prefer a somewhat enhanced strange sea as compared to the fixed-target determination, cf. Fig. 13. However, the NNLO QCD corrections to this process are still unknown. They are not taken into account in the analysis of  $W + c$  data so far and may have a substantial influence on the fit. The strange sea extracted by ATLAS from an analysis of the combined inclusive data on the  $W$ - and  $Z$ -boson production is even further enhanced, which suggests a restoration of SU(3) flavor symmetry in the sea distributions. However, the accuracy of this determination is poor due to a limited potential of the inclusive data in disentangling the quark flavors. Therefore, the ATLAS result is in fact comparable with other determinations within uncertainties.

In general, the existing experimental constraints on the strange PDF are relatively poor. Therefore, the results of various determinations demonstrate a significant spread, which is mainly driven by the data selection. An additional spread between results of earlier PDF analyses appears due to implementation issues. In particular, the strong strange-sea suppression observed in the NNPDF analysis [152] is related to the error in Eq. (34) of Ref. [152] for the DIS charm-quark production cross section. The CT10 analysis [50], which reported an enhanced strange sea, is flawed due to a wrong sign for the structure function  $xF_3$  [3]. This error also concerns the earlier results on the strange–anti-strange asymmetry [172]. Finally, the MSTW [173] analysis suffered from an error in the gluon-induced contribution to the NLO QCD correction for the charged-current DIS charm-quark production, which was corrected in MMHT14 [6].

## 3.5. Nuclear corrections

Many global PDF analyses make use of data with deuterium targets, such as lepton-deuteron DIS and proton-deuteron DY, as a way of obtaining stronger constraints on the flavor dependence of PDFs that are not possible with proton data alone. The use of deuterium data requires that one take into account differences between PDFs in the deuteron and those in the free nucleon, which arise from effects such as nuclear Fermi motion and binding of the nucleons in the nucleus, as well as nucleon off-shell corrections and nuclear shadowing. While some analyses assume that nuclear corrections in the deuteron are negligible, a number of recent global PDF studies have incorporated nuclear effects into their analyses [1, 5, 48, 169, 174–176].

Generally, the nuclear effects become increasingly important at large values of  $x$  ( $x \gtrsim 0.4$ ), as




FIG. 14: (Left panel) Ratio of deuteron to isoscalar nucleon structure functions  $F_2^d/F_2^N$  computed from the CJ15 PDFs [1] for different values of  $Q^2$ . The pink envelope represents the fit uncertainties for  $Q^2 = 10 \text{ GeV}^2$ . The downturn in the ratio at  $Q^2 = 2 \text{ GeV}^2$  is due to target mass corrections. (Right panel) Impact on the  $d/u$  ratio from the CJ15 fit [1] (red band) of removing the deuterium nuclear corrections (cross-hatched band), and omitting all deuterium data (green band).

Fig. 14 illustrates for the ratio of the deuteron to isoscalar nucleon structure functions. In this region the nuclear PDFs can be computed through convolutions of the bound nucleon PDFs and nuclear smearing functions describing the momentum distributions of nucleons in the deuteron. The latter can be expressed in terms of deuteron wave functions, calculated from modern potentials based on high-precision fits to nucleon–nucleon scattering data. These potentials differ primarily in their treatment of the short range  $NN$  interaction, and the different strengths of the high-momentum tails of the wave functions translate directly to the magnitude of the nuclear corrections at large  $x$  [175, 177].

The nucleon off-shell corrections, on the other hand, are somewhat more model dependent, and several model studies have been performed to estimate their effect on nuclear PDFs [178–182]. Some earlier PDF analyses [169, 175, 176] used specific physics-motivated models for the off-shell corrections, while more recent approaches have fitted the off-shell parameters directly to data [1, 181]. Other analyses [6, 183] have attempted to parametrize the entire nuclear correction in terms of a universal,  $Q^2$ -independent function, without appealing to physical constraints. In this approach, to account for the effects of Fermi smearing a functional form must be used that produces the steep rise in the  $F_2^d/F_2^N$  ratio at high  $x$ , such as with a logarithm raised to a high power [183].

The effects of the nuclear corrections are most directly visible in the extraction of the  $d$ -quark PDF at large  $x$ , see [1]. Figure 14 shows that omitting nuclear smearing effects in the deuteron leads to an overestimated  $d/u$  ratio at  $x \gtrsim 0.6$ . In fact, omitting nuclear corrections induces a strong tension between the SLAC deuteron DIS data (see, e.g., [184]) and the recent high precision  $W$ -boson asymmetry data from the DØ collaboration at the Tevatron [20], which are sensitive to the  $d$ -quark PDF in a similar large- $x$  range as the SLAC data, but are not affected by nuclear corrections. It also causes an artificial deformation of the  $d$ -quark distribution, leading to essentially uncontrolled systematic errors when quark distributions are needed beyond the  $x$  range constrained by the data. This illustrates not only the theoretical but also the phenomenological need for such

corrections when considering data at large  $x$ . Of course, one can choose to avoid nuclear effects altogether by using only proton data; however, doing so increases the uncertainty on the  $d$ -quark PDF at both small and large values of  $x$ , as Fig. 14 illustrates. Additional details concerning the role of nuclear corrections when using deuterium target data in global fits can be found in Ref. [1]. The extrapolation of nuclear effects from the deuteron to heavy nuclei is unclear, especially in view of the differences between the off-shell quark deformation fitted using deuteron targets [1] and using the ratio of heavy nuclei to deuteron structure functions [181]. Care should also be exercised when using neutrino-nucleus scattering data to obtain, for example, constraints on strange-quark PDFs [170] due to the currently poor understanding of the interaction dynamics of the final state heavy-quark propagating through the target nucleus.

### 3.6. Software and tools

Data used in the PDF fits cover a wide range of kinematics and stem from a large number of different scattering processes. In order to achieve an accurate theoretical description of both the PDF evolution and the hard scattering cross sections, well-tested software and tools are necessary. Benchmark numbers for the PDF evolution have long been established, see e.g., the Les Houches report [185], and open-source evolution codes such as QCDNUM [186, 187] and Hoppet [188] are available in Bjorken  $x$ -space and QCD-Pegasus [106] in Mellin  $N$ -space. This is an important development as it allows to expose the software used in the PDF fits to systematic validation, the need of which can be illustrated with recent theory improvements published by various groups. For example, MSTW [173] has tested its NNLO evolution code against QCD-Pegasus [106] and corrected the implementation of one of the heavy-quark OMEs.

For the hard scattering cross sections of the various processes, fast fitting methods like fastNLO [189, 190] and APPLGrid [160] have been developed. In addition, some groups have also published open-source code for the theory predictions of all physical cross sections employed in their analyses. The ABM11 and ABM12 fits [2, 48] use OPENQCDRAD [191] code, which is publicly available. The HERAPDF2.0 fit [4] relies on the QCD fit platform xFitter (formerly known as HERAFitter) [149, 150], which is an open-source package that provides a framework for the determination of PDFs and enables the choice of theoretical options for obtaining PDF-dependent cross section predictions. In particular, xFitter allows for a choice of different available schemes for treatment of heavy quarks in DIS. In Mellin  $N$ -space, an efficient method exists [192, 193] which improves on that by [194] and which has been widely used in analyses, e.g. [193]. However, no code has been made publicly available so far.

Given the increasing precision of PDF analyses, which is driven by the accuracy of the experimental data, there is ongoing demand to provide theoretical predictions that are as precise as possible. This has stimulated recent checks of the analysis software used by various groups and has resulted in a number of documented improvements. This list includes, for example, NNPDF2.1 [152] which used an incorrect cross section for charged-current DIS that was off by a factor of two for low scales due to an additional factor of  $(1 + m_c^2/Q^2)$  in Eq. (34) of Ref. [152], see Sec. 4.1 in Ref. [48] for a detailed discussion. The MSTW [173] analysis had omitted the gluon Wilson coefficient at NLO in the same process, and corrected it in MMHT14 [6], while CT14 [3] has reported a computational error in previous analyses involving a term with a wrong sign in the treatment of heavy-quark mass effects in charged-current DIS. The impact of these errors on the determination of the PDFs for strange sea has been discussed above.

This illustrates that there is a continued need for benchmarking of hard scattering cross sections of relevance for PDF determinations in order to consolidate the accuracy of theory predictions for

PDF sets	$\alpha_s(M_Z)$	method of determination
ABM12 [2]	$0.1132 \pm 0.0011$	fit at NNLO
CJ15 [1]	$0.1183 \pm 0.0002$	fit at NLO
CT14 [3]	0.118	assumed at NNLO
HERAPDF2.0 [4] <sup>a</sup>	$0.1183^{+0.0040}_{-0.0034}$	fit at NLO
JR14 [5]	$0.1136 \pm 0.0004$	dynamical fit at NNLO
	$0.1162 \pm 0.0006$	standard fit at NNLO
MMHT14 [9] <sup>b</sup>	0.118	assumed at NNLO
NNPDF3.0 [7]	0.118	assumed at NNLO
PDF4LHC15 [8]	0.118	assumed at NNLO

<sup>a</sup>In detail HERAPDF2.0 obtains at NLO  $\alpha_s(M_Z) = 0.1183 \pm 0.0009(\text{exp}) \pm 0.0005(\text{model/parameterisation}) \pm 0.0012(\text{hadronisation})^{+0.0037}_{-0.0030}(\text{scale})$ , which have been added in quadrature in the table entry. The HERAPDF2.0 central variant uses a fixed value  $\alpha_s(M_Z) = 0.118$ .

<sup>b</sup>MMHT14 obtains  $\alpha_s(M_Z) = 0.1172 \pm 0.0013$  at NNLO as a best fit.

TABLE 5: Values of  $\alpha_s(M_Z)$  obtained or used in the nominal PDF sets of the various groups.

those observables. In this respect, open-source software may facilitate future theory improvements and may help to establish standards for precision theory predictions.

#### 4. STRONG COUPLING CONSTANT

The value of the strong coupling constant  $\alpha_s(M_Z)$  has a direct impact on the size of a number of cross sections at the LHC, such as Higgs boson production, see Sec. 5, and is therefore an important parameter. Due to QCD factorization,  $\alpha_s$  exhibits a significant correlation with the gluon PDF and also with the charm-quark mass. Therefore, the strong coupling constant has come to require particular attention in the context of global PDF analyses.

Current precision determinations of  $\alpha_s(M_Z)$  require NNLO accuracy in QCD because of the small uncertainties in the experimental data analyzed and the significantly reduced dependence from the variation of the renormalization scale indicating the uncertainty due to the truncation of the perturbative series. Extractions of  $\alpha_s$  at NLO typically yield  $\alpha_s(M_Z) \simeq 0.118$ , however, the NLO scale uncertainty is large, giving sizable variations  $\Delta\alpha_s(M_Z) = 0.005$  for  $\mu_r \in [Q/2, 2Q]$  in DIS analyses. Determinations of  $\alpha_s$  to NNLO accuracy benefit from a significantly reduced renormalization scale dependence, but generally result in smaller central values for  $\alpha_s(M_Z)$ , with shifts downwards from NLO to NNLO of a few percent in DIS analyses. Beyond NNLO, the perturbative expansion converges, as illustrated in DIS in a valence analysis [42] at N<sup>3</sup>LO which yields  $\alpha_s(M_Z) = 0.1141^{+0.0020}_{-0.0022}$ , in agreement with the NNLO values listed in Tab. 6.

Of course, measurements of  $\alpha_s(M_Z)$  are not limited to global fits of PDFs, but stem from a large number of different processes and methods at different scales, see, e.g., [195–197] for discussions



	year	$\alpha_s(M_Z)$	method/data sets/reference
SY	2001	$0.1166 \pm 0.0013$	$F_2^{ep}$ [199]
	2001	$0.1153 \pm 0.0063$	$xF_3^{\nu N}$ (DIS off heavy nuclei) [199]
A02	2002	$0.1143 \pm 0.0020$	[200]
MRST03	2003	$0.1153 \pm 0.0020$	[201]
BBG	2004(06,12)	$0.1134^{+0.0019}_{-0.0021}$	valence analysis, NNLO [42, 202, 203]
GRS	2006	0.112	valence analysis, NNLO [204]
AMP06	2006	$0.1128 \pm 0.0015$	[205]
JR	2008	$0.1128 \pm 0.0010$	dynamical approach [206]
	2008	$0.1162 \pm 0.0006$	with jet hadroproduction at NLO [206]
ABKM	2009	$0.1135 \pm 0.0014$	$n_f = 3$ FFNS heavy quark scheme [174]
	2009	$0.1129 \pm 0.0014$	BMSN heavy quark scheme [174]
MSTW	2009	$0.1171 \pm 0.0014$	[207]
ABM11 <sub>J</sub>	2010	$0.1134 - 0.1149 \pm 0.0012$	with jet production at Tevatron (NLO) [208]
NNPDF2.1	2011	$0.1174 \pm 0.0006 \pm 0.0001$	with data for DIS off heavy nuclei [209, 210]
ABM11	2012	$0.1134 \pm 0.0011$	[48]
ABM12	2013	$0.1132 \pm 0.0011$	[2]
Thorne	2013	0.1136	DIS, Drell-Yan data; with account of higher twist [211]
CT10	2013	0.1140	with jet hadroproduction data [50]
JR	2014	$0.1136 \pm 0.0004$	dynamical approach [5]
	2014	$0.1162 \pm 0.0006$	standard fit [5]
CT14	2015	$0.1150^{+0.0060}_{-0.0040}$	$\Delta\chi^2 > 1$ and with data for DIS off heavy nuclei [3]
MMHT	2015	$0.1172 \pm 0.0013$	with data for DIS off heavy nuclei [212]

TABLE 6: Determinations of  $\alpha_s(M_Z)$  values at NNLO from QCD analyses of the deep-inelastic world data and, partly, including other hard scattering data. For recent compilations, see [195–197].

and comparisons. Here we restrict ourselves to issues of  $\alpha_s$  arising in PDF fits. In Tab. 5 we give an overview of the  $\alpha_s$  values currently used in the PDF analyses. There, two aspects are important. Firstly, some PDF analyses leave  $\alpha_s$  as a free parameter in their fits, which obviously allows one to control its correlation with other PDF parameters and avoids potential biases. Secondly, among the NNLO values of  $\alpha_s(M_Z)$  used there exists a large spread of  $\alpha_s$  values, ranging from  $\alpha_s(M_Z) = 0.1132$  to  $0.1183$ . Some of those fitted values of  $\alpha_s(M_Z)$  are significantly smaller than, for example, an average provided by the PDG [39] in 2014, which gives  $\alpha_s(M_Z) = 0.1185 \pm 0.0006$  at NNLO, and is often quoted as a motivation for fixing  $\alpha_s(M_Z) = 0.118$  as in some entries in Tab. 5. In the recent 2015 update, the PDG [198] reports the value  $\alpha_s(M_Z) = 0.1181 \pm 0.0013$  with

PDF set	theory accuracy	data sets used
CT14 [3]	NLO with the scale set to the individual $p_T$ of jet	Tevatron + LHC
MMHT14 [9]	NLO + threshold corrections [213]	Tevatron
NNPDF3.0 [7]	NLO + threshold corrections [214]	Tevatron + LHC (“safety cuts”)

TABLE 7: The jet data sets and the theory approximations used in the NNLO PDF fits.

the uncertainty increased by a factor of two.

While the potential agreement or disagreement with the PDG average is beyond the scope of this study, it is instructive to focus on  $\alpha_s(M_Z)$  measurements from PDF analyses as listed in Tab. 6 which have been performed since the NNLO QCD corrections in DIS first became available. This series of measurements has led to  $\alpha_s(M_Z)$  values which are not only mostly lower than the PDG average, but also exhibit a large spread in the range  $\alpha_s(M_Z) = 0.1120 - 0.1174$ . This spread is significant given the small size of the experimental uncertainties in the data. As it turns out, the differences in the values of  $\alpha_s(M_Z)$  can be traced back to different data sets used or to different theory assumptions applied, as indicated in Tab. 6.

For instance, the inclusion of data for the hadro-production of jets, e.g., from the LHC, does have an impact on the value of  $\alpha_s(M_Z)$  and can therefore provide valuable constraints. However, it is important to note that the perturbative QCD corrections to the hard scattering cross section are only known completely to NLO, while the exact NNLO result for the  $gg$  channel [14] and approximations based on soft gluon enhancement [213–216] indicate corrections as large as 15%–20%. Those corrections and their magnitude depend, of course, on the details of the kinematics and on the jet parameters (jet radius  $R$ ). For high  $p_T$  they are dominated by threshold logarithms  $\ln(p_T)$  accompanied by logarithms  $\ln(R)$  for small jet radii [216].

The  $\alpha_s(M_Z)$  values in PDF analyses currently determined with the help of jet data (cf. Tab. 6) are, strictly speaking, valid to NLO accuracy only and therefore subject to significantly larger theory uncertainties due to the variation of the renormalization scale. The various groups employ different approaches in their NNLO analyses to cope with this inconsistency, such as using dynamical scales or applying some variant of threshold corrections, as detailed in Tab. 7. As a results of these efforts, the gluon PDF and  $\alpha_s$  obtained, for example, in the MMHT14 and NNPDF3.0 analyses are in a good agreement.

Different modeling of important theory aspects, such as whether or not to include target mass corrections, higher twist contributions and nuclear corrections in the description of DIS data, or whether or not to use a VFNS in the description of DIS heavy-quark data, can account for the range of  $\alpha_s(M_Z)$  in Tab. 6. With largely similar model assumptions, NNPDF2.1 [209, 210], MSTW [207] and MMHT [212] obtained the range  $\alpha_s(M_Z) = 0.1171 - 0.1174$ . All these choices can lead to systematic shifts of the value of  $\alpha_s(M_Z)$ . Let us briefly mention some of the issues in detail.

Higher twist contributions do have a big impact, because these terms are fitted within a combined analysis. Alternatively, the part of the DIS data significantly affected by these terms has to be removed by suitable kinematical cuts on the scale  $Q^2$  and center-of-mass energies  $W^2$ . In a variant of the ABM11 analysis [48], higher twist terms have been omitted and the cuts  $W^2 > 12.5 \text{ GeV}^2$  and  $Q^2 > 2.5 \text{ GeV}^2$  as used by MSTW [207] have been applied. This resulted in a sizable shift

upwards to  $\alpha_s(M_Z^2) = 0.1191 \pm 0.0016$  in line with earlier studies in [217]. Yet more conservative cuts of  $W^2 > 12.5 \text{ GeV}^2$  and  $Q^2 > 10 \text{ GeV}^2$  in the ABM11 variant with higher twist terms set to zero led to  $\alpha_s(M_Z^2) = 0.1134 \pm 0.0008$ , well in agreement with the nominal value in the ABM11 analysis, cf. Tab. 6. Thus, in PDF analyses without account of higher twist contributions to DIS data such tight cuts are essential. We also note that NNPDF3.0 [7] uses a cut of  $Q^2 > 5 \text{ GeV}^2$  which removes some but not all data affected by the higher twist contributions.

Higher order constraints from fixed-target DIS data can also lead to shifts in  $\alpha_s(M_Z)$  [43]. For instance, NMC has measured the DIS differential cross sections and extracted the DIS structure functions  $F_2^{\text{NMC}}$  [218]. At the time of the NMC analysis, however, the relevant DIS corrections to  $\mathcal{O}(\alpha_s^3)$  [41] were not available (see discussion after Eq. (5) above). This information is, however, important and has to be taken into account now. In case of fitting  $F_2^{\text{NMC}}$  and not describing  $F_L(x, Q^2)$  at NNLO, much larger values of  $\alpha_s(M_Z^2)$  are obtained. It is therefore strongly recommended to fit the published differential scattering cross sections using  $F_L(x, Q^2)$  at  $\mathcal{O}(\alpha_s^3)$ . Presently, the MMHT [212] analysis uses  $F_L(x, Q^2)$  only at NLO. One should note, however, that the values of  $F_L(x, Q^2)$  at NNLO are significantly different in the small- $x$  region.

Finally, great care needs to be exercised when DIS data off nuclei are included in global fits, see Sec. 3. Details of modeling of nuclear corrections can in fact also cause systematic shifts in the value of  $\alpha_s(M_Z)$ . Therefore, Tab. 6 indicates if scattering data on heavy nuclei have been included in the determination. For example, MMHT [212] has reported a comparatively high value of  $\alpha_s(M_Z)$  as a consequence of fitting the NuTeV  $\nu\text{Fe}$  DIS data [163]. In general, determinations of  $\alpha_s(M_Z)$  should be based upon, or at least cross-checked with, fits using proton and deuteron DIS data only.

## 5. CROSS SECTION PREDICTIONS FOR THE LHC

### 5.1. Higgs boson production

The dominant production mechanism for the SM Higgs boson at the LHC is the gluon-gluon fusion process. The large size of the QCD radiative corrections to the inclusive cross section at NLO, see, e.g. Ref. [219], together with the sizable scale uncertainty have motivated systematic theory improvements. In the effective theory based on the limit of a large top-quark mass ( $m_t \rightarrow \infty$ , integrating out the top-quark loop, but using the full  $m_t$  dependence in the Born cross section), this has led to the computation of the corresponding corrections at NNLO [220–222] and even to N<sup>3</sup>LO in QCD [89, 223]. This shows an apparent, if slow, convergence of the perturbative expansion, along with greatly reduced sensitivity to the choice for the renormalization and factorization scales  $\mu_r$  and  $\mu_f$ . At N<sup>3</sup>LO the total scale variation amounts to 3% and estimates of the four-loop corrections support these findings [224].

This leaves, as the largest remaining source of uncertainties in the predictions of the physical cross section, the input for the strong coupling constant  $\alpha_s$  and the PDFs. Despite the impressive progress in theory and experiment, the situation resembles that after the completion of the NLO QCD corrections, when it was pointed out in Ref. [219] that one of the main residual uncertainties in the predictions was due to the gluon PDF.

In Tab. 8 we summarize the PDF dependence of the inclusive cross section  $\sigma(H)^{\text{NNLO}}$  in the effective theory (i.e., in the limit of  $m_t \gg m_H$ ) at  $\sqrt{s} = 13 \text{ TeV}$  for a Higgs boson mass  $m_H = 125.0 \text{ GeV}$ ,  $\mu_r = \mu_f = m_H$ , and  $m_t^{\text{pole}} = 172.5 \text{ GeV}$  with uncertainties  $\sigma(H)^{\text{NNLO}} + \Delta\sigma(\text{PDF} + \alpha_s)$ , and compare the results for various PDF sets. The PDF uncertainties are typically given at the

PDF sets	$\sigma(H)^{\text{NNLO}}$ [pb] nominal $\alpha_s(M_Z)$	$\sigma(H)^{\text{NNLO}}$ [pb] $\alpha_s(M_Z) = 0.115$	$\sigma(H)^{\text{NNLO}}$ [pb] $\alpha_s(M_Z) = 0.118$
ABM12 [2]	$39.80 \pm 0.84$	$41.62 \pm 0.46$	$44.70 \pm 0.50$
CJ15 [1] <sup>a</sup>	$42.45^{+1.73}_{-1.12}$	$39.48^{+1.60}_{-1.04}$	$42.45^{+1.73}_{-1.12}$
CT14 [3] <sup>b</sup>	$42.33^{+1.43}_{-1.68}$	$39.41^{+1.33}_{-1.56}$ (40.10)	$42.33^{+1.43}_{-1.68}$
HERAPDF2.0 [4] <sup>c</sup>	$42.62^{+0.35}_{-0.43}$	$39.68^{+0.32}_{-0.40}$ (40.88)	$42.62^{+0.35}_{-0.43}$
JR14 (dyn) [5]	$38.01 \pm 0.34$	$39.34 \pm 0.22$	$42.25 \pm 0.24$
MMHT14 [6]	$42.36^{+0.56}_{-0.78}$	$39.43^{+0.53}_{-0.73}$ (40.48)	$42.36^{+0.56}_{-0.78}$
NNPDF3.0 [7]	$42.59 \pm 0.80$	$39.65 \pm 0.74$ (40.74 $\pm$ 0.88)	$42.59 \pm 0.80$
PDF4LHC15 [8]	$42.42 \pm 0.78$	$39.49 \pm 0.73$	$42.42 \pm 0.78$

<sup>a</sup> The CJ15 PDFs have been determined at NLO accuracy in QCD, with the cross section errors here computed for  $T = 10$  (i.e., the errors provided by the CJ15 set are multiplied 10).

<sup>b</sup> The PDF uncertainties quoted by CT14 denote the 90% c.l. and should be reduced by a factor of 1.645 for comparison with the 68% c.l. uncertainties quoted by other groups.

<sup>c</sup> The model uncertainties of the HERAPDF20\_NNLO\_VAR set are not included in the uncertainty estimates.

TABLE 8: The Higgs cross section at NNLO in QCD (computed in the effective theory) at  $\sqrt{s} = 13$  TeV for  $m_H = 125.0$  GeV at the nominal scale  $\mu_r = \mu_f = m_H$  with the PDF (and, if available, also  $\alpha_s$ ) uncertainties. The columns correspond to different choices for the central value of  $\alpha_s(M_Z)$ . The numbers in parenthesis are obtained using PDF sets with the value of  $\alpha_s(M_Z)$  fixed.

1 $\sigma$  c.l. We list the results for  $\sigma(H)^{\text{NNLO}}$  using either the values for the strong coupling constant  $\alpha_s(M_Z)$  at NNLO, corresponding to the respective PDF set, or fixed values of  $\alpha_s(M_Z) = 0.115$  and  $\alpha_s(M_Z) = 0.118$ . This is done to illustrate the fact that in some PDFs the value of  $\alpha_s(M_Z)$  is not obtained from a fit to data (including faithful uncertainties) but fixed beforehand, e.g., to the world average [39]. Often the same fixed value of  $\alpha_s(M_Z)$  is chosen at NLO and at NNLO independent of the order of perturbation theory, see also Sec. 4. Table 8 shows a large spread for predictions from different PDFs with a range  $\sigma(H)^{\text{NNLO}} = 38.0 - 42.6$  pb using the nominal value of  $\alpha_s(M_Z)$ . Specifically, the PDF and  $\alpha_s$  differences between different sets are up to 11% and are significantly larger than the residual scale uncertainty due to N<sup>3</sup>LO QCD corrections. In addition, the cross sections shift in the range  $\sigma(H)^{\text{NNLO}} = 39.0 - 44.7$  pb if a fixed value of  $\alpha_s(M_Z)$  in the range  $\alpha_s(M_Z) = 0.115 - 0.118$  is used. This amounts to a relative difference of more than 13% and contradicts the most recent estimates of the combined PDF and  $\alpha_s$  uncertainties in the inclusive cross section [89], which quotes 3.2%. In general, the findings underpin the importance of controlling the accuracy and the correlation of the strong coupling constant with the PDF parameters in fits.

Of particular interest is the impact of additional parameters in the PDF fits, such as the charm-

$m_c^{\text{pole}}$ [GeV]	$\alpha_s(M_Z)$ (best fit)	$\chi^2/\text{NDP}$ (HERA data [148])	$\sigma(H)^{\text{NNLO}}$ [pb] best fit $\alpha_s(M_Z)$	$\sigma(H)^{\text{NNLO}}$ [pb] $\alpha_s(M_Z) = 0.118$
1.05	0.1157	73/52	40.65	(41.63)
1.1	0.1159	69/52	40.85	(41.70)
1.15	0.1160	66/52	41.04	(41.78)
1.2	0.1162	64/52	41.25	(41.85)
1.25	0.1164	64/52	41.47	(41.93)
1.3	0.1166	63/52	41.69	(42.00)
1.35	0.1168	63/52	41.93	(42.09)
1.4	0.1171	65/52	42.16	(42.16)
1.45	0.1173	68/52	42.42	(42.24)
1.5	0.1175	73/52	42.64	(42.31)
1.55	0.1177	80/52	42.88	(42.38)
1.6	0.1180	88/52	43.16	(42.46)
1.65	0.1182	99/52	43.34	(42.51)
1.7	0.1184	112/52	43.59	(42.58)
1.75	0.1186	127/52	43.81	(42.63)

TABLE 9: The values of the charm-quark mass (on-shell scheme  $m_c^{\text{pole}}$ ) and the strong coupling  $\alpha_s(M_Z)$  in the MSTW analysis [153] using the set `MSTW2008nnlo_mcrange` together with the value for  $\chi^2/\text{NDP}$  for the HERA data [148] and the Higgs cross section at NNLO in QCD (computed in the effective theory) at  $\sqrt{s} = 13$  TeV for  $m_H = 125.0$  GeV at the nominal scale  $\mu_r = \mu_f = m_H$ . The numbers in parentheses are obtained using the PDF set `MSTW2008nnlo_mcrange_fixasmz` with the value of  $\alpha_s(M_Z)$  fixed to  $\alpha_s(M_Z) = 0.118$ .

quark mass, on the Higgs cross section. The differences in the treatment of heavy quarks and the consequences for the quality of the description of charm-quark DIS data have already been discussed in Sec. 3. ABM12 [2] fits the value of  $m_c(m_c)$  in the  $\overline{\text{MS}}$  scheme and the uncertainties in the charm-quark mass are included in the uncertainties quoted in Tab. 8. Other groups keep a fixed value of the charm-quark mass in the on-shell scheme, cf. Tab. 3, and vary the value of  $m_c^{\text{pole}}$  within some range. Such studies have been performed in the past by NNPDF2.1 [152] and MSTW [153] and more recently by MMHT [9].

In Tabs. 9, 10 and 11 we display the results of these fits together with the values of  $\chi^2/\text{NDP}$  for the DIS charm-quark data [148], mostly computed with `xFitter` [149, 150], as well as the corresponding cross section for Higgs boson production to NNLO accuracy. The MSTW analysis in Tab. 9 shows a linear rise of the cross section for increasing values  $m_c^{\text{pole}} = 1.05 - 1.75$  GeV in the range  $\sigma(H) = 40.6 - 43.8$  pb, which amounts to a variation of more than 7%. Even if

$m_c^{\text{pole}}$ [GeV]	$\alpha_s(M_Z)$ (best fit)	$\chi^2/\text{NDP}$ (HERA data [148])	$\sigma(H)^{\text{NNLO}}$ [pb] best fit $\alpha_s(M_Z)$	$\sigma(H)^{\text{NNLO}}$ [pb] $\alpha_s(M_Z) = 0.118$
1.15	0.1164	78/52 (71/52)	40.48	(42.05)
1.2	0.1166	76/52 (70/52)	40.74	(42.11)
1.25	0.1167	75/52 (76/52)	40.89	(42.17)
1.3	0.1169	76/52 (77/52)	41.16	(42.25)
1.35	0.1171	78/52 (79/52)	41.41	(42.30)
1.4	0.1172	82/52 (83/52)	41.56	(42.36)
1.45	0.1173	88/52 (89/52)	41.75	(42.45)
1.5	0.1173	96/52 (96/52)	41.81	(42.51)
1.55	0.1175	105/52 (106/52)	42.08	(42.58)

TABLE 10: Same as Tab. 9 for the MMHT14 analysis [9] using the set `MMHT2014nnlo_mcrange_nf5` and the best fit value of  $\alpha_s(M_Z)$ . The values of  $\chi^2/\text{NDP}$  for the HERA data [148] are those quoted in [9] for the best fit value of  $\alpha_s(M_Z)$ . The numbers in parentheses are obtained with the value of  $\alpha_s(M_Z)$  fixed to  $\alpha_s(M_Z) = 0.118$ .

PDF sets	$m_c^{\text{pole}}$ [GeV]	$\alpha_s(M_Z)$ (fixed)	$\chi^2/\text{NDP}$ (HERA data [148])	$\sigma(H)^{\text{NNLO}}$ [pb] fixed $\alpha_s(M_Z)$
NNPDF2.1 [152]	$\sqrt{2}$	0.119	65/52	$44.18 \pm 0.49$
	1.5	0.119	78/52	$44.54 \pm 0.51$
	1.6	0.119	92/52	$44.74 \pm 0.50$
	1.7	0.119	110/52	$44.95 \pm 0.51$
NNPDF2.3 [225]	$\sqrt{2}$	0.118	71/52	$43.77 \pm 0.41$
NNPDF3.0 [7]	1.275	0.118	67/52	$42.59 \pm 0.80$

TABLE 11: Same as Tab. 9 for various NNPDF analyses. The values of the strong coupling  $\alpha_s(M_Z)$  have been fixed in those fits. The values of  $\chi^2/\text{NDP}$  for the description of the HERA data have been determined with the FONLL-C [145] scheme.

$\alpha_s(M_Z) = 0.118$  is kept fixed, the cross section varies in the range  $\sigma(H) = 41.6 - 42.6$  pb, which is equivalent to 2%. The best fit in the MSTW analysis with  $\chi^2/\text{NDP} = 63/52$  leads to  $m_c^{\text{pole}} = 1.3$  GeV and  $\alpha_s(M_Z) = 0.1166$ , both of which are lower than the ones of the nominal fit with  $m_c^{\text{pole}} = 1.4$  GeV and  $\alpha_s(M_Z) = 0.1171$ . In Tab. 10 the same study is performed for the MMHT PDFs [9], where the reduced quark mass range  $m_c^{\text{pole}} = 1.15 - 1.55$  GeV still leads to cross section variations  $\sigma(H) = 40.5 - 42.1$  pb (i.e., 4%) for the best fit  $\alpha_s(M_Z)$ , or  $\sigma(H) = 42.1 - 42.6$  pb (i.e.,

1%) for a fixed  $\alpha_s(M_Z) = 0.118$ . The latter case leads to a best fit of  $m_c^{\text{pole}} = 1.2$  GeV with  $\chi^2/\text{NPD} = 70/52$ , which is significantly smaller than the nominal fit with  $m_c^{\text{pole}} = 1.4$  GeV and  $\chi^2/\text{NPD} = 82/52$ .

NNPDF has performed a study of the  $m_c$  dependence in [152], which shows the same trend as for MSTW and MMHT, i.e., the smaller the chosen value of  $m_c^{\text{pole}}$ , the better the goodness-of-fit for the HERA data [148]. In addition, Tab. 11 displays the changes in the charm-quark mass values from  $m_c^{\text{pole}} = \sqrt{2}$  GeV to  $m_c^{\text{pole}} = 1.275$  GeV in the evolution of the NNPDF fits from v2.1 [152] and v2.3 [225] to v3.0 [7], with the obvious correlation of smaller cross sections for Higgs boson production with smaller chosen values of  $m_c^{\text{pole}}$ .

As pointed out already in Sec. 3, on-shell masses  $m_c^{\text{pole}} = 1.2 - 1.3$  GeV as preferred by the goodness-of-fit analyses in Tabs. 9, 10 and 11 for the charm-quark data from HERA [148], are not compatible with the world average of the PDG [39]. Thus, in some PDF fits, the numerical value of the charm-quark mass effectively takes over the role of a “tuning” parameter for the Higgs cross section.

## 5.2. Hadro-production of heavy quarks

### 5.2.1. Top-quark hadro-production: inclusive cross section

The cross section for the hadro-production of top-quark pairs has been measured with unprecedented accuracy at the LHC in Run 1 with  $\sqrt{s} = 7$  TeV and 8 TeV. The inclusive cross section is known to NNLO in QCD [226–229], featuring good convergence of the perturbation series and reduced sensitivity to the renormalization and factorization scales  $\mu_r$  and  $\mu_f$ . These theory predictions adopt the on-shell renormalization scheme for the heavy-quark mass. The conversion to the  $\overline{\text{MS}}$  scheme for the heavy-quark mass has been discussed in Refs. [230–232]. For observables such as the inclusive cross section which are dominated by hard scales  $\mu_r \simeq \mu_f \simeq m_t$ , the theory predictions in terms of the  $\overline{\text{MS}}$  mass for the top quark show an even better scale stability and perturbative convergence.

In a similar study as for Higgs boson production in Tab. 8 we illustrate in Tab. 12 the PDF dependence of the inclusive cross section  $\sigma(t\bar{t})^{\text{NNLO}}$  for various sets with uncertainties  $\Delta\sigma(\text{PDF} + \alpha_s)$ . The computation is performed in the theoretical framework as implemented in the HATHOR code [231]. In Tab. 12 we choose  $\sqrt{s} = 13$  TeV and fix the pole mass  $m_t^{\text{pole}} = 172.0$  GeV and the scales at  $\mu_r = \mu_f = m_t^{\text{pole}}$ . For this fixed value of  $m_t^{\text{pole}}$ , we show the impact of different values for the strong coupling constant at NNLO. We choose  $\alpha_s(M_Z)$  either corresponding to the respective PDF set or fixed to the values 0.115 and 0.118. The results in Tab. 12 display a spread in a range  $\sigma(t\bar{t})^{\text{NNLO}} = 715 - 834$  pb using the nominal value of  $\alpha_s(M_Z)$  for each PDF set, which amounts to a relative range of more than 15%. This decreases to about 6%, if the values of  $\alpha_s(M_Z)$  are fixed to 0.115 or 0.118.

The theoretical predictions at leading order depend parametrically on the strong coupling constant and the top-quark mass to second power, as well as on the convolution of the gluon PDFs,  $\sigma(t\bar{t})^{\text{LO}} \propto (\alpha_s^2/m_t^2)(g \otimes g)$ . Therefore, it is necessary to fully account for the correlations between the top-quark mass, the gluon PDF and the strong coupling when comparing to experimental data. A number of analyses have considered  $t\bar{t}$  hadro-production data. ABM12 [2] has included data for top-quark pair-production in a variant of the fit to determine the  $\overline{\text{MS}}$  mass  $m_t(m_t)$ , keeping the full correlation with  $\alpha_s(M_Z)$  and the gluon PDF. On the other hand, CMS has determined the top-quark

PDF sets	$\sigma(t\bar{t})^{\text{NNLO}}$ [pb] nominal $\alpha_s(M_Z)$	$\sigma(t\bar{t})^{\text{NNLO}}$ [pb] $\alpha_s(M_Z) = 0.115$	$\sigma(t\bar{t})^{\text{NNLO}}$ [pb] $\alpha_s(M_Z) = 0.118$
ABM12 [2]	$715.1 \pm 21.3$	$741.7 \pm 10.3$	$786.5 \pm 10.9$
CJ15 [1] <sup>a</sup>	$786.7^{+32.2}_{-53.3}$	$742.0^{+30.6}_{-50.2}$	$786.7^{+32.2}_{-53.3}$
CT14 [3] <sup>b</sup>	$834.2^{+36.0}_{-36.5}$	$786.7^{+34.0}_{-34.4}$ (791.4)	$834.2^{+36.0}_{-36.5}$
HERAPDF2.0 [4] <sup>c</sup>	$804.1^{+11.4}_{-17.0}$	$757.8^{+10.8}_{-16.0}$ (756.8)	$804.1^{+11.4}_{-17.0}$
JR14 (dyn) [5]	$719.3 \pm 9.1$	$739.6 \pm 5.7$	$784.2 \pm 6.0$
MMHT14 [6]	$831.8^{+13.9}_{-17.5}$	$784.5^{+13.1}_{-16.5}$ (794.8)	$831.8^{+13.9}_{-17.5}$
NNPDF3.0 [7]	$831.8 \pm 15.0$	$784.4 \pm 14.2$ (800.9 $\pm$ 16.5)	$831.8 \pm 15.0$
PDF4LHC15 [8]	$832.5 \pm 16.4$	$785.1 \pm 15.5$	$832.5 \pm 16.4$

<sup>a</sup> The CJ15 PDFs have been determined at NLO accuracy in QCD, with the cross section errors here computed for  $T = 10$  (i.e., the errors provided by the CJ15 set are multiplied 10).

<sup>b</sup> The PDF uncertainties quoted by CT14 denote the 90% c.l. and should be reduced by a factor of 1.645 for comparison with the 68% c.l. uncertainties quoted by other groups.

<sup>c</sup> The model uncertainties of the HERAPDF20\_NNLO\_VAR set are not included in the uncertainty estimates.

TABLE 12: The inclusive cross section for top-quark pair production at NNLO in QCD at  $\sqrt{s} = 13$  TeV for a pole mass of  $m_t^{\text{pole}} = 172.0$  GeV at the nominal scale  $\mu_r = \mu_f = m_t^{\text{pole}}$  with the PDF (and, if available, also  $\alpha_s$ ) uncertainties. The columns correspond to different choices for the central value of  $\alpha_s(M_Z)$ . The numbers in parenthesis are obtained using PDF sets with the value of  $\alpha_s(M_Z)$  fixed.

pole mass as well as the strong coupling constant in a fit which kept all other parameters mutually fixed [233], while Ref. [234] has explored constraints on the gluon PDF from  $t\bar{t}$  hadro-production data using fixed values for  $\alpha_s(M_Z)$  and the pole mass  $m_t^{\text{pole}}$ .

In the global analyses by MMHT14 [6] and NNPDF3.0 [7] those data were also used to fit  $\alpha_s(M_Z)$  and the gluon PDF. These analyses employ a fixed value for the pole mass  $m_t^{\text{pole}}$ , which is motivated by precisely measured top-quark masses from kinematic reconstructions, i.e., Monte Carlo masses, but does not account for the above mentioned correlation with  $\alpha_s(M_Z)$  and the gluon PDF. Moreover, the Monte Carlo mass requires additional calibration [235].

For the inclusive top-quark cross section we explore in Tabs. 13 and 14 the implicit dependence of the cross section on the charm-quark mass  $m_c$  used in the GM-VFNS of the PDF fits and list the corresponding values of  $\chi^2/\text{NPD}$  for the DIS charm-quark data [148]. This is analogous to the study for the Higgs cross section in Tabs. 10 and 11. For MMHT [9] the best fit with  $m_c^{\text{pole}} = 1.25$  GeV and  $\alpha_s(M_Z) = 0.1167$  leads to an inclusive cross section of  $\sigma(t\bar{t})^{\text{NNLO}} = 814$  pb, which is 2% lower than the value obtained for the nominal MMHT fit, cf. Tab. 12. Likewise, the changes in the NNPDF fits from v2.1 [152] and v2.3 [225] to v3.0 [7] are documented in Tab. 14. The effects amount to almost 2% when comparing  $\sigma(t\bar{t})^{\text{NNLO}}$  for the best fit of NNPDF2.1



$m_c^{\text{pole}}$ [GeV]	$\alpha_s(M_Z)$ (best fit)	$\chi^2/\text{NDP}$ (HERA data [148])	$\sigma(t\bar{t})^{\text{NNLO}}$ [pb] best fit $\alpha_s(M_Z)$	$\sigma(t\bar{t})^{\text{NNLO}}$ [pb] $\alpha_s(M_Z) = 0.118$
1.15	0.1164	78/52 (71/52)	810.2	(835.8)
1.2	0.1166	76/52 (70/52)	813.0	(835.4)
1.25	0.1167	75/52 (76/52)	814.0	(834.8)
1.3	0.1169	76/52 (77/52)	816.5	(834.2)
1.35	0.1171	78/52 (79/52)	819.0	(833.4)
1.4	0.1172	82/52 (83/52)	819.0	(831.8)
1.45	0.1173	88/52 (89/52)	820.2	(831.5)
1.5	0.1173	96/52 (96/52)	818.8	(830.0)
1.55	0.1175	105/52 (106/52)	821.0	(829.0)

TABLE 13: The values of the charm-quark mass (on-shell scheme  $m_c^{\text{pole}}$ ) and the strong coupling  $\alpha_s(M_Z)$  in the MMHT14 analysis [9] together the inclusive cross section for top-quark pair production at NNLO in QCD computed with the set `MMHT2014nnlo_mcrange_nf5` at  $\sqrt{s} = 13$  TeV for a pole mass of  $m_t^{\text{pole}} = 172.0$  GeV at the nominal scale  $\mu_r = \mu_f = m_t^{\text{pole}}$ . The values of  $\chi^2/\text{NDP}$  for the HERA data [148] are those quoted in [9] for the best fit value of  $\alpha_s(M_Z)$ . The numbers in parentheses for the cross section and  $\chi^2/\text{NDP}$  are obtained using the PDF set with the value of  $\alpha_s(M_Z)$  fixed to  $\alpha_s(M_Z) = 0.118$ .

PDF sets	$m_c^{\text{pole}}$ [GeV]	$\alpha_s(M_Z)$ (fixed)	$\chi^2/\text{NDP}$ (HERA data [148])	$\sigma(t\bar{t})^{\text{NNLO}}$ [pb] fixed $\alpha_s(M_Z)$
NNPDF2.1 [152]	$\sqrt{2}$	0.119	65/52	$847.1 \pm 16.3$
	1.5	0.119	78/52	$850.8 \pm 14.3$
	1.6	0.119	92/52	$842.9 \pm 13.6$
	1.7	0.119	110/52	$840.1 \pm 14.1$
NNPDF2.3 [225]	$\sqrt{2}$	0.118	71/52	$835.7 \pm 14.9$
NNPDF3.0 [7]	1.275	0.118	67/52	$831.8 \pm 15.0$

TABLE 14: Same as Tab. 13 for various NNPDF analyses. The values of the strong coupling  $\alpha_s(M_Z)$  have always been fixed in those fits. The values of  $\chi^2/\text{NDP}$  for the description of the HERA data have been determined with the FONLL-C [145] scheme.

with  $m_c^{\text{pole}} = \sqrt{2}$  GeV and  $\alpha_s(M_Z) = 0.119$  to the cross section computed with NNPDF3.0 with  $m_c^{\text{pole}} = 1.275$  GeV and  $\alpha_s(M_Z) = 0.118$ . In both Tables 13 and 14 there is a correlation showing decreasing cross sections with decreasing values of  $m_c^{\text{pole}}$ , although less pronounced than in the case of the Higgs production cross section. The potential bias in the prediction of the inclusive




FIG. 15: (Left panel) Predictions for top-quark pair production cross sections at approximate NNLO as a function of the top-quark rapidity using different PDFs at NNLO with the respective PDF uncertainty (depicted by bands of different style). (Right panel) The acceptance and extrapolation estimators with the respective PDF uncertainties, obtained by using different PDF sets.

top-quark pair production cross section due to a particular “tuning” of the value of the charm-quark mass for some PDFs is, however, of the same order of magnitude or larger than the quoted PDF uncertainties. Therefore, this needs to be accounted for as an additional modeling uncertainty.

### 5.2.2. Top-quark hadro-production: differential distributions

The differential cross section of the top-quark pair production is also known to NNLO in QCD [236]. Publicly available codes such as `Difftop` [237] provide differential distributions to approximate NNLO accuracy based on soft-gluon threshold resummation results. We use `Difftop` to calculate the distribution in the top-quark rapidity  $y_t$  for proton-proton collisions at  $\sqrt{s} = 13$  TeV at NNLO<sub>approx</sub> accuracy using the ABM12, CT14, MMHT14, NNPDF3.0, and the PDF4LHC15 PDF sets at NNLO with their respective  $\alpha_s$  values. Here, we take the top-quark pole mass to be  $m_t^{\text{pole}} = 172.5$  GeV, following the preferences in the LHC analyses. The renormalization and factorization scales are set to  $m_t^{\text{pole}}$  and the choice of a dynamical scale does not change the following discussions.

By using differential cross sections, not only the sensitivity of top-quark pair production to the PDFs can be estimated, but also possible effects on the experimental acceptance by changing the PDF choice. In the experimental analysis, the PDF dependent acceptance corrections arise mostly from the PDF dependent normalization of the production cross section and originate from the phase space regions uncovered by the detector. Usually, the acceptances are determined by using Monte Carlo simulations as a ratio of the number of reconstructed events in the fiducial volume of the detector (visible phase space) to the number of events generated in the full phase space. In the case of top-quark pair production, the visible (full) phase space would correspond to the top-quark rapidity range of  $|y_t| < 2.5$  ( $|y_t| < 3$ ). Here, an acceptance estimator and a related extrapolation factor are calculated by using `Difftop` predictions for the respective cross section ratios  $\sigma_{\text{vis}}/\sigma_{\text{tot}}$

and  $\sigma_{\text{unmeasured}}/\sigma_{\text{tot}}$ . Such estimators are not expected to describe the true experimental efficiency, but are helpful for drawing conclusions about PDF related effects.

The predictions of the top-quark rapidity and the acceptance estimates obtained by using `DiffTop` with different PDFs are shown in Fig. 15. The largest difference in the global normalization of the predicted cross sections is observed if the ABM12 PDFs are used instead of the CT14, NNPDF3.0 or MMHT14 sets. The origin of this effect is again the smaller nominal value of  $\alpha_s$  in ABM12 in combination with a smaller gluon PDF in the  $x$  range relevant to top-quark pair production at  $\sqrt{s} = 13$  TeV. The corresponding acceptance estimators and their uncertainties, obtained from the error propagation of the corresponding PDF uncertainties at 68% c.l., however, demonstrate significant differences also in the expected acceptance corrections, obtained by using ABM12 alternative to other PDFs.

The recent PDF4LHC recommendation [8] for calculation of the acceptance corrections for precision observables, such as the top-quark pair-production cross section in the LHC Run 2 data taking period, is to use the set PDF4LHC15\_100, which is obtained by averaging the CT14, MMHT14 and NNPDF3.0 PDFs. While the central prediction obtained by using PDF4LHC15 is indeed very close to those obtained with the CT14, MMHT14 or NNPDF3.0 PDFs, the error on the corresponding acceptance estimator somewhat underestimates the acceptance spread of the individual PDFs with their uncertainties. Furthermore, it does not cover the difference in the acceptances to the one using the ABM12 PDF. Therefore, for the conservative estimate of the acceptance correction and its uncertainty, as demanded in the measurement of SM precision observables, the use of the PDF4LHC15\_100 set would lead to a significant underestimation of the uncertainty on the resulting cross section measurement.

A further conclusion from Fig. 15 is that in the case of top-quark pair production, once calculational speed is needed, it seems to be sufficient to consider a reduced choice of PDF sets. For instance, instead of using the averaged set PDF4LHC15\_100 one can take just one of the three PDFs, CT14, MMHT14 or NNPDF3.0. In this way, one can also avoid the theoretical problems due to the differences in the treatment of heavy quarks, which is introduced by the averaging procedure, as illustrated in Sec. 3. Alternative PDF choices can then always be studied to some approximation with a reweighting method. In spite of the valiant effort in Ref. [8] to provide a uniform solution, the PDF choice for measurements of precision observables must be decided on a case-by-case basis for each particular process.

### 5.2.3. Bottom-quark hadro-production

Bottom-quark production in proton-proton collisions at the LHC is also dominated by the gluon-gluon fusion process. Therefore, the LHCb measurements of  $B$ -meson production in the forward region [238] with rapidities  $2.0 < y < 4.5$  at  $\sqrt{s} = 7$  TeV probe the gluon distributions simultaneously at small  $x$  up to  $x \sim 2 \times 10^{-5}$  and at large  $x \simeq 1$ . The small- $x$  region is not accessible with HERA DIS data, for example. The potential improvements of PDFs near the edges of the currently covered kinematical region, namely, at small  $x$  and low scales, was first illustrated in [239, 240] using differential LHCb data on hadro-production of  $c\bar{c}$  and  $b\bar{b}$  pairs.

In the present comparison in Tab. 15, the normalized cross sections,  $(d\sigma/dy)/(d\sigma/dy_0)$ , for bottom-quark production are calculated from the absolute measurements published by LHCb, with  $d\sigma/dy_0$  being the cross section in the center bin,  $3 < y_0 < 3.5$ , of the measured rapidity range in each  $p_T$  bin [239]. In the absence of NNLO QCD corrections, the theoretical predictions are obtained at NLO in QCD [241–243] using a fixed number of flavors,  $n_f = 3$ , for the hard scattering cross sections. Since data for the hadro-production of heavy quarks other than top have not been

PDF sets	$\chi^2/\text{NDP}$ (with unc.)	$\chi^2/\text{NDP}$ (nominal)
ABM11 [48] <sup>a</sup>	222/244	394/244
CJ15 [1]	244/244	272/244
CT14 [3]	166/244	241/244
HERAPDF2.0 [4]	219/244	366/244
JR14 [5]	205/244	217/244
MMHT14 [6]	165/244	202/244
NNPDF3.0 [7]	160/244	197/244
PDF4LHC15 [8]	173/244	218/244

<sup>a</sup> The set ABM11 fit [48] is used here, because ABM12 [2] sets are only available at NNLO.

TABLE 15: The values of  $\chi^2/\text{NDP}$  for the normalised bottom-quark cross sections measured at LHCb [238] using the NLO PDFs of the individual groups. The left column accounts for the quoted PDF uncertainties (with the CJ15 and CT14 PDF uncertainties rescaled to 68% c.l.), while the right column uses the central prediction of each PDF set.

considered for publicly available PDF fits thus far, issues of any model dependence such as in [141] due to the use of GM-VFNS cannot be quantified. In the calculation of the normalized cross sections, the theoretical uncertainty is strongly reduced, since variations of the renormalization and factorization scales as well as of the fragmentation parameters do not significantly affect the shape of the  $y$  distributions for heavy-flavor production, while this shape remains sensitive to PDFs.

The values for  $\chi^2/\text{NDP}$  given in Tab. 15 are computed with the QCD fit platform `xFitter` for the individual PDF sets obtained at NLO, namely, ABM11 [48], CJ15 [1], CT14 [3], HERAPDF2.0 [4], JR14 [5], MMHT14 [6], NNPDF3.0 [7], as well as the averaged set PDF4LHC15 [8]. All PDFs provide a good description of the data, despite the fact that none of the groups use any data sensitive to the gluons at very low  $x$ , in the region directly probed by the LHCb  $B$ -meson measurement. Remarkably, one finds that  $\chi^2/\text{NDP} < 1$  for the vast majority of the groups (left column in Tab. 15), suggesting that the derived PDF uncertainties at the edges of the so far measured regions might be inflated.

#### 5.2.4. Charm-quark hadro-production

Charm-quark hadro-production offers another possibility to illustrate the consistency of the theory predictions for the various PDF sets. The exclusive production of charmed mesons in the forward region at LHCb probe the gluon distribution down to small- $x$  values of  $x \sim 5 \times 10^{-6}$  at  $\sqrt{s} = 7$  TeV, and data can be confronted with QCD predictions at NLO accuracy, see, e.g., [244, 245].

For the inclusive cross section of the reaction  $pp \rightarrow c\bar{c}$  the QCD predictions are known up to NNLO in the  $\overline{\text{MS}}$  scheme for the charm-quark mass and display good convergence of the perturbative expansion and stability under variation of the renormalization and factorization scales [244].




FIG. 16: Theoretical predictions for the total  $pp \rightarrow c\bar{c}$  cross section as a function of the center-of-mass energy  $\sqrt{s}$  at NLO (dashed lines) and NNLO (solid lines) QCD accuracy in the  $\overline{\text{MS}}$  mass scheme with  $m_c(m_c) = 1.27$  GeV and scale choice  $\mu_R = \mu_F = 2m_c(m_c)$  using the central PDF sets (solid lines) of ABM12 [2], CJ15 [1], CT14 [3] and JR14 [5] and the respective PDF uncertainties (dashed lines). The predictions for ABM12 (CJ15) use the NNLO (NLO) PDFs independent of the order of perturbation theory. See text for details and references on the experimental data from fixed target experiments and colliders (STAR, PHENIX, ALICE, ATLAS, LHCb).

In Figs. 16 and 17 we compare the theory predictions at NLO and NNLO with  $m_c(m_c) = 1.27$  GeV in the  $\overline{\text{MS}}$  scheme, see Sec. 3, for the scale choice  $\mu_r = \mu_f = 2m_c(m_c)$  as a function of the center-of-mass energy  $\sqrt{s}$  to available experimental data. These data span a large range in  $\sqrt{s}$ , which starts with fixed target experiments at energies up to  $\sqrt{s} = 50$  GeV summarized in [246] and HERA-B data [247] (purple points in Figs. 16, 17). At higher energies RHIC data from PHENIX and STAR [248, 249] (black points in Figs. 16, 17) are available and the LHC contributes measurements at energies  $\sqrt{s} = 2.76$  TeV from ALICE [250], at  $\sqrt{s} = 7$  TeV from ALICE [250], ATLAS [251] and LHCb [252], and at the highest available energy  $\sqrt{s} = 13$  TeV from LHCb [253] (blue points in Figs. 16, 17). The total cross sections of LHCb have been obtained from charmed hadron production measurements in a limited phase space region [252, 253] using extrapolations based on NLO QCD predictions matched with parton shower Monte Carlo generators.

The theory predictions for the PDF sets ABM12, CJ15, CT14 and JR14 at NLO and NNLO




FIG. 17: Same as Fig. 16 using the central PDF sets of HERAPDF2.0 [4], MMHT14 [6], NNPDF3.0 [7] and PDF4LHC15 [8] together with the respective PDF uncertainties.

are shown in Fig. 16, together with the respective PDF uncertainties. For all these PDF sets the perturbative expansion is stable, the theory computations agree well with the data and predictions, e.g., for a future collider with  $\sqrt{s} \simeq 100$  TeV, yield positive cross sections. The PDF uncertainties obtained for CT14, however, do increase significantly above energies of  $\sqrt{s} \simeq 1$  TeV.

The same information for the sets HERAPDF2.0, MMHT14, NNPDF3.0 and PDF4LHC15 is displayed in Fig. 17. These predictions all agree with data at low energies but start to behave very differently for HERAPDF2.0, MMHT14 or NNPDF3.0 at energies above  $\sqrt{s} \simeq \mathcal{O}(10)$  TeV and for PDF4LHC15 above  $\sqrt{s} \simeq \mathcal{O}(100)$  TeV. At the same time, the associated PDF uncertainties in this region of phase space become very large, thereby limiting the predictive power. Typically, the PDF uncertainties of the NNLO sets are even larger than at NLO. In the case of MMHT14 the consistency of the NNLO predictions with LHC data from ALICE [250], ATLAS [251] and LHCb [252, 253] at energies of  $\sqrt{s} = 7$  TeV and 13 TeV deteriorates. For NNPDF3.0 the central prediction at NNLO displays a change in slope for energies above  $\sqrt{s} \simeq 3$  TeV leading to a steeply rising cross section. The most striking feature, however, are the negative cross sections for HERAPDF2.0, MMHT14 and PDF4LHC15 at energies above  $\sqrt{s} \simeq \mathcal{O}(30 - 100)$  TeV, depending on the chosen set. This is an effect of the negative gluon PDF for those sets at values of  $x$  within the kinematic reach of current or future hadron colliders up to  $\sqrt{s} \simeq 100$  TeV. This results in

an instability of the perturbative expansion of the  $\sigma_{pp \rightarrow c\bar{c}}$  cross section at high energies when the contribution from the quark-gluon channel dominates. While the reason for a negative gluon PDF in the NNLO set of PDF4LHC15 (being some average of the CT14, MMHT14 and NNPDF3.0 sets) is unclear, we note that both HERAPDF2.0 and MMHT14 do apply the so-called RT optimal GM-VFNS [146] for the description of DIS charm-quark production at low scales and in the  $x$  range relevant for the case at hand, see Sec. 3 and Tab. 3. In contrast, other PDFs shown in Fig. 16 demonstrate stability of the perturbative expansion through NNLO up to very high energies and good consistency of the predictions with the experimental data.

### 5.3. $W'/Z'$ production

Cross sections sensitive to large- $x$  parton distributions typically fall rapidly with increasing  $x$  values, leading to limitations in the quantity and precision of experimental data and the kinematic range over which they can be obtained. Consequently, the precision to which one can constrain large- $x$  PDFs decreases with  $x$ , and systematic uncertainties due to extrapolations into unmeasured regions of  $x$  (or those excluded by cuts) increase. Similarly, the theoretical uncertainties due to various approximations in the treatment of nuclear corrections for deuterium data, or target mass and higher twist effects, also become larger.

To illustrate this, consider the production of a heavy  $W'$  boson as a function of the  $W'$  rapidity  $y_{W'}$  [254]. Assuming Standard Model couplings, the parton luminosity for a produced negatively charged  $W'^-$  boson is given by

$$\mathcal{L}_{W'^-} = \frac{2\pi G_F}{3\sqrt{2}} x_1 x_2 \left[ \cos^2 \theta_C (\bar{u}(x_2)d(x_1) + \bar{c}(x_2)s(x_1)) + \sin^2 \theta_C (\bar{u}(x_2)s(x_1) + \bar{c}(x_2)d(x_1)) \right] + (x_1 \leftrightarrow x_2), \quad (22)$$

where  $G_F$  is the Fermi constant and  $\theta_C$  the Cabibbo angle. The uncertainty  $\delta\mathcal{L}_{W'^-}$  in the luminosity is shown in Fig. 18 for various PDF sets as a function of  $y_{W'}$ , for several fixed values of the boson mass from the SM  $W$  up to  $M_{W'} = 7$  TeV.

Note that as the rapidity or mass of the produced boson increases, so does the momentum fraction  $x_{1,2} = (M_{W'}/\sqrt{s})e^{\pm y_{W'}}$  of one or both partons, in which case the luminosity behaves as  $\mathcal{L}^- \sim \bar{u}(x_2)d(x_1)$ . Except for the highest  $M_{W'}$  values, the PDF uncertainty typically remains small up to large values of  $y_{W'}$ , corresponding to  $x_1 \approx 0.65$ , beyond which it rises dramatically for all  $M_{W'}$ . This is precisely the region where data constraining the  $d$ -quark PDF are scarce, and theoretical assumptions play an important role [1]. This is particularly pronounced for fits that exclude DIS data at low invariant masses, such as the three fits included in the PDF4LHC combination [8]. For large  $W'$  masses, the  $\bar{u}$  PDF is evaluated at  $x_2 \sim 0.2 - 0.5$ , where data are either nonexistent or have large errors, giving rise to the increased uncertainties in some of the PDF sets at  $y_{W'} \sim 0$ .

Interestingly, at large  $y_{W'}$  the relative uncertainties from the CJ15, MMHT14, CT14 and PDF4LHC15 PDF sets displayed in Fig. 18 almost coincide, even though the quoted confidence levels for each vary. For instance, the MMHT14 and PDF4LHC15 sets are for the 68% c.l., while CT14 corresponds to 90% c.l., and CJ15 assumes a  $\Delta\chi^2 = 100$  (or tolerance  $T = 10$ ). The fact that similar uncertainties are found in several modern PDF analyses, that generally give good descriptions of the data but use very different methodologies for computing PDF errors and quote different confidence levels, illustrates the difficulty in obtaining statistically meaningful uncertainties from combinations of different PDF sets.




FIG. 18: Relative uncertainty  $\delta\mathcal{L}_{W'} / \mathcal{L}_{W'}$  in the  $W'$  luminosity as a function of rapidity  $y_{W'}$  for the combined PDF4LHC15 (68% c.l., dotted), CJ15 ( $\Delta\chi^2 = 100$ , solid), MMHT14 (68% c.l., dot-dashed), and CT14 (90% c.l., dashed), for various  $W'$  masses from 80 GeV (SM) to 7.0 TeV.




FIG. 19: Ratio of central values of the  $W'$  luminosity  $\mathcal{L}_{W'}$  to the PDF4LHC value (dotted, 68% c.l. shaded band) as a function of rapidity  $y_{W'}$ . The PDF sets CJ15 (red solid curve), MMHT14 (blue dot-dashed curve), CT14 (blue dashed curve), and NNPDF3.0 (green dashed curve) are compared for a  $W'$  mass  $M_{W'} = 3.5$  TeV.

This is also illustrated in Fig. 19, where the central values for the  $W'$  luminosity for several PDF sets are compared relative to the luminosity computed from the central PDF4LHC15 distributions. The combination PDF4LHC15 set is unable to adequately represent the systematic large- $x$  uncertainties due to the different theoretical assumptions utilized in the fits; indeed, the ratios of central values are seen to be of the same order as the overall PDF4LHC15 68% c.l. uncertainty, and in the case of the NNPDF3.0 set are about twice as large.

A comparison with the luminosity computed using the CJ15 PDF set, which is not included in



the PDF4LHC15 combination, is instructive here. The two main theoretical assumptions affecting the  $W'^-$  luminosity are the nuclear corrections in deuterium (applied or fitted in the CJ15 and MMHT14 analyses, as well as in JR14 and ABM12), and the parametrization of the  $d$ -quark PDF. For the latter, the choices made in PDF analyses are to either allow the  $d/u$  ratio to reach a finite, nonzero limiting value at  $x \rightarrow 1$  (CJ15 and CT14), or allow it to become very small,  $d/u \rightarrow 0$ , or very large,  $d/u \rightarrow \infty$  (MMHT14 and NNPDF3.0). Furthermore, the CJ15 distributions were also fitted to low invariant mass ( $W^2 < 12.5 \text{ GeV}^2$ ) DIS data, which were excluded by kinematic cuts by the MMHT14, CT14 and NNPDF3.0 groups. Consequently, the following features can be observed in Fig. 19:

- The MMHT14 curve follows CJ15 closely until  $y_{W'} \approx 1$  ( $x \approx 0.65$ ), after which the chosen  $d$ -quark parametrization forces an upward turn in the region not constrained by the large- $x$  and low- $W^2$  SLAC data utilized in CJ15.
- The CT14 curve is lower than CJ15 at  $y_{W'} \lesssim 0.6$  ( $x \lesssim 0.45$ ), and higher at larger  $y_{W'}$ , because of the neglect of nuclear corrections. At  $y_{W'} > 1$  the  $d$ -quark PDF is essentially unconstrained since neither the low- $W^2$  SLAC data nor the reconstructed Tevatron  $W$ -boson production data are included in the fit.
- The NNPDF3.0 fit, which excludes low- $W^2$  DIS data and does not utilize nuclear or hadronic corrections, consistently deviates from all others. It is, however, compatible with those within its own uncertainties, which at large  $x$  are about four times larger than that of the other fits.

In summary, in extreme kinematic regions, such as at large rapidity or for large-mass observables, caution must be exercised when utilizing PDF error bands and nominal confidence levels provided by the various PDF groups for precision calculations and statistically meaningful comparisons to data. Calculations performed with a PDF combination set should always be cross-checked with as many individual PDF sets as possible, taking into account the amount and kind of data included in each fit, as well as the different theoretical inputs. The latter explore different physics issues and can vary considerably from one PDF set to another. When differences arise, further scrutiny of the PDF fit results themselves may be needed before drawing any definitive conclusions.

## 6. RECOMMENDATIONS FOR PDF USAGE

Recommendations for the usage of PDFs generally aim to provide some guidance in estimating the magnitude and the uncertainties of cross sections in a reliable but also efficient way. First attempts in this direction have been provided by the *PDF4LHC Working Group Interim Recommendations* [255], which based its recommendations for predictions at NNLO in QCD on the central set of the MSTW [173] PDFs, along with an elaborate procedure to compute PDF uncertainties based on an envelope with some other sets. However, this approach has been criticized for being impractical. The 2015 PDF4LHC recommendations [8] have evolved from these discussions and aim at improving the efficiency of cross section computations by averaging several PDFs along with their respective uncertainties. We briefly recall these suggestions and put them into context of the findings of the previous sections. We comment on several shortcomings of the recommendations proposed in [8] and present an alternative set of recommendations for PDF usage at the LHC that do not suffer from these problems.

## 6.1. The 2015 PDF4LHC recommendations: A critical appraisal

The 2015 PDF4LHC recommendations [8] distinguish in total four cases: (i) *Comparisons between data and theory for Standard Model measurements*, (ii) *Searches for Beyond the Standard Model phenomena*, (iii) *Calculation of PDF uncertainties in situations when computational speed is needed, or a more limited number of error PDFs may be desirable* and (iv) *Calculation of PDF uncertainties in precision observables*.

In case (i) the recommendation is to use the individual PDF sets ABM12 [2], CJ12 [169], CT14 [3], JR14 [5], HERAPDF2.0 [4], MMHT14 [6], and NNPDF3.0 [7] for comparisons between data and theory for SM measurements. This is obvious and follows similar advice given in [255]. It is not clear, however, that the full account of the PDF dependence should be limited to SM processes only. We find that differences observed in the theory predictions with the various PDFs can often be traced to using different underlying theoretical descriptions and models in the PDF fits. With the availability of more LHC data, tests of the compatibility of those data sets in the individual PDF fits will become more important, as well as studies to quantify the constraining potential of certain processes, e.g., hadro-production of  $t\bar{t}$  pairs, jets or  $W^\pm$  and  $Z$  bosons. Likewise, for PDF fits with ranges of different  $\alpha_s(M_Z)$  values it will be interesting to provide the profile for the value of  $\chi^2/\text{NDP}$ .

For case (ii), in searching for BSM phenomena it is recommended to employ the PDF4LHC15 sets [8]. Those sets are constructed as a statistical combination of the CT14 [3], MMHT14 [6], and NNPDF3.0 [7] PDF sets, using a Monte Carlo approach with various levels of precision leading to, e.g., the PDF4LHC15\_30 or PDF4LHC15\_100 sets, which differ in the number of error PDFs. The restriction to CT14, MMHT14 and NNPDF3.0 implies a bias both for the central value and for the PDF uncertainties of BSM cross section predictions. The bias is best illustrated by recalling one of the selection requirements [8], namely that the central value of  $\alpha_s(M_Z)$  is fixed at an agreed common value, currently chosen to be  $\alpha_s(M_Z) = 0.118$  at both NLO and NNLO. Such choices are in clear contradiction with the precision determinations of  $\alpha_s(M_Z)$  at different orders in perturbation theory, as summarized in Sec. 4. Moreover, for searches at the highest energies, the PDFs are probed close to the hadronic threshold near  $x \simeq 1$ , where nuclear corrections and other hadronic effects, considered for instance in the CJ15 [1] and JR14 [5] analyses, are important.

Case (iii) advises the use of the PDF4LHC15\_30 sets in the calculation of PDF uncertainties when computational speed is needed, or a more limited number of error PDFs may be desirable. This is certainly more efficient for Monte Carlo simulations than the previous envelope procedure [255]. At the same time, this recommendation is particularly worrisome because of the large deviations between data and theory predictions at low scales and also at the edges of the kinematical ranges of data currently used in PDF fits, as the illustrations based on exemplary cross sections in Secs. 3 and 5 have shown. Also, the average of various GM-VFNS for heavy quark production, such as ACOT [142], FONLL [145] and RT [146], leaves a large degree of arbitrariness in the theory predictions, cf. Fig. 10. Note also that the PDF4LHC15\_30 sets were updated around the end of December 2015 [256] to account for an extension of their validity ranges down to small values of  $Q$ , as the originally published sets were only valid for  $Q > 8$  GeV. This restriction has neither been described in [8] nor have the changes been mentioned so far in the published literature.

Finally, case (iv) concerns the calculation of PDF uncertainties in precision observables and advises to use the set PDF4LHC15\_100. However, it is unclear why PDF uncertainties should be dealt with differently than in recommendation (i), i.e., by considering all individual PDF sets. As far as fiducial cross sections are concerned, where extrapolations with additional assumptions or techniques are not required, each individual set is supposed to provide faithful uncertainty bands

from the propagation of the errors of the experimental data used in the PDF fit. If extrapolations are required, such as near the boundary of the kinematics covered in PDF fits, additional care should be taken as the results of the PDF4LHC15 sets cannot be considered reliable, see, e.g. Figs. 17 and 19 for the PDF uncertainties provided by the PDF4LHC15\_100 set at NLO and NNLO.

In summary, the recent PDF4LHC recommendations [8] cannot be viewed as definitive in the case of precision theory predictions, as the averaging procedure recently advocated in the PDF4LHC recommendations [8] introduces bias, artificially inflates the uncertainties, and makes it very difficult to quantify potential discrepancies between individual sets.

## 6.2. New recommendations for PDF usage at the LHC

With the aim of simplifying the usage of PDFs while maintaining predictive capability of the individual sets, here we present a new, streamlined set of recommendations for PDF usage at the LHC. We distinguish two cases: (i) *Precision theory predictions*, a class of predictions, either within or beyond the SM, which encompasses any type of cross section prediction including radiative corrections of any kind, whether at fixed-order or via resummation to some logarithmic accuracy; (ii) *Other theory predictions*, the complementary class containing all other cross section predictions where high precision is not required, such as those based on Born approximations and/or magnitude estimates, or in cases where precision may be sacrificed in favor of computational speed.

For these two cases, the recommendations are:

### (i) Precision theory predictions

**Recommendation:** Use the individual PDF sets ABM12, CJ15, CT14, JR14, HERAPDF2.0, MMHT14 and NNPDF3.0 (or as many as possible), together with the respective uncertainties for the chosen PDF set, the strong coupling  $\alpha_s(M_Z)$  and the heavy quark masses  $m_c$ ,  $m_b$  and  $m_t$ .

**Rationale:** Precise theory predictions as needed for any comparisons between theory and data for processes in the SM or beyond (such as hadro-production of jets,  $W^\pm$ - or Z-boson production, either singly or in pairs, heavy-quark hadro-production, or generally the production of new massive particles at the TeV scale) often depend on details of the PDF fits and the underlying theory assumptions and schemes used. Differences in the theory predictions based on the individual sets ABM12 [2], CJ15 [1], CT14 [3], JR14 [5], HERAPDF2.0 [4], MMHT14 [6], and NNPDF3.0 [7] can give an indication of residual systematic uncertainties or shed light on drawbacks and potential improvements of the physics models used in the extraction of those PDFs. Of course, this advice applies as well to measurements of SM parameters such as the strong coupling  $\alpha_s(M_Z)$ , heavy quark masses  $m_c$ ,  $m_b$  and  $m_t$  or the W-boson mass  $M_W$ , because these parameters are directly correlated to the PDFs used in their extraction from some observable at the LHC.

### (ii) Other theory predictions

**Recommendation:** Use any one of the PDF sets listed in LHAPDF(v6).

**Rationale:** Often in phenomenological applications for the LHC one is interested in quick, order of magnitude estimates for the cross sections. These are directly proportional to the parton luminosity, and often to the value of  $\alpha_s(M_Z)$ . Restricting the recommendation to

any of the parton densities sets provided by the LHAPDF (v6) [257] interface excludes parton luminosities with lesser precision in the interpolation of the underlying grids (e.g., in LHAPDF (v5) [258]) or “partonometers” [259] with outdated calibration. In other situations, such as in complex MC simulations of LHC processes, or in studies of new accelerators and detectors, one may be willing to sacrifice precision in favor of speed of calculation. In this and in the previous cases, usage of the sets PDF4LHC15\_30 and PDF4LHC15\_100 may provide an efficient way of simultaneously estimating PDF statistical and systematic uncertainties, although care must be exercised in their interpretation depending on the observable and covered kinematic range. Given an initial PDF choice, the impact of different PDF sets can then be efficiently studied with reweighting methods to provide additional control over PDF systematic uncertainties.

The new PDF recommendations provide an efficient alternative both for the use with Monte Carlo event generators in order to generate large MC samples for the LHC data analysis, as well as for precision theory predictions together with the assessment of the PDF uncertainties themselves.

The proposal also meets the developments of modern tools for NLO QCD computations matched with Shower Monte Carlo generators, which provide nowadays the workhorse for many experimental analyses at the LHC. The latest version of the most widely used tools, like MadGraph5\_aMC@NLO [260], POWHEG-BOX (v2) [261, 262] and SHERPA (v2) [263, 264], or of other recently developed generators, like Geneva [265], implement the reweighting feature. This allows to generate weighted events for a given setup, and then to reweight a-posteriori each event in a fast and efficient way, by generating new weights associated with different choices of renormalization and factorization scales and/or PDFs. Those events can then be used to perform experimental analyses with different cuts, extracting differential distributions at the hadron level. Actually, at present and to the best of our knowledge, PDF reweighting is performed by assuming that PDF dependence of weights is linear, whereas it is not, because PDFs are present even in the Sudakov form-factor. Efforts to extend the reweighting to the entire Sudakov form-factor and to the full shower are indeed ongoing. The reweighting technique turns out to be particularly useful to compute in a fast (although at the moment approximate) way PDF uncertainties affecting the predictions obtained by the tools mentioned above, especially when a large number of PDF sets are available at this purpose, allowing to avoid to repeat from the beginning CPU intensive calculations. Thus, the use of as many as possible of the individual PDF sets ABM12, CJ15, CT14, JR14, HERAPDF2.0, MMHT14 and NNPDF3.0 can be realized in a straightforward manner.

## 7. CONCLUSION

In this report we have reviewed recent developments in the determination of PDFs in global QCD analyses. Thanks to high precision experimental measurements and continuous theoretical improvements, the parton content of the proton is generally well constrained and PDFs, along with the strong coupling constant  $\alpha_s(M_Z)$  and the heavy-quark masses  $m_c$ ,  $m_b$  and  $m_t$ , have been determined with good accuracy, at least at NNLO in QCD. This forms the foundation for precise cross section predictions at the LHC in Run 2.

We have briefly discussed the available data used in PDF extractions and the kinematic range covered, and emphasized the importance of selecting mutually consistent sets of data in PDF fits in order to achieve acceptable  $\chi^2$  values for the goodness-of-fit estimate. The main thrust of the study has been the computation of benchmark cross sections for a variety of processes at hadron colliders, including Higgs boson production in gluon-gluon fusion. We have illustrated

how different choices for the theoretical description of the hard scattering process and choices of parameters have an impact on the predicted cross sections, and lead to systematic shifts that are often significantly larger than the associated PDF and  $\alpha_s(M_Z)$  uncertainties. A particular example has been the treatment of heavy quarks in DIS, where the quality of the various scheme choices has been quantified in terms of  $\chi^2/\text{NPD}$  values when comparing predicted cross sections to data. We have also pointed out the inconsistently low values for the pole mass of the charm quark used in some fits, and have stressed the correlation of the strong coupling constant  $\alpha_s(M_Z)$  with the PDF parameters. Ideally,  $\alpha_s(M_Z)$  should be determined simultaneously with the PDFs, and we have summarized here the state of the art in the context of PDF analyses.

Our findings expose a number of shortcomings in the recent PDF4LHC recommendations [8]. We have shown that these do not provide sufficient control over some theoretical uncertainties, and may therefore be problematic for precision predictions in Run 2 of the LHC. Instead, we suggest new recommendations for the usage of PDFs to meet the precision requirements of the LHC era. At the same time, we seek to simplify the computational overhead for the user in the determination of uncertainties assigned to the parton luminosity and give the user a larger choice of possible PDF sets.

### Acknowledgments

We would like to thank S. Alioli, M. Botje and K. Rabbertz for discussions and K. Rabbertz for valuable comments on the manuscript. We are also deeply grateful to L. Harland-Lang and to R. Thorne for a check of the Higgs cross sections in Tab. 10.

This work has been supported by Bundesministerium für Bildung und Forschung through contract (05H2015), by the DOE contract No. DE-AC05-06OR23177, under which Jefferson Science Associates, LLC operates Jefferson Lab, and by the European Commission through PITN-GA-2012-316704 (*HIGGSTOOLS*). The work of A.A. and J.F.O. was supported in part by DOE contracts No. DE-SC0008791 and No. DE-FG02-97ER41922, respectively. Two of the authors (J.B. and S.M.) would like to thank the Mainz Institute for Theoretical Physics (MITP) for its hospitality and support.

- 
- [1] A. Accardi, L. T. Brady, W. Melnitchouk, J. F. Owens, and N. Sato, *Constraints on large- $x$  parton distributions from new weak boson production and deep-inelastic scattering data*, arXiv:1602.03154.
  - [2] S. Alekhin, J. Blümlein, and S. Moch, *The ABM parton distributions tuned to LHC data*, *Phys. Rev. D* **89** (2014) 054028, [arXiv:1310.3059].
  - [3] S. Dulat, T. J. Hou, J. Gao, M. Guzzi, J. Huston, P. Nadolsky, J. Pumplin, C. Schmidt, D. Stump, and C. P. Yuan, *The CT14 global analysis of Quantum Chromodynamics*, arXiv:1506.07443.
  - [4] **ZEUS, H1** Collaboration, H. Abramowicz et al., *Combination of measurements of inclusive deep-inelastic  $e^\pm p$  scattering cross sections and QCD analysis of HERA data*, arXiv:1506.06042.
  - [5] P. Jimenez-Delgado and E. Reya, *Delineating parton distributions and the strong coupling*, *Phys. Rev. D* **89** (2014) 074049, [arXiv:1403.1852].
  - [6] L. A. Harland-Lang, A. D. Martin, P. Motylinski, and R. S. Thorne, *Parton distributions in the LHC era: MMHT 2014 PDFs*, *Eur. Phys. J. C* **75** (2015) 204, [arXiv:1412.3989].

- [7] **NNPDF** Collaboration, R. D. Ball et al., *Parton distributions for the LHC Run II*, *JHEP* **04** (2015) 040, [arXiv:1410.8849].
- [8] J. Butterworth et al., *PDF4LHC recommendations for LHC Run II*, *J. Phys.* **G43** (2016) 023001, [arXiv:1510.03865].
- [9] L. A. Harland-Lang, A. D. Martin, P. Motylinski, and R. S. Thorne, *Charm and beauty quark masses in the MMHT2014 global PDF analysis*, arXiv:1510.02332.
- [10] H. Paukkunen and C. A. Salgado, *Agreement of neutrino deep-inelastic scattering data with global fits of parton distributions*, *Phys. Rev. Lett.* **110** (2013) 212301, [arXiv:1302.2001].
- [11] G. Altarelli, *QCD and experiment: Status of  $\alpha_s$* , in: P.M. Zerwas and H.A. Kastrup eds., *Workshop on QCD: 20 Years Later Aachen, Germany, June 9-13 (1992)* p. 172–204.
- [12] J. Blümlein, M. Klein, G. Ingelman, and R. Rückl, *Testing QCD scaling violations in the HERA energy range*, *Z. Phys.* **C45** (1990) 501.
- [13] J. P. Berge et al., *A measurement of differential cross sections and nucleon structure functions in charged current neutrino interactions on iron*, *Z. Phys.* **C49** (1991) 187.
- [14] A. Gehrmann-De Ridder, T. Gehrmann, E. W. N. Glover, and J. Pires, *Second order QCD corrections to jet production at hadron colliders: the all-gluon contribution*, *Phys. Rev. Lett.* **110** (2013) 162003, [arXiv:1301.7310].
- [15] **H1** Collaboration, F. D. Aaron et al., *Inclusive deep-inelastic scattering at high  $Q^2$  with longitudinally polarised lepton beams at HERA*, *JHEP* **09** (2012) 061, [arXiv:1206.7007].
- [16] **ATLAS** Collaboration, G. Aad et al., *Measurement of the inclusive  $W^\pm$  and  $Z/\gamma$  cross sections in the electron and muon decay channels in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector*, *Phys. Rev.* **D85** (2012) 072004, [arXiv:1109.5141].
- [17] **CMS** Collaboration, S. Chatrchyan et al., *Measurement of the muon charge asymmetry in inclusive  $pp \rightarrow W + X$  production at  $\sqrt{s} = 7$  TeV and an improved determination of light parton distribution functions*, *Phys. Rev.* **D90** (2014) 032004, [arXiv:1312.6283].
- [18] **CMS** Collaboration, V. Khachatryan et al., *Measurement of the differential cross section and charge asymmetry for inclusive  $pp \rightarrow W + X$  production at  $\sqrt{s} = 8$  TeV*, arXiv:1603.01803.
- [19] **DØ** Collaboration, V. M. Abazov et al., *Measurement of the muon charge asymmetry in  $p\bar{p} \rightarrow W + X \rightarrow \mu\nu + X$  events at  $\sqrt{s}=1.96$  TeV*, *Phys.Rev.* **D88** (2013) 091102, [arXiv:1309.2591].
- [20] **DØ** Collaboration, V. M. Abazov et al., *Measurement of the electron charge asymmetry in  $p\bar{p} \rightarrow W + X \rightarrow e\nu + X$  decays in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV*, *Phys. Rev.* **D91** (2015) 032007, [arXiv:1412.2862]. [Erratum: *Phys. Rev.* **D91**,no.7,079901(2015)].
- [21] **LHCb** Collaboration, R. Aaij et al., *Measurement of the forward  $Z$  boson production cross section in  $pp$  collisions at  $\sqrt{s} = 7$  TeV*, *JHEP* **08** (2015) 039, [arXiv:1505.07024].
- [22] **LHCb** Collaboration, R. Aaij et al., *Measurement of forward  $Z \rightarrow e^+e^-$  production at  $\sqrt{s} = 8$  TeV*, *JHEP* **05** (2015) 109, [arXiv:1503.00963].
- [23] **LHCb** Collaboration, R. Aaij et al., *Measurement of forward  $W$  and  $Z$  boson production in  $pp$  collisions at  $\sqrt{s} = 8$  TeV*, *JHEP* **01** (2016) 155, [arXiv:1511.08039].
- [24] S. Alekhin, J. Blümlein, S. Moch, and R. Placakyte, *Iso-spin asymmetry of quark distributions and implications for single top-quark production at the LHC*, arXiv:1508.07923.
- [25] **CDF** Collaboration, D. Acosta et al., *Measurement of the forward-backward charge asymmetry from  $W \rightarrow e\nu$  production in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV*, *Phys. Rev.* **D71** (2005) 051104, [hep-ex/0501023].
- [26] **CDF** Collaboration, T. Aaltonen et al., *Direct measurement of the  $W$  production charge asymmetry in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV*, *Phys. Rev. Lett.* **102** (2009) 181801, [arXiv:0901.2169].
- [27] **DØ** Collaboration, V. M. Abazov et al., *Measurement of the  $W$  boson production charge asymmetry*

- in  $p\bar{p} \rightarrow W + X \rightarrow e\nu + X$  events at  $\sqrt{s} = 1.96$  TeV, *Phys. Rev. Lett.* **112** (2014) 151803, [arXiv:1312.2895]. [Erratum: *Phys. Rev. Lett.* 114,no.4,049901(2015)].
- [28] **HERAFitter developers' Team** Collaboration, S. Camarda et al., *QCD analysis of W- and Z-boson production at the Tevatron*, arXiv:1503.05221.
- [29] A. Kwiatkowski, H. Spiesberger, and H. J. Möhring, *Heracles: An event generator for ep interactions at HERA energies including radiative processes: version 1.0*, *Comput. Phys. Commun.* **69** (1992) 155.
- [30] A. Arbuzov, D. Yu. Bardin, J. Blümlein, L. Kalinovskaya, and T. Riemann, *Hector 1.00: A program for the calculation of QED, QCD and electroweak corrections to ep and  $l^{\pm}N$  deep-inelastic neutral and charged current scattering*, *Comput. Phys. Commun.* **94** (1996) 128, [hep-ph/9511434].
- [31] J. Blümlein, *Leading log radiative corrections to ep scattering including jet measurement*, *Phys. Lett.* **B271** (1991) 267.
- [32] J. Blümlein,  *$O(\alpha^2 L^2)$  radiative corrections to deep-inelastic ep scattering for different kinematical variables*, *Z. Phys.* **C65** (1995) 293, [hep-ph/9403342].
- [33] J. Blümlein and H. Kawamura,  *$O(\alpha^2 L)$  radiative corrections to deep-inelastic ep scattering*, *Phys. Lett.* **B553** (2003) 242, [hep-ph/0211191].
- [34] J. Kripfganz and H. Perl, *Electroweak radiative corrections and quark mass singularities*, *Z. Phys.* **C41** (1988) 319.
- [35] J. Blümlein, *Leading log radiative corrections to deep-inelastic neutral and charged current scattering at HERA*, *Z. Phys.* **C47** (1990) 89.
- [36] H. Spiesberger, *QED radiative corrections for parton distributions*, *Phys. Rev.* **D52** (1995) 4936, [hep-ph/9412286].
- [37] M. Roth and S. Weinzierl, *QED corrections to the evolution of parton distributions*, *Phys. Lett.* **B590** (2004) 190, [hep-ph/0403200].
- [38] J. Blümlein, *The theory of deeply inelastic scattering*, *Prog. Part. Nucl. Phys.* **69** (2013) 28, [arXiv:1208.6087].
- [39] **Particle Data Group** Collaboration, K. A. Olive et al., *Review of particle physics*, *Chin. Phys.* **C38** (2014) 090001.
- [40] S. Moch, J. A. M. Vermaseren, and A. Vogt, *The longitudinal structure function at the third order*, *Phys. Lett.* **B606** (2005) 123, [hep-ph/0411112].
- [41] J. A. M. Vermaseren, A. Vogt, and S. Moch, *The third-order QCD corrections to deep-inelastic scattering by photon exchange*, *Nucl. Phys.* **B724** (2005) 3, [hep-ph/0504242].
- [42] J. Blümlein, H. Böttcher, and A. Guffanti, *Non-singlet QCD analysis of deep-inelastic world data at  $O(\alpha_s^3)$* , *Nucl. Phys.* **B774** (2007) 182, [hep-ph/0607200].
- [43] S. Alekhin, J. Blümlein, and S. Moch, *Higher order constraints on the Higgs production rate from fixed-target DIS data*, *Eur. Phys. J.* **C71** (2011) 1723, [arXiv:1101.5261].
- [44] H. Georgi and H. D. Politzer, *Freedom at moderate energies: masses in color dynamics*, *Phys. Rev.* **D14** (1976) 1829.
- [45] F. M. Steffens, M. D. Brown, W. Melnitchouk, and S. Sanches, *Parton distributions in the presence of target mass corrections*, *Phys. Rev.* **C86** (2012) 065208, [arXiv:1210.4398].
- [46] J. Blümlein and H. Böttcher, *Higher twist contributions to the structure functions  $F_2^p(x, Q^2)$  and  $F_2^d(x, Q^2)$  at large x and higher orders*, *Phys. Lett.* **B662** (2008) 336, [arXiv:0802.0408].
- [47] S. I. Alekhin, *Global fit to the charged leptons DIS data:  $\alpha_s$  parton distributions, and high twists*, *Phys. Rev.* **D63** (2001) 094022, [hep-ph/0011002].
- [48] S. Alekhin, J. Blümlein, and S. Moch, *Parton distribution functions and benchmark cross sections at NNLO*, *Phys. Rev.* **D86** (2012) 054009, [arXiv:1202.2281].

- [49] L. A. Harland-Lang, A. D. Martin, P. Motylinski, and R. S. Thorne, *The impact of the final HERA combined data on PDFs obtained from a global fit*, arXiv:1601.03413.
- [50] J. Gao, M. Guzzi, J. Huston, H.-L. Lai, Z. Li, P. Nadolsky, J. Pumplin, D. Stump, and C. P. Yuan, *CT10 next-to-next-to-leading order global analysis of QCD*, *Phys. Rev.* **D89** (2014) 033009, [arXiv:1302.6246].
- [51] A. Chuvakin, J. Smith, and W. L. van Neerven, *Comparison between variable flavor number schemes for charm quark electroproduction*, *Phys. Rev.* **D61** (2000) 096004, [hep-ph/9910250].
- [52] K. G. Wilson, *Nonlagrangian models of current algebra*, *Phys. Rev.* **179** (1969) 1499.
- [53] W. Zimmermann, *Lectures on elementary particle physics and quantum field theory, Brandeis Summer Institute, Vol. 1 (MIT Press, Cambridge, 1970)*, p. 395 (1970).
- [54] R. A. Brandt and G. Preparata, *Operator product expansions near the light cone*, *Nucl. Phys.* **B27** (1972) 541.
- [55] Y. Frishman, *Operator products at almost light like distances*, *Annals Phys.* **66** (1971) 373.
- [56] W. Furmanski and R. Petronzio, *Lepton - hadron processes beyond leading order in Quantum Chromodynamics*, *Z. Phys.* **C11** (1982) 293.
- [57] E. B. Zijlstra and W. L. van Neerven, *Contribution of the second order gluonic Wilson coefficient to the deep-inelastic structure function*, *Phys. Lett.* **B273** (1991) 476.
- [58] W. L. van Neerven and E. B. Zijlstra, *Order  $\alpha_s^2$  contributions to the deep-inelastic Wilson coefficient*, *Phys. Lett.* **B272** (1991) 127.
- [59] E. B. Zijlstra and W. L. van Neerven, *Order  $\alpha_s^2$  QCD corrections to the deep-inelastic proton structure functions  $F_2$  and  $F_L$* , *Nucl. Phys.* **B383** (1992) 525.
- [60] D. I. Kazakov and A. V. Kotikov, *Total  $\alpha_s$  correction to deep-inelastic scattering cross section ratio,  $R = \sigma_L/\sigma_T$  in QCD. Calculation of longitudinal structure function*, *Nucl. Phys.* **B307** (1988) 721. [Erratum: *Nucl. Phys.* B345,299(1990)].
- [61] D. I. Kazakov, A. V. Kotikov, G. Parente, O. A. Sampayo, and J. Sanchez Guillen, *Complete quartic ( $\alpha_s^4$ ) correction to the deep-inelastic longitudinal structure function  $F_L$  in QCD*, *Phys. Rev. Lett.* **65** (1990) 1535. [Erratum: *Phys. Rev. Lett.* 65 (1990) 2921].
- [62] J. Sanchez Guillen, J. Miramontes, M. Miramontes, G. Parente, and O. A. Sampayo, *Next-to-leading order analysis of the deep-inelastic  $R = \sigma_L/\sigma_{\text{total}}$* , *Nucl. Phys.* **B353** (1991) 337.
- [63] E. B. Zijlstra and W. L. van Neerven, *Order  $\alpha_s^2$  correction to the structure function  $F_3(x, Q^2)$  in deep-inelastic neutrino-hadron scattering*, *Phys. Lett.* **B297** (1992) 377.
- [64] S. Moch and J. A. M. Vermaseren, *Deep-inelastic structure functions at two loops*, *Nucl. Phys.* **B573** (2000) 853, [hep-ph/9912355].
- [65] S. Moch and M. Rogal, *Charged current deep-inelastic scattering at three loops*, *Nucl. Phys.* **B782** (2007) 51, [arXiv:0704.1740].
- [66] S. Moch, M. Rogal, and A. Vogt, *Differences between charged-current coefficient functions*, *Nucl. Phys.* **B790** (2008) 317, [arXiv:0708.3731].
- [67] S. Moch, J. A. M. Vermaseren, and A. Vogt, *Third-order QCD corrections to the charged-current structure function  $F_3$* , *Nucl. Phys.* **B813** (2009) 220, [arXiv:0812.4168].
- [68] D. J. Gross and F. Wilczek, *Asymptotically free gauge theories. 1*, *Phys. Rev.* **D8** (1973) 3633.
- [69] D. J. Gross and F. Wilczek, *Asymptotically free gauge theories. 2.*, *Phys. Rev.* **D9** (1974) 980.
- [70] H. Georgi and H. D. Politzer, *Electroproduction scaling in an asymptotically free theory of strong interactions*, *Phys. Rev.* **D9** (1974) 416.
- [71] G. Parisi, *An introduction to scaling violations*, in *Weak Interactions and Neutrino Physics. Proceedings: 11th Rencontre de Moriond, Flaine 1976, Feb 28-Mar 12, 1976. 2.*, pp. 83–114, 1976.
- [72] K. J. Kim and K. Schilcher, *Scaling violation in the infinite momentum frame*, *Phys. Rev.* **D17**



- (1978) 2800.
- [73] G. Altarelli and G. Parisi, *Asymptotic freedom in parton language*, *Nucl. Phys.* **B126** (1977) 298.
  - [74] E. G. Floratos, D. A. Ross, and C. T. Sachrajda, *Higher order effects in asymptotically free gauge theories: The anomalous dimensions of Wilson operators*, *Nucl. Phys.* **B129** (1977) 66. [Erratum: *Nucl. Phys.*B139,545(1978)].
  - [75] E. G. Floratos, D. A. Ross, and C. T. Sachrajda, *Higher order effects in asymptotically free gauge theories. 2. Flavor singlet Wilson operators and coefficient functions*, *Nucl. Phys.* **B152** (1979) 493.
  - [76] A. Gonzalez-Arroyo, C. Lopez, and F. J. Yndurain, *Second order contributions to the structure functions in deep-inelastic scattering. 1. Theoretical calculations*, *Nucl. Phys.* **B153** (1979) 161.
  - [77] A. Gonzalez-Arroyo, C. Lopez, and F. J. Yndurain, *Second order contributions to the structure functions in deep-inelastic scattering. 2. Comparison with experiment for the nonsinglet contributions to  $e, \mu$  nucleon scattering*, *Nucl. Phys.* **B159** (1979) 512.
  - [78] G. Curci, W. Furmanski, and R. Petronzio, *Evolution of parton densities beyond leading order: The nonsinglet case*, *Nucl. Phys.* **B175** (1980) 27.
  - [79] W. Furmanski and R. Petronzio, *Singlet parton densities beyond leading order*, *Phys. Lett.* **B97** (1980) 437.
  - [80] E. G. Floratos, R. Lacaze, and C. Kounnas, *Space and timelike cut vertices in QCD beyond the leading order. 1. Nonsinglet sector*, *Phys. Lett.* **B98** (1981) 89.
  - [81] E. G. Floratos, C. Kounnas, and R. Lacaze, *Higher order QCD effects in inclusive annihilation and deep-inelastic scattering*, *Nucl. Phys.* **B192** (1981) 417.
  - [82] A. Gonzalez-Arroyo and C. Lopez, *Second order contributions to the structure functions in deep-inelastic scattering. 3. The singlet case*, *Nucl. Phys.* **B166** (1980) 429.
  - [83] E. G. Floratos, R. Lacaze, and C. Kounnas, *Space and timelike cut vertices in QCD beyond the leading order. 2. The singlet sector*, *Phys. Lett.* **B98** (1981) 285.
  - [84] R. Hamberg and W. L. van Neerven, *The correct renormalization of the gluon operator in a covariant gauge*, *Nucl. Phys.* **B379** (1992) 143.
  - [85] R. K. Ellis and W. Vogelsang, *The evolution of parton distributions beyond leading order: The singlet case*, [hep-ph/9602356](https://arxiv.org/abs/hep-ph/9602356).
  - [86] S. Moch, J. A. M. Vermaseren, and A. Vogt, *The three-loop splitting functions in QCD: The nonsinglet case*, *Nucl. Phys.* **B688** (2004) 101–134, [[hep-ph/0403192](https://arxiv.org/abs/hep-ph/0403192)].
  - [87] A. Vogt, S. Moch, and J. A. M. Vermaseren, *The three-loop splitting functions in QCD: The singlet case*, *Nucl. Phys.* **B691** (2004) 129–181, [[hep-ph/0404111](https://arxiv.org/abs/hep-ph/0404111)].
  - [88] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel, and C. Schneider, *The 3-loop pure singlet heavy flavor contributions to the structure function  $F_2(x, Q^2)$  and the anomalous dimension*, *Nucl. Phys.* **B890** (2014) 48, [[arXiv:1409.1135](https://arxiv.org/abs/1409.1135)].
  - [89] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog, A. Lazopoulos, and B. Mistlberger, *High precision determination of the gluon fusion Higgs boson cross section at the LHC*, [arXiv:1602.00695](https://arxiv.org/abs/1602.00695).
  - [90] I. B. Khriplovich, *Green's functions in theories with non-abelian gauge group*, *Sov. J. Nucl. Phys.* **10** (1969) 235. [*Yad. Fiz.*10,409(1969)].
  - [91] G. 't Hooft, *unpublished*, .
  - [92] H. D. Politzer, *Reliable perturbative results for strong interactions?*, *Phys. Rev. Lett.* **30** (1973) 1346.
  - [93] D. J. Gross and F. Wilczek, *Ultraviolet behavior of nonabelian gauge theories*, *Phys. Rev. Lett.* **30** (1973) 1343.
  - [94] W. E. Caswell, *Asymptotic behavior of nonabelian gauge theories to two loop order*, *Phys. Rev.*

- Lett.* **33** (1974) 244.
- [95] D. R. T. Jones, *Two loop diagrams in Yang-Mills theory*, *Nucl. Phys.* **B75** (1974) 531.
- [96] O. V. Tarasov, A. A. Vladimirov, and A. Yu. Zharkov, *The Gell-Mann-Low function of QCD in the three-loop approximation*, *Phys. Lett.* **B93** (1980) 429.
- [97] S. A. Larin and J. A. M. Vermaseren, *The three loop QCD beta function and anomalous dimensions*, *Phys. Lett.* **B303** (1993) 334, [hep-ph/9302208].
- [98] T. van Ritbergen, J. A. M. Vermaseren, and S. A. Larin, *The four loop beta function in Quantum Chromodynamics*, *Phys. Lett.* **B400** (1997) 379, [hep-ph/9701390].
- [99] M. Czakon, *The four-loop QCD beta-function and anomalous dimensions*, *Nucl. Phys.* **B710** (2005) 485, [hep-ph/0411261].
- [100] M. Diemoz, F. Ferroni, E. Longo, and G. Martinelli, *Parton densities from deep-inelastic scattering to hadronic processes at super collider energies*, *Z. Phys.* **C39** (1988) 21.
- [101] M. Glück, E. Reya, and A. Vogt, *Parton structure of the photon beyond the leading order*, *Phys. Rev.* **D45** (1992) 3986.
- [102] R. K. Ellis, Z. Kunszt, and E. M. Levin, *The evolution of parton distributions at small  $x$* , *Nucl. Phys.* **B420** (1994) 517. [Erratum: *Nucl. Phys.*B433,498(1995)].
- [103] J. Blümlein and A. Vogt, *The evolution of unpolarized singlet structure functions at small  $x$* , *Phys. Rev.* **D58** (1998) 014020, [hep-ph/9712546].
- [104] J. Blümlein, S. Riemersma, W. L. van Neerven, and A. Vogt, *Theoretical uncertainties in the QCD evolution of structure functions and their impact on  $\alpha_s(M_Z^2)$* , *Nucl. Phys. Proc. Suppl.* **51C** (1996) 97, [hep-ph/9609217].
- [105] G. Rossi,  *$x$  space analysis for the photon structure functions in QCD*, *Phys. Rev.* **D29** (1984) 852.
- [106] A. Vogt, *Efficient evolution of unpolarized and polarized parton distributions with QCD-PEGASUS*, *Comput. Phys. Commun.* **170** (2005) 65, [hep-ph/0408244].
- [107] S. J. Brodsky, P. Hoyer, C. Peterson, and N. Sakai, *The intrinsic charm of the proton*, *Phys. Lett.* **B93** (1980) 451.
- [108] P. Jimenez-Delgado, T. J. Hobbs, J. T. Londergan, and W. Melnitchouk, *New limits on intrinsic charm in the nucleon from global analysis of parton distributions*, *Phys. Rev. Lett.* **114** (2015) 082002, [arXiv:1408.1708].
- [109] J. Blümlein, *A kinematic condition on intrinsic charm*, *Phys. Lett.* **B753** (2016) 619, [arXiv:1511.00229].
- [110] E. Witten, *Heavy quark contributions to deep-inelastic scattering*, *Nucl. Phys.* **B104** (1976) 445.
- [111] J. Babcock, D. W. Sivers, and S. Wolfram, *QCD estimates for heavy particle production*, *Phys. Rev.* **D18** (1978) 162.
- [112] V. A. Novikov, M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, *Charm photoproduction and Quantum Chromodynamics*, *Nucl. Phys.* **B136** (1978) 125. [*Yad. Fiz.*27,771(1978)].
- [113] J. P. Leveille and T. J. Weiler, *Characteristics of heavy quark leptonproduction in QCD*, *Nucl. Phys.* **B147** (1979) 147.
- [114] M. Glück, E. Hoffmann, and E. Reya, *Scaling violations and the gluon distribution of the nucleon*, *Z. Phys.* **C13** (1982) 119.
- [115] T. Gottschalk, *Chromodynamic corrections to neutrino production of heavy quarks*, *Phys. Rev.* **D23** (1981) 56.
- [116] M. Glück, S. Kretzer, and E. Reya, *The strange sea density and charm production in deep-inelastic charged current processes*, *Phys. Lett.* **B380** (1996) 171, [hep-ph/9603304]. [Erratum: *Phys. Lett.*B405,391(1997)].
- [117] J. Blümlein, A. Hasselhuhn, P. Kovacikova, and S. Moch,  *$O(\alpha_s)$  heavy flavor corrections to charged*

- current deep-inelastic scattering in Mellin space, *Phys. Lett.* **B700** (2011) 294, [arXiv:1104.3449].
- [118] E. Laenen, S. Riemersma, J. Smith, and W. L. van Neerven, *Complete  $O(\alpha_s)$  corrections to heavy flavor structure functions in electroproduction*, *Nucl. Phys.* **B392** (1993) 162.
- [119] E. Laenen, S. Riemersma, J. Smith, and W. L. van Neerven,  *$O(\alpha_s)$  corrections to heavy flavor inclusive distributions in electroproduction*, *Nucl. Phys.* **B392** (1993) 229.
- [120] S. Riemersma, J. Smith, and W. L. van Neerven, *Rates for inclusive deep-inelastic electroproduction of charm quarks at HERA*, *Phys. Lett.* **B347** (1995) 143, [hep-ph/9411431].
- [121] I. Bierenbaum, J. Blümlein, and S. Klein, *The gluonic operator matrix elements at  $O(\alpha_s^2)$  for DIS heavy flavor production*, *Phys. Lett.* **B672** (2009) 401, [arXiv:0901.0669].
- [122] A. Behring, I. Bierenbaum, J. Blümlein, A. De Freitas, S. Klein, and F. Wißbrock, *The logarithmic contributions to the  $O(\alpha_s^3)$  asymptotic massive Wilson coefficients and operator matrix elements in deeply inelastic scattering*, *Eur. Phys. J.* **C74** (2014) 3033, [arXiv:1403.6356].
- [123] I. Bierenbaum, J. Blümlein, and S. Klein, *Mellin moments of the  $O(\alpha_s^3)$  heavy flavor contributions to unpolarized deep-inelastic scattering at  $Q^2 \gg m^2$  and anomalous dimensions*, *Nucl. Phys.* **B820** (2009) 417, [arXiv:0904.3563].
- [124] S. Alekhin and S. Moch, *Heavy-quark deep-inelastic scattering with a running mass*, *Phys. Lett.* **B699** (2011) 345, [arXiv:1011.5790].
- [125] P. Marquard, A. V. Smirnov, V. A. Smirnov, and M. Steinhauser, *Quark mass relations to four-loop order in perturbative QCD*, *Phys. Rev. Lett.* **114** (2015) 142002, [arXiv:1502.01030].
- [126] J. Blümlein, A. De Freitas, W. L. van Neerven, and S. Klein, *The longitudinal heavy quark structure function  $F_L^{Q\bar{Q}}$  in the region  $Q^2 \gg m^2$  at  $O(\alpha_s^3)$* , *Nucl. Phys.* **B755** (2006) 272, [hep-ph/0608024].
- [127] J. Ablinger, J. Blümlein, S. Klein, C. Schneider, and F. Wißbrock, *The  $O(\alpha_s^3)$  massive operator matrix elements of  $O(n_f)$  for the structure function  $F_2(x, Q^2)$  and transversity*, *Nucl. Phys.* **B844** (2011) 26, [arXiv:1008.3347].
- [128] J. Blümlein, A. Hasselhuhn, S. Klein, and C. Schneider, *The  $O(\alpha_s^3 n_f T_F^2 C_{A,F})$  contributions to the gluonic massive operator matrix elements*, *Nucl. Phys.* **B866** (2013) 196, [arXiv:1205.4184].
- [129] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel, M. Round, C. Schneider, and F. Wißbrock, *The 3-loop non-singlet heavy flavor contributions and anomalous dimensions for the structure function  $F_2(x, Q^2)$  and transversity*, *Nucl. Phys.* **B886** (2014) 733, [arXiv:1406.4654].
- [130] H. Kawamura, N. A. Lo Presti, S. Moch, and A. Vogt, *On the next-to-next-to-leading order QCD corrections to heavy-quark production in deep-inelastic scattering*, *Nucl. Phys.* **B864** (2012) 399, [arXiv:1205.5727].
- [131] S. Catani, M. Ciafaloni, and F. Hautmann, *High-energy factorization and small  $x$  heavy flavor production*, *Nucl. Phys.* **B366** (1991) 135.
- [132] I. Bierenbaum, J. Blümlein, and S. Klein, *Two-loop massive operator matrix elements and unpolarized heavy flavor production at asymptotic values  $Q^2 \gg m^2$* , *Nucl. Phys.* **B780** (2007) 40, [hep-ph/0703285].
- [133] I. Bierenbaum, J. Blümlein, S. Klein, and C. Schneider, *Two-loop massive operator matrix elements for unpolarized heavy flavor production to  $O(\epsilon)$* , *Nucl. Phys.* **B803** (2008) 1, [arXiv:0803.0273].
- [134] T. Appelquist and J. Carazzone, *Infrared singularities and massive fields*, *Phys. Rev.* **D11** (1975) 2856.
- [135] K. G. Chetyrkin, J. H. Kühn, and M. Steinhauser, *RunDec: A Mathematica package for running and decoupling of the strong coupling and quark masses*, *Comput. Phys. Commun.* **133** (2000) 43, [hep-ph/0004189].

- [136] M. Buza, Y. Matiounine, J. Smith, and W. L. van Neerven, *Charm electroproduction viewed in the variable flavor number scheme versus fixed order perturbation theory*, *Eur. Phys. J.* **C1** (1998) 301–320, [hep-ph/9612398].
- [137] M. Buza, Y. Matiounine, J. Smith, R. Migneron, and W. L. van Neerven, *Heavy quark coefficient functions at asymptotic values  $Q^2 \gg m^2$* , *Nucl. Phys.* **B472** (1996) 611, [hep-ph/9601302].
- [138] J. Ablinger, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel, M. Round, C. Schneider, and F. Wissbrock, *The transition matrix element  $A_{gq}(N)$  of the variable flavor number scheme at  $\mathcal{O}(\alpha_s^3)$* , *Nucl. Phys.* **B882** (2014) 263, [arXiv:1402.0359].
- [139] A. G. Grozin, M. Höschele, J. Hoff, and M. Steinhauser, *Simultaneous decoupling of bottom and charm quarks*, *JHEP* **09** (2011) 066, [arXiv:1107.5970].
- [140] F. Maltoni, G. Ridolfi, and M. Ubiali, *b-initiated processes at the LHC: a reappraisal*, *JHEP* **07** (2012) 022, [arXiv:1203.6393]. [Erratum: JHEP04,095(2013)].
- [141] R. Harlander, M. Krämer, and M. Schumacher, *Bottom-quark associated Higgs-boson production: reconciling the four- and five-flavour scheme approach*, arXiv:1112.3478.
- [142] M. A. G. Aivazis, J. C. Collins, F. I. Olness, and W.-K. Tung, *Leptoproduction of heavy quarks. 2. A Unified QCD formulation of charged and neutral current processes from fixed target to collider energies*, *Phys. Rev.* **D50** (1994) 3102, [hep-ph/9312319].
- [143] M. Krämer, F. I. Olness, and D. E. Soper, *Treatment of heavy quarks in deeply inelastic scattering*, *Phys. Rev.* **D62** (2000) 096007, [hep-ph/0003035].
- [144] W.-K. Tung, S. Kretzer, and C. Schmidt, *Open heavy flavor production in QCD: Conceptual framework and implementation issues*, *J. Phys.* **G28** (2002) 983, [hep-ph/0110247].
- [145] S. Forte, E. Laenen, P. Nason, and J. Rojo, *Heavy quarks in deep-inelastic scattering*, *Nucl. Phys.* **B834** (2010) 116, [arXiv:1001.2312].
- [146] R. S. Thorne, *Effect of changes of variable flavor number scheme on parton distribution functions and predicted cross sections*, *Phys. Rev.* **D86** (2012) 074017, [arXiv:1201.6180].
- [147] S. Alekhin, J. Blümlein, and S. Moch, *Heavy-quark production in deep-inelastic scattering*, *PoS DIS2013* (2013) 297, [arXiv:1307.7258].
- [148] **ZEUS, H1** Collaboration, H. Abramowicz et al., *Combination and QCD analysis of charm production cross section measurements in deep-inelastic ep scattering at HERA*, *Eur. Phys. J.* **C73** (2013) 2311, [arXiv:1211.1182].
- [149] S. Alekhin, O. Behnke, P. Belov, S. Borroni, M. Botje, et al., **HERAFitter**, *Open source QCD fit project*, arXiv:1410.4412.
- [150] **xFitter**, *An open source QCD fit framework*, <http://xFitter.org> [xFitter.org].
- [151] S. Alekhin, J. Blümlein, K. Daum, K. Lipka, and S. Moch, *Precise charm-quark mass from deep-inelastic scattering*, *Phys. Lett.* **B720** (2013) 172, [arXiv:1212.2355].
- [152] R. D. Ball, V. Bertone, F. Cerutti, L. Del Debbio, S. Forte, A. Guffanti, J. I. Latorre, J. Rojo, and M. Ubiali, *Impact of heavy quark masses on parton distributions and LHC phenomenology*, *Nucl. Phys.* **B849** (2011) 296, [arXiv:1101.1300].
- [153] A. D. Martin, W. J. Stirling, R. S. Thorne, and G. Watt, *Heavy-quark mass dependence in global PDF analyses and 3- and 4-flavour parton distributions*, *Eur. Phys. J.* **C70** (2010) 51, [arXiv:1007.2624].
- [154] K. G. Chetyrkin, J. H. Kühn, A. Maier, P. Maierhöfer, P. Marquard, M. Steinhauser, and C. Sturm, *Charm and bottom quark masses: An update*, *Phys. Rev.* **D80** (2009) 074010, [arXiv:0907.2110].
- [155] B. Dehnadi, A. H. Hoang, and V. Mateu, *Bottom and charm mass determinations with a convergence test*, *JHEP* **08** (2015) 155, [arXiv:1504.07638].
- [156] Y. Kiyo, G. Mishima, and Y. Sumino, *Determination of  $m_c$  and  $m_b$  from quarkonium  $1S$  energy*

- levels in perturbative QCD, arXiv:1510.07072.
- [157] Y. Li and F. Petriello, *Combining QCD and electroweak corrections to dilepton production in FEWZ*, *Phys. Rev.* **D86** (2012) 094034, [arXiv:1208.5967].
  - [158] S. Catani, G. Ferrera, and M. Grazzini, *W Boson Production at Hadron Colliders: The Lepton Charge Asymmetry in NNLO QCD*, *JHEP* **05** (2010) 006, [arXiv:1002.3115].
  - [159] C. Balazs, J.-W. Qiu, and C. P. Yuan, *Effects of QCD resummation on distributions of leptons from the decay of electroweak vector bosons*, *Phys. Lett.* **B355** (1995) 548, [hep-ph/9505203].
  - [160] T. Carli, D. Clements, A. Cooper-Sarkar, C. Gwenlan, G. P. Salam, F. Siegert, P. Starovoitov, and M. Sutton, *A posteriori inclusion of parton density functions in NLO QCD final-state calculations at hadron colliders: The APPLGRID Project*, *Eur. Phys. J.* **C66** (2010) 503, [arXiv:0911.2985].
  - [161] **NNPDF** Collaboration, R. D. Ball, V. Bertone, F. Cerutti, L. Del Debbio, S. Forte, A. Guffanti, J. I. Latorre, J. Rojo, and M. Ubiali, *Unbiased global determination of parton distributions and their uncertainties at NNLO and at LO*, *Nucl. Phys.* **B855** (2012) 153, [arXiv:1107.2652].
  - [162] S. Alekhin, J. Blümlein, L. Caminadac, K. Lipka, K. Lohwasser, S. Moch, R. Petti, and R. Placakyte, *Determination of Strange Sea Quark Distributions from Fixed-target and Collider Data*, *Phys. Rev.* **D91** (2015) 094002, [arXiv:1404.6469].
  - [163] **NuTeV** Collaboration, M. Goncharov et al., *Precise measurement of dimuon production cross sections in muon neutrino Fe and muon anti-neutrino Fe deep-inelastic scattering at the Tevatron*, *Phys. Rev.* **D64** (2001) 112006, [hep-ex/0102049].
  - [164] A. Kayis-Topaksu et al., *Measurement of charm production in neutrino charged-current interactions*, *New J. Phys.* **13** (2011) 093002, [arXiv:1107.0613].
  - [165] **NOMAD** Collaboration, O. Samoylov et al., *A precision measurement of charm dimuon production in neutrino interactions from the NOMAD experiment*, *Nucl. Phys.* **B876** (2013) 339, [arXiv:1308.4750].
  - [166] **CMS** Collaboration, S. Chatrchyan et al., *Measurement of associated W+charm production in pp collisions at  $\sqrt{s} = 7$  TeV*, *JHEP* **02** (2014) 013, [arXiv:1310.1138].
  - [167] **ATLAS** Collaboration, G. Aad et al., *Measurement of the production of a W boson in association with a charm quark in pp collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector*, *JHEP* **05** (2014) 068, [arXiv:1402.6263].
  - [168] **ATLAS** Collaboration, G. Aad et al., *Determination of the strange quark density of the proton from ATLAS measurements of the  $W \rightarrow \ell\nu$  and  $Z \rightarrow \ell\ell$  cross sections*, *Phys. Rev. Lett.* **109** (2012) 012001, [arXiv:1203.4051].
  - [169] J. F. Owens, A. Accardi, and W. Melnitchouk, *Global parton distributions with nuclear and finite- $Q^2$  corrections*, *Phys. Rev.* **D87** (2013) 094012, [arXiv:1212.1702].
  - [170] A. Accardi, *PDFs from nucleons to nuclei*, *PoS DIS2015* (2015) 001, [arXiv:1602.02035].
  - [171] **NuSea** Collaboration, R. S. Towell et al., *Improved measurement of the  $\bar{d}/\bar{u}$  asymmetry in the nucleon sea*, *Phys. Rev.* **D64** (2001) 052002, [hep-ex/0103030].
  - [172] H. L. Lai, P. M. Nadolsky, J. Pumplin, D. Stump, W. K. Tung, and C. P. Yuan, *The strange parton distribution of the nucleon: Global analysis and applications*, *JHEP* **04** (2007) 089, [hep-ph/0702268].
  - [173] A. D. Martin, W. J. Stirling, R. S. Thorne, and G. Watt, *Parton distributions for the LHC*, *Eur. Phys. J.* **C63** (2009) 189, [arXiv:0901.0002].
  - [174] S. Alekhin, J. Blümlein, S. Klein, and S. Moch, *The 3, 4, and 5-flavor NNLO parton from deep-inelastic scattering data and at hadron colliders*, *Phys. Rev.* **D81** (2010) 014032, [arXiv:0908.2766].
  - [175] A. Accardi, M. E. Christy, C. E. Keppel, P. Monaghan, W. Melnitchouk, J. G. Morfin, and J. F.

- Owens, *New parton distributions from large- $x$  and low- $Q^2$  data*, *Phys. Rev.* **D81** (2010) 034016, [arXiv:0911.2254].
- [176] A. Accardi, W. Melnitchouk, J. F. Owens, M. E. Christy, C. E. Keppel, L. Zhu, and J. G. Morfin, *Uncertainties in determining parton distributions at large  $x$* , *Phys. Rev.* **D84** (2011) 014008, [arXiv:1102.3686].
- [177] J. Arrington, J. G. Rubin, and W. Melnitchouk, *How well do we know the neutron structure function?*, *Phys. Rev. Lett.* **108** (2012) 252001, [arXiv:1110.3362].
- [178] W. Melnitchouk, A. W. Schreiber, and A. W. Thomas, *Deep-inelastic scattering from off-shell nucleons*, *Phys. Rev.* **D49** (1994) 1183, [nucl-th/9311008].
- [179] W. Melnitchouk, A. W. Schreiber, and A. W. Thomas, *Relativistic deuteron structure function*, *Phys. Lett.* **B335** (1994) 11, [nucl-th/9407007].
- [180] S. A. Kulagin, G. Piller, and W. Weise, *Shadowing, binding and off-shell effects in nuclear deep-inelastic scattering*, *Phys. Rev.* **C50** (1994) 1154, [nucl-th/9402015].
- [181] S. A. Kulagin and R. Petti, *Global study of nuclear structure functions*, *Nucl. Phys.* **A765** (2006) 126, [hep-ph/0412425].
- [182] P. J. Ehlers, A. Accardi, L. T. Brady, and W. Melnitchouk, *Nuclear effects in the proton-deuteron Drell-Yan process*, *Phys. Rev.* **D90** (2014) 014010, [arXiv:1405.2039].
- [183] A. D. Martin, A. J. T. M. Mathijssen, W. J. Stirling, R. S. Thorne, B. J. A. Watt, and G. Watt, *Extended Parameterisations for MSTW PDFs and their effect on Lepton Charge Asymmetry from  $W$  Decays*, *Eur. Phys. J.* **C73** (2013) 2318, [arXiv:1211.1215].
- [184] S. Dasu, P. deBarbaro, A. Bodek, H. Harada, M. Krasny, et al., *Measurement of kinematic and nuclear dependence of  $R = \sigma_L/\sigma_T$  in deep-inelastic electron scattering*, *Phys. Rev.* **D49** (1994) 5641.
- [185] W. Giele et al., *The QCD/SM working group: Summary report*, hep-ph/0204316.
- [186] M. Botje, QCDNUM: *Fast QCD evolution and convolution*, *Comput. Phys. Commun.* **182** (2011) 490, [arXiv:1005.1481].
- [187] M. Botje, *Erratum for the time-like evolution in QCDNUM*, arXiv:1602.08383.
- [188] G. P. Salam and J. Rojo, *A higher order perturbative parton evolution toolkit (HOPPET)*, *Comput. Phys. Commun.* **180** (2009) 120, [arXiv:0804.3755].
- [189] T. Kluge, K. Rabbertz, and M. Wobisch, *fastNLO: Fast pQCD calculations for PDF fits*, hep-ph/0609285.
- [190] D. Britzger, K. Rabbertz, F. Stober, and M. Wobisch, *New features in version 2 of the fastNLO project*, arXiv:1208.3641.
- [191] OPENQCDRAD, <http://www-zeuthen.desy.de/~alekhin/OPENQCDRAD>.
- [192] D. A. Kosower, *Extracting parton densities from collider data*, *Nucl. Phys.* **B520** (1998) 263, [hep-ph/9708392].
- [193] M. Stratmann and W. Vogelsang, *Towards a global analysis of polarized parton distributions*, *Phys. Rev.* **D64** (2001) 114007, [hep-ph/0107064].
- [194] D. Graudenz, M. Hampel, A. Vogt, and C. Berger, *The Mellin transform technique for the extraction of the gluon density*, *Z. Phys.* **C70** (1996) 77, [hep-ph/9506333].
- [195] S. Bethke et al., *Workshop on precision measurements of  $\alpha_s$* , arXiv:1110.0016.
- [196] S. Moch et al., *High precision fundamental constants at the TeV scale*, arXiv:1405.4781.
- [197] D. d'Enterria et al., *High-precision  $\alpha_s$  measurements from LHC to FCC-ee*, arXiv:1512.05194.
- [198] **Particle Data Group** Collaboration  
<http://pdg.lbl.gov/2015/reviews/rpp2015-rev-qcd.pdf>.
- [199] J. Santiago and F. J. Yndurain, *Improved calculation of  $F_2$  in electroproduction and  $xF_3$  in neutrino scattering to NNLO and determination of  $\alpha_s$* , *Nucl. Phys.* **B611** (2001) 447, [hep-ph/0102247].

- [200] S. I. Alekhin, *Value of  $\alpha_s$  from deep-inelastic scattering data*, *JHEP* **02** (2003) 015, [[hep-ph/0211294](#)].
- [201] A. D. Martin, R. G. Roberts, W. J. Stirling, and R. S. Thorne, *Uncertainties of predictions from parton distributions. 2. Theoretical errors*, *Eur. Phys. J.* **C35** (2004) 325, [[hep-ph/0308087](#)].
- [202] J. Blümlein, H. Böttcher, and A. Guffanti, *Non-singlet QCD analysis of the structure function  $F_2$  in 3-loops*, *Nucl. Phys. Proc. Suppl.* **135** (2004) 152, [[hep-ph/0407089](#)].
- [203] J. Blümlein and H. Böttcher, *Higher twist contributions to the structure functions  $F_2(x, Q^2)$  and  $g_2(x, Q^2)$* , [arXiv:1207.3170](#).
- [204] M. Glück, E. Reya, and C. Schuck, *Non-singlet QCD analysis of  $F_2(x, Q^2)$  up to NNLO*, *Nucl. Phys.* **B754** (2006) 178, [[hep-ph/0604116](#)].
- [205] S. Alekhin, K. Melnikov, and F. Petriello, *Fixed target Drell-Yan data and NNLO QCD fits of parton distribution functions*, *Phys. Rev.* **D74** (2006) 054033, [[hep-ph/0606237](#)].
- [206] P. Jimenez-Delgado and E. Reya, *Dynamical NNLO parton distributions*, *Phys. Rev.* **D79** (2009) 074023, [[arXiv:0810.4274](#)].
- [207] A. D. Martin, W. J. Stirling, R. S. Thorne, and G. Watt, *Uncertainties on  $\alpha_s$  in global PDF analyses and implications for predicted hadronic cross sections*, *Eur. Phys. J.* **C64** (2009) 653, [[arXiv:0905.3531](#)].
- [208] S. Alekhin, J. Blümlein, and S. Moch, *Update of the NNLO PDFs in the 3-, 4-, and 5-flavour scheme*, *PoS DIS2010* (2010) 021, [[arXiv:1007.3657](#)].
- [209] S. Lionetti, R. D. Ball, V. Bertone, F. Cerutti, L. Del Debbio, S. Forte, A. Guffanti, J. I. Latorre, J. Rojo, and M. Ubiali, *Precision determination of  $\alpha_s$  using an unbiased global NLO parton set*, *Phys. Lett.* **B701** (2011) 346, [[arXiv:1103.2369](#)].
- [210] R. D. Ball, V. Bertone, L. Del Debbio, S. Forte, A. Guffanti, J. I. Latorre, S. Lionetti, J. Rojo, and M. Ubiali, *Precision NNLO determination of  $\alpha_s(M_Z)$  using an unbiased global parton set*, *Phys. Lett.* **B707** (2012) 66, [[arXiv:1110.2483](#)].
- [211] R. S. Thorne, *Theoretical procedures and the effect on PDFs and  $\alpha_s(M_Z^2)$* , *PoS DIS2013* (2013) 042, [[arXiv:1306.3907](#)].
- [212] L. A. Harland-Lang, A. D. Martin, P. Motylinski, and R. S. Thorne, *Uncertainties on  $\alpha_s$  in the MMHT2014 global PDF analysis and implications for SM predictions*, *Eur. Phys. J.* **C75** (2015) 435, [[arXiv:1506.05682](#)].
- [213] N. Kidonakis and J. F. Owens, *Effects of higher order threshold corrections in high  $E_T$  jet production*, *Phys. Rev.* **D63** (2001) 054019, [[hep-ph/0007268](#)].
- [214] S. Carrazza and J. Pires, *Perturbative QCD description of jet data from LHC Run-I and Tevatron Run-II*, *JHEP* **10** (2014) 145, [[arXiv:1407.7031](#)].
- [215] M. C. Kumar and S. Moch, *Phenomenology of threshold corrections for inclusive jet production at hadron colliders*, *Phys. Lett.* **B730** (2014) 122–129, [[arXiv:1309.5311](#)].
- [216] D. de Florian, P. Hinderer, A. Mukherjee, F. Ringer, and W. Vogelsang, *Approximate next-to-next-to-leading order corrections to hadronic jet production*, *Phys. Rev. Lett.* **112** (2014) 082001, [[arXiv:1310.7192](#)].
- [217] A. Vogt, *Structure function evolution at next-to-leading order and beyond*, *Nucl. Phys. Proc. Suppl.* **79** (1999) 102, [[hep-ph/9906337](#)].
- [218] **New Muon** Collaboration, M. Arneodo et al., *Measurement of the proton and deuteron structure functions,  $F_2^p$  and  $F_2^d$ , and of the ratio  $\sigma_L/\sigma_T$* , *Nucl. Phys.* **B483** (1997) 3, [[hep-ph/9610231](#)].
- [219] M. Spira, A. Djouadi, D. Graudenz, and P. M. Zerwas, *Higgs boson production at the LHC*, *Nucl. Phys.* **B453** (1995) 17, [[hep-ph/9504378](#)].
- [220] R. V. Harlander and W. B. Kilgore, *Next-to-next-to-leading order Higgs production at hadron*

- colliders*, *Phys. Rev. Lett.* **88** (2002) 201801, [hep-ph/0201206].
- [221] C. Anastasiou and K. Melnikov, *Higgs boson production at hadron colliders in NNLO QCD*, *Nucl. Phys.* **B646** (2002) 220, [hep-ph/0207004].
- [222] V. Ravindran, J. Smith, and W. L. van Neerven, *NNLO corrections to the total cross section for Higgs boson production in hadron hadron collisions*, *Nucl. Phys.* **B665** (2003) 325, [hep-ph/0302135].
- [223] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog, and B. Mistlberger, *Higgs boson gluon fusion production in QCD at three loops*, *Phys. Rev. Lett.* **114** (2015) 212001, [arXiv:1503.06056].
- [224] D. de Florian, J. Mazzitelli, S. Moch, and A. Vogt, *Approximate  $N^3LO$  Higgs-boson production cross section using physical-kernel constraints*, *JHEP* **10** (2014) 176, [arXiv:1408.6277].
- [225] R. D. Ball et al., *Parton distributions with LHC data*, *Nucl. Phys.* **B867** (2013) 244, [arXiv:1207.1303].
- [226] P. Bärnreuther, M. Czakon, and A. Mitov, *Percent Level Precision Physics at the Tevatron: First Genuine NNLO QCD Corrections to  $q\bar{q} \rightarrow t\bar{t} + X$* , *Phys. Rev. Lett.* **109** (2012) 132001, [arXiv:1204.5201].
- [227] M. Czakon and A. Mitov, *NNLO corrections to top-pair production at hadron colliders: the all-fermionic scattering channels*, *JHEP* **12** (2012) 054, [arXiv:1207.0236].
- [228] M. Czakon and A. Mitov, *NNLO corrections to top pair production at hadron colliders: the quark-gluon reaction*, *JHEP* **01** (2013) 080, [arXiv:1210.6832].
- [229] M. Czakon, P. Fiedler, and A. Mitov, *Total top-quark pair-production cross section at hadron colliders through  $O(\alpha_s^4)$* , *Phys. Rev. Lett.* **110** (2013) 252004, [arXiv:1303.6254].
- [230] U. Langenfeld, S. Moch, and P. Uwer, *Measuring the running top-quark mass*, *Phys. Rev.* **D80** (2009) 054009, [arXiv:0906.5273].
- [231] M. Aliev, H. Lacker, U. Langenfeld, S. Moch, P. Uwer, and M. Wiedermann, *HATHOR: HAdronic Top and Heavy quarks crOss section calculatoR*, *Comput. Phys. Commun.* **182** (2011) 1034, [arXiv:1007.1327].
- [232] M. Dowling and S. Moch, *Differential distributions for top-quark hadro-production with a running mass*, *Eur. Phys. J.* **C74** (2014) 3167, [arXiv:1305.6422].
- [233] CMS Collaboration, S. Chatrchyan et al., *Determination of the top-quark pole mass and strong coupling constant from the  $t\bar{t}$  production cross section in  $pp$  collisions at  $\sqrt{s} = 7$  TeV*, *Phys. Lett.* **B728** (2014) 496, [arXiv:1307.1907]. [Erratum: *Phys. Lett.* **B728**, 526 (2014)].
- [234] M. Czakon, M. L. Mangano, A. Mitov, and J. Rojo, *Constraints on the gluon PDF from top quark pair production at hadron colliders*, *JHEP* **07** (2013) 167, [arXiv:1303.7215].
- [235] J. Kieseler, K. Lipka, and S. Moch, *Calibration of the top-quark Monte Carlo mass*, arXiv:1511.00841.
- [236] M. Czakon, D. Heymes, and A. Mitov, *High-precision differential predictions for top-quark pairs at the LHC*, arXiv:1511.00549.
- [237] M. Guzzi, K. Lipka, and S. Moch, *Top-quark pair production at hadron colliders: differential cross section and phenomenological applications with DiffTop*, *JHEP* **01** (2015) 082, [arXiv:1406.0386].
- [238] LHCb Collaboration, R. Aaij et al., *Measurement of B meson production cross sections in proton-proton collisions at  $\sqrt{s} = 7$  TeV*, *JHEP* **08** (2013) 117, [arXiv:1306.3663].
- [239] O. Zenaiev, A. Geiser, K. Lipka, J. Blümlein, A. Cooper-Sarkar, et al., *Impact of heavy-flavour production cross sections measured by the LHCb experiment on parton distribution functions at low  $x$* , arXiv:1503.04581.
- [240] O. Zenaiev, *Charm production and QCD analysis at HERA and LHC*. PhD thesis, U. Hamburg,



- Dept. Phys., DESY-THESIS-2015-012, 2015.
- [241] P. Nason, S. Dawson, and R. K. Ellis, *The total cross section for the production of heavy quarks in hadronic collisions*, *Nucl. Phys.* **B303** (1988) 607.
- [242] W. Beenakker, H. Kuijf, W. L. van Neerven, and J. Smith, *QCD corrections to heavy quark production in  $p\bar{p}$  collisions*, *Phys. Rev.* **D40** (1989) 54.
- [243] P. Nason, S. Dawson, and R. K. Ellis, *The one particle inclusive differential cross section for heavy quark production in hadronic collisions*, *Nucl. Phys.* **B327** (1989) 49. [Erratum: *Nucl. Phys.* **B335**, 260(1990)].
- [244] M. V. Garzelli, S. Moch, and G. Sigl, *Lepton fluxes from atmospheric charm revisited*, *JHEP* **10** (2015) 115, [arXiv:1507.01570].
- [245] R. Gauld, J. Rojo, L. Rottoli, and J. Talbert, *Charm production in the forward region: constraints on the small- $x$  gluon and backgrounds for neutrino astronomy*, *JHEP* **11** (2015) 009, [arXiv:1506.08025].
- [246] C. Lourenco and H. Wöhri, *Heavy flavour hadro-production from fixed-target to collider energies*, *Phys.Rept.* **433** (2006) 127, [hep-ph/0609101].
- [247] **HERA-B** Collaboration, I. Abt et al., *Measurement of  $D^0$ ,  $D^+$ ,  $D_s^+$  and  $D^{*+}$  production in fixed target 920-GeV proton-nucleus collisions*, *Eur.Phys.J.* **C52** (2007) 531, [arXiv:0708.1443].
- [248] **PHENIX** Collaboration, A. Adare et al., *Measurement of high- $p_T$  single electrons from heavy-flavor decays in  $p+p$  collisions at  $s^{1/2} = 200$  GeV*, *Phys.Rev.Lett.* **97** (2006) 252002, [hep-ex/0609010].
- [249] **STAR** Collaboration, L. Adamczyk et al., *Measurements of  $D^0$  and  $D^*$  production in  $p+p$  collisions at  $\sqrt{s} = 200$  GeV*, *Phys.Rev.* **D86** (2012) 072013, [arXiv:1204.4244].
- [250] **ALICE** Collaboration, B. Abelev et al., *Measurement of charm production at central rapidity in proton-proton collisions at  $\sqrt{s} = 2.76$  TeV*, *JHEP* **1207** (2012) 191, [arXiv:1205.4007].
- [251] **ATLAS** Collaboration, *Measurement of  $D^{(*)}$  meson production cross sections in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector*, ATLAS-CONF-2011-017, ATLAS-COM-CONF-2011-030.
- [252] **LHCb** Collaboration, R. Aaij et al., *Prompt charm production in  $pp$  collisions at  $\sqrt{s} = 7$  TeV*, *Nucl.Phys.* **B871** (2013) 1, [arXiv:1302.2864].
- [253] **LHCb** Collaboration, R. Aaij et al., *Measurements of prompt charm production cross sections in  $pp$  collisions at  $\sqrt{s} = 13$  TeV*, arXiv:1510.01707.
- [254] L. T. Brady, A. Accardi, W. Melnitchouk, and J. F. Owens, *Impact of PDF uncertainties at large  $x$  on heavy boson production*, *JHEP* **06** (2012) 019, [arXiv:1110.5398].
- [255] M. Botje et al., *The PDF4LHC working group interim recommendations*, arXiv:1101.0538.
- [256] A. Buckley and J. Rojo, *private communication*, .
- [257] A. Buckley, J. Ferrando, S. Lloyd, K. Nordström, B. Page, M. Rüfenacht, M. Schönherr, and G. Watt, *LHAPDF6: parton density access in the LHC precision era*, *Eur. Phys. J.* **C75** (2015) 132, [arXiv:1412.7420].
- [258] M. R. Whalley, D. Bourilkov, and R. C. Group, *The Les Houches accord PDFs (LHAPDF) and LHAGLUE*, in *HERA and the LHC: A workshop on the implications of HERA for LHC physics. Proceedings, Part B*, 2005. hep-ph/0508110.
- [259] J. P. Ralston, *Pocket partonometer*, *Phys. Lett.* **B172** (1986) 430.
- [260] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079, [arXiv:1405.0301].
- [261] POWHEG-BOX (v2), <http://powhegbox.mib.infn.it/>.
- [262] S. Alioli, P. Nason, C. Oleari, and E. Re, *A general framework for implementing NLO calculations*

- in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043, [arXiv:1002.2581].
- [263] SHERPA (v2), <https://sherpa.hepforge.org/doc/SHERPA-MC-2.2.0.html>.
- [264] T. Gleisberg, S. Höche, F. Krauss, M. Schönherr, S. Schumann, F. Siegert, and J. Winter, *Event generation with SHERPA 1.1*, *JHEP* **02** (2009) 007, [arXiv:0811.4622].
- [265] S. Alioli, C. W. Bauer, C. Berggren, F. J. Tackmann, and J. R. Walsh, *Drell-Yan production at NNLL'+NNLO matched to parton showers*, *Phys. Rev.* **D92** (2015) 094020, [arXiv:1508.01475].