

A Higgs Boson at 96 GeV?!

S. Heinemeyer*

*Instituto de Física Teórica, (UAM/CSIC), Universidad Autónoma de Madrid,
Cantoblanco, E-28049 Madrid, Spain*

Campus of International Excellence UAM+CSIC, Cantoblanco, E-28049, Madrid, Spain

Instituto de Física de Cantabria (CSIC-UC), E-39005 Santander, Spain

E-mail: Sven.Heinemeyer@cern.ch

T. Stefaniak

DESY, Notkestraße 85, D-22607 Hamburg, Germany

E-mail: tim.stefaniak@desy.de

We briefly summarize some searches for Higgs bosons with a mass of $m_\phi \lesssim 110$ GeV at LEP and the LHC. We discuss a possible signal in the diphoton decay mode at $m_\phi \sim 96$ GeV as reported by CMS, together with a $\sim 2\sigma$ hint in the $b\bar{b}$ final state at LEP. We briefly review possible interpretation of such a new particle in various BSM models. We focus on possible explanations as reported within the NMSSM and the $\mu\nu$ SVM. Conclusions for future collider projects are briefly outlined.

Prospects for Charged Higgs Discovery at Colliders - CHARGED2018

25-28 September 2018

Uppsala, Sweden

*Speaker.

1. Introduction

In the year 2012 the ATLAS and CMS collaborations have discovered a new particle that – within theoretical and experimental uncertainties – is consistent with the existence of a Standard-Model (SM) Higgs boson at a mass of ~ 125 GeV [1–3]. No conclusive signs of physics beyond the SM have been found so far at the LHC. However, the measurements of Higgs-boson couplings, which are known experimentally to a precision of roughly $\sim 20\%$, leave room for Beyond Standard Model (BSM) interpretations. Many BSM models possess extended Higgs-boson sectors. Consequently, one of the main tasks of the LHC Run II and beyond will be to determine whether the observed scalar boson forms part of the Higgs sector of an extended model.

Motivated by the Hierarchy Problem, Supersymmetry (SUSY) extensions of the SM play a prominent role in the exploration of new physics. SUSY doubles the particle degrees of freedom by predicting two scalar partners for all SM fermions, as well as fermionic partners to all SM bosons. The simplest SUSY extension is the Minimal Supersymmetric Standard Model (MSSM) [4, 5]. In contrast to the single Higgs doublet of the SM, in the MSSM two Higgs doublets, H_u and H_d , are required. In the $\mathcal{C}\mathcal{P}$ conserving case the MSSM Higgs sector consists of two $\mathcal{C}\mathcal{P}$ -even, one $\mathcal{C}\mathcal{P}$ -odd and two charged Higgs bosons. The light (or the heavy) $\mathcal{C}\mathcal{P}$ -even MSSM Higgs boson can be interpreted as the signal discovered at ~ 125 GeV [6] (see Refs. [7, 8] for recent updates).

Going beyond the MSSM, a well-motivated extension is given by the Next-to-MSSM (NMSSM), see e.g. [9, 10] for reviews. The NMSSM provides a solution for the so-called “ μ problem” by naturally associating an adequate scale to the μ parameter appearing in the MSSM superpotential [11, 12]. In the NMSSM a new singlet superfield is introduced, which only couples to the Higgs- and sfermion-sectors, giving rise to an effective μ -term, proportional to the vacuum expectation value (vev) of the scalar singlet. In the $\mathcal{C}\mathcal{P}$ conserving case the NMSSM Higgs sector consists of three $\mathcal{C}\mathcal{P}$ -even Higgs bosons, h_i ($i = 1, 2, 3$), two $\mathcal{C}\mathcal{P}$ -odd Higgs bosons, a_j ($j = 1, 2$), and the charged Higgs boson pair H^\pm . In the NMSSM the lightest but also the second lightest $\mathcal{C}\mathcal{P}$ -even Higgs boson can be interpreted as the signal observed at about 125 GeV, see, e.g., [13, 14].

A natural extension of the NMSSM is the $\mu\nu$ SUSM, in which the singlet superfield is interpreted as a right-handed neutrino superfield [15, 16] (here we focus on the “one generation case”), see Refs. [17–19] for reviews. The $\mu\nu$ SUSM is the simplest extension of the MSSM that can provide massive neutrinos through a see-saw mechanism at the electroweak scale. A Yukawa coupling for the right-handed neutrino of the order of the electron Yukawa coupling is introduced that induces the explicit breaking of R -parity. One consequence is that there is no lightest stable SUSY particle anymore. Nevertheless, the model can still provide a dark matter candidate with a gravitino that has a life time longer than the age of the observable universe [20–23]. The explicit violation of lepton number and lepton flavor can modify the spectrum of the neutral and charged fermions in comparison to the NMSSM. The three families of charged leptons will mix with the chargino and the Higgsino and form five massive charged fermions. Within the scalar sector, due to R -parity breaking, the left- and right-handed sneutrinos will mix with the doublet Higgses and form six massive $\mathcal{C}\mathcal{P}$ -even and five massive $\mathcal{C}\mathcal{P}$ -odd states, assuming that there is no $\mathcal{C}\mathcal{P}$ -violation. Also in the $\mu\nu$ SUSM the signal at ~ 125 GeV can be interpreted as the lightest or the second lightest $\mathcal{C}\mathcal{P}$ -even scalar. As SUSY models, but also other BSM Higgs-boson sector extensions can possess a scalar below 125 GeV the search for such light scalars is of high importance at the LHC.

2. Experimental data

Searches for Higgs bosons below 125 GeV have been performed at LEP, the Tevatron and the LHC. LEP reported a 2.3σ local excess observed in the $e^+e^- \rightarrow Z(H \rightarrow b\bar{b})$ searches [24], which would be consistent with a scalar mass of ~ 98 GeV (but due to the final state the mass resolution is rather coarse). The “excess” corresponds to

$$\mu_{\text{LEP}} = \frac{\sigma(e^+e^- \rightarrow Z\phi \rightarrow Zb\bar{b})}{\sigma^{\text{SM}}(e^+e^- \rightarrow ZH_{\text{SM}} \rightarrow Zb\bar{b})} = 0.117 \pm 0.057, \quad (2.1)$$

where the signal strength μ_{LEP} is the measured cross section normalized to the SM expectation, with the SM Higgs-boson mass at ~ 96 GeV. The value for μ_{LEP} was extracted in Ref. [25] using methods described in Ref. [26].

Interestingly, recent CMS Run II results [27] for Higgs searches in the diphoton final state show a local excess of $\sim 3\sigma$ around ~ 96 GeV, with a similar excess of 2σ in the Run I data at a comparable mass. In this case the “excess” corresponds to (combining 7, 8 and 13 TeV data)

$$\mu_{\text{CMS}} = \frac{\sigma(gg \rightarrow \phi \rightarrow \gamma\gamma)}{\sigma^{\text{SM}}(gg \rightarrow H_{\text{SM}} \rightarrow \gamma\gamma)} = 0.6 \pm 0.2. \quad (2.2)$$

First Run II results from ATLAS with 80 fb^{-1} in the $\gamma\gamma$ searches below 125 GeV were recently published [28]. No significant excess above the SM expectation was observed in the mass range between 65 and 110 GeV. However, the limit on cross section times branching ratio obtained in the diphoton final state by ATLAS is not only well above μ_{CMS} , but even weaker than the corresponding upper limit obtained by CMS at ~ 96 GeV. This is illustrated in Fig. 1, where we compare the expected (dashed) and observed (solid) limits in the $gg \rightarrow \phi \rightarrow \gamma\gamma$ channel (normalized to the SM value) as reported by CMS (red) and ATLAS (blue) as a function of m_ϕ . Shown in magenta is μ_{CMS} of Eq. (2.2). The “weaker” expected (and observed) exclusion around 91 GeV corresponds to the Z peak, where a larger background is expected.

3. BSM models

Several analyses attempted to explain the *combined* “excess” of LEP and CMS in a variety of BSM models¹. To our knowledge explanations exist in the following frameworks (see also [29]):

- Higgs singlet with additional vector-like matter, as well as Type-I 2HDM [30].
- Radion model [31].
- Type-I 2HDM with a moderately-to-strongly fermiophobic $\mathcal{C}\mathcal{P}$ -even Higgs [32].
- $\mu\nu\text{SSM}$ [33], as will be discussed in Sect. 3.2.
- Higgs associated with the breakdown of an $U(1)_{L_\mu L_\tau}$ symmetry [34].
- NMSSM [35], as will be discussed in Sect. 3.1.
- Higgs inflation inspired μNMSSM [36].

On the other hand, in the MSSM the CMS excess cannot be realized [7].

¹More analyses attempted to explain one of the two “excesses”, but we will not discuss these further here.




Figure 1: Limits on the cross section $gg \rightarrow \phi \rightarrow \gamma\gamma$ normalized to the SM value as a function of $m_\phi (\equiv M_h)$. Compared are the expected (dashed) and observed (solid) limits from CMS (red) and ATLAS (blue). Shown in magenta is $\mu_{\text{CMS}} = 0.6 \pm 0.2$.

3.1 The NMSSM solution

The results in this section are based on Ref. [35]. Within the NMSSM a natural candidate to explain the LEP “excess” consists in a mostly singlet-like Higgs with a doublet component of about 10% (mixing squared). Relatively large Higgs branching fractions into $\gamma\gamma$ are possible due to the three-state mixing, in particular when the effective Higgs coupling to $b\bar{b}$ becomes small, see e.g. Refs. [37,38]. In our numerical analysis we display the quantities ξ_b and ξ_γ , defined as follows:

$$\begin{aligned} \xi_b &\equiv \frac{\Gamma(h_1 \rightarrow ZZ) \cdot \text{BR}(h_1 \rightarrow b\bar{b})}{\Gamma(H_{\text{SM}}(M_{h_1}) \rightarrow ZZ) \cdot \text{BR}(H_{\text{SM}}(M_{h_1}) \rightarrow b\bar{b})} \sim \frac{\sigma(e^+e^- \rightarrow Z(h_1 \rightarrow b\bar{b}))}{\sigma(e^+e^- \rightarrow Z(H_{\text{SM}}(M_{h_1}) \rightarrow b\bar{b}))} \\ \xi_\gamma &\equiv \frac{\Gamma(h_1 \rightarrow gg) \cdot \text{BR}(h_1 \rightarrow \gamma\gamma)}{\Gamma(H_{\text{SM}}(M_{h_1}) \rightarrow gg) \cdot \text{BR}(H_{\text{SM}}(M_{h_1}) \rightarrow \gamma\gamma)} \sim \frac{\sigma(gg \rightarrow h_1 \rightarrow \gamma\gamma)}{\sigma(gg \rightarrow H_{\text{SM}}(M_{h_1}) \rightarrow \gamma\gamma)}. \end{aligned} \quad (3.1)$$

These definitions of $\xi_{b,\gamma}$ give estimates of the signals that h_1 would generate in the LEP searches for $e^+e^- \rightarrow Z(H \rightarrow b\bar{b})$ and the LHC searches for $pp \rightarrow H \rightarrow \gamma\gamma$, normalized to the SM cross-sections. In the analysis in Ref. [35] constraints from “other sectors” (such as Dark Matter or $(g-2)_\mu$) are not taken into account, as they are not closely related to Higgs sector physics.

The NMSSM parameters are chosen as (see Ref. [35] for definitions and details),

$$\begin{aligned} \lambda &= 0.6, \quad \kappa = 0.035, \quad \tan\beta = 2, \quad M_{H^\pm} = 1000 \text{ GeV}, \quad A_\kappa = -325 \text{ GeV}, \\ \mu_{\text{eff}} &= (397 + 15 \cdot x) \text{ GeV} \quad (x \text{ is varied in the interval } [0, 1]), \\ \text{the third generation squark mass scale } m_{\tilde{Q}} &= 1000 \text{ GeV}, A_t = A_b = 0. \end{aligned}$$

In our analysis we vary μ_{eff} in a narrow interval as indicated above. It was tested with HiggsBounds-4.3.1 (and 5.1.1beta) [39–43] and HiggsSignals-1.3.1 (and 2.1.0beta) [43–46] that our parameter points are in agreement with the Higgs rate measurements at the LHC as well as with the Higgs boson searches at LEP, the Tevatron and the LHC.

With growing μ_{eff} , the mixing between the two light $\mathcal{C}\mathcal{P}$ -even states increases, eventually pushing the singlet mass down to ~ 90 GeV and the mass of the SM-like state up to ~ 128 GeV. Consistency with the experimental results obtained on the observed state at 125 GeV is achieved for a mass of the SM-like state that is compatible with the LHC discovery within experimental and theoretical uncertainties. The $\mathcal{C}\mathcal{P}$ -odd singlet has a mass of ~ 150 GeV, while the heavy doublet states are at ~ 1 TeV in this scenario. The decay properties of h_1 and h_2 are given in Tab. 1 (for a specific point). The quantities ξ_b and ξ_γ of Eq. (3.1), are shown in Fig. 2, estimating the signals associated with h_1 in the $b\bar{b}$ channel at LEP and in the $\gamma\gamma$ channel at the LHC, as compared to an SM Higgs at the same mass. The magnitude of the estimated $e^+e^- \rightarrow Z(h_1 \rightarrow b\bar{b})$ signal reaches $\sim 13\%$ of that of an SM Higgs at $M_{h_1} \sim 95$ GeV, while $pp \rightarrow h_1 \rightarrow \gamma\gamma$ corresponds to more than 40% of an SM signal in the same mass range. In this example, $\text{BR}(h_1 \rightarrow \gamma\gamma)$ (or $\text{BR}(h_1 \rightarrow gg)$) is only moderately enhanced with respect to the SM branching fraction due to an H_u^0 -dominated doublet composition of h_1 , while $\text{BR}(h_1 \rightarrow b\bar{b})$ remains dominant, albeit slightly suppressed. This scenario would thus simultaneously address the LEP and the CMS excesses in a phenomenologically consistent manner.

Table 1: The Higgs properties for one example point in the NMSSM. The Higgs width into xx' is denoted by $\Gamma_{xx'}$, and the width normalized to the SM width at the same mass is represented by $\widehat{\Gamma}_{xx'}$. The symbol $\widehat{\text{BR}}_{xx'}$ represents the Higgs branching ratio into xx' , normalized to the SM branching ratio at the same mass.

	h_1	h_2		h_1
$\Gamma_{\gamma\gamma}$ [GeV]	$6.3 \cdot 10^{-7}$	$7.6 \cdot 10^{-6}$	$\widehat{\text{BR}}_{\gamma\gamma}$	2.2
$\Gamma_{b\bar{b}}$ [GeV]	$1.4 \cdot 10^{-4}$	$2.1 \cdot 10^{-3}$	$\widehat{\text{BR}}_{b\bar{b}}$	0.88
Γ_{gg} [GeV]	$2.2 \cdot 10^{-5}$	$2.2 \cdot 10^{-4}$	$\widehat{\Gamma}_{gg}$	0.18
Γ_{ZZ} [GeV]	$1.0 \cdot 10^{-7}$	$9.7 \cdot 10^{-5}$	$\widehat{\Gamma}_{ZZ}$	0.15
Γ_{WW} [GeV]	$8.4 \cdot 10^{-6}$	$7.9 \cdot 10^{-4}$	ξ_γ	0.41
$\Gamma_{\tau\tau}$ [GeV]	$1.5 \cdot 10^{-5}$	$2.4 \cdot 10^{-4}$	ξ_b	0.13
$\Gamma_{c\bar{c}}$ [GeV]	$1.8 \cdot 10^{-5}$	$1.2 \cdot 10^{-4}$		
M_{h_i} [GeV]	95.0	125.5		

3.2 The $\mu\nu$ S SM solution

The results in this section are based on Ref. [33]. Within the $\mu\nu$ S SM (in the simplified “one generation case”) we will interpret the light scalar as the $\mathcal{C}\mathcal{P}$ -even right-handed sneutrino. Since the singlet of the NMSSM and the right-handed sneutrino of the $\mu\nu$ S SM are both gauge-singlets, they share very similar properties. However, the explanation of the excesses in the $\mu\nu$ S SM avoids bounds from direct detection experiments, because R -parity is broken in the $\mu\nu$ S SM and the dark matter candidate is not a neutralino as in the NMSSM but a gravitino with a lifetime longer than the age of the universe [18]. This can be important since the direct detection measurements were shown to be very constraining in the NMSSM while trying to explain the dark matter abundance on top of the excesses from LEP and CMS [25].

In Tab. 2 we list the values of the parameters we used to account for the lightest $\mathcal{C}\mathcal{P}$ -even scalar as the right-handed sneutrino and the second lightest one the SM-like Higgs boson (see




Figure 2: The quantities ξ_b and ξ_γ of Eq. (3.1), estimating the signals associated with h_1 in the $b\bar{b}$ channel at LEP and in the $\gamma\gamma$ channel at the LHC, as compared to an SM Higgs at the same mass. Explicit values at $M_{h_1} = 95.0$ GeV are given in Tab. 1.

Ref. [33] for definitions and details). As in the NMSSM `HiggsBounds` and `HiggsSignals` were used to ensure the compatibility with experimental data. In the analysis in Ref. [33] constraints from “other sectors” (such as flavor physics or $(g-2)_\mu$) are not taken into account, as they are not closely related to Higgs sector physics. λ is chosen to be large to account for a sizable mixing of the right-handed sneutrino and the doublet Higgs bosons. In the regime where the SM-like Higgs boson is not the lightest scalar, one does not need large quantum corrections to the Higgs boson mass (which were evaluated according to Ref. [33]), because the tree-level mass is already well above 100 GeV. This is why $\tan\beta$ can be low and the soft trilinear couplings $A^{u,d,e}$ are set to zero. The values of A^λ and $-A^V$ are chosen to be around 1 TeV to get masses for the heavy MSSM-like Higgs and the left-handed sneutrinos of this order, so they do not play an important role in the following discussion. On the other hand, κ is small to bring the mass of the right-handed sneutrino below the SM-like Higgs boson mass. Finally, the two parameters that are varied are μ and A^κ . By increasing μ the mixing of the right-handed sneutrino with the SM-like Higgs boson is increased, which is needed to couple the gauge-singlet to quarks and gauge-bosons. At the same time we used the value of A^κ to keep the mass of the right-handed sneutrino in the correct range.

The result are shown in Fig. 3, the CMS (left) and the LEP excesses (right) in the μ - A^κ plane. While the LEP excess is easily reproduced in the observed parameter space, we cannot achieve the central value for μ_{CMS} , but only slightly smaller values. As already observed in Ref. [25], the reason for this is that for explaining the LEP excess a sizable coupling to the bottom quark is

Table 2: Input parameters for the scenario featuring the right-handed sneutrino in the mass range of the LEP and CMS excesses and a SM-like Higgs boson as next-to-lightest $\mathcal{C}\mathcal{P}$ -even scalar; all masses and values for trilinear parameters are in GeV.

$v_{iL}/\sqrt{2}$	Y_i^V	A_i^V	$\tan\beta$	μ	λ	A^λ	κ	A^κ	M_1
10^{-5}	10^{-7}	-1000	2	[413; 418]	0.6	956	0.035	[-300; -318]	100
M_2	M_3	$m_{Q_{iL}}^2$	$m_{U_{iR}}^2$	$m_{D_{iR}}^2$	A_i^u	A_i^d	$(m_e^2)_{ii}$	A_{33}^e	$A_{11,22}^e$
200	1500	800^2	800^2	800^2	0	0	800^2	0	0




Figure 3: Signal strengths for the lightest $\tilde{\nu}_R$ -like neutral scalar at CMS ($pp \rightarrow h_1 \rightarrow \gamma\gamma$) (left) and LEP ($e^+e^- \rightarrow h_1 Z \rightarrow b\bar{b}Z$) (right) in the μ - A^κ plane. The gray area is excluded because the right-handed sneutrino becomes tachyonic at tree-level.

needed. On the contrary, the CMS excess demands a small value for the $h_1 b\bar{b}$ coupling so that the total width of the h_1 becomes small and $h_1 \rightarrow \gamma\gamma$ is enhanced. Nevertheless, considering the large experimental uncertainties in μ_{CMS} and μ_{LEP} , the scenario reviewed in this section accommodates both excesses comfortably well (at approximately 1σ).

4. Conclusions (for future colliders)

Searches for Higgs bosons below 125 GeV have been performed at LEP, the Tevatron and the LHC. We have briefly reviewed that LEP reported a 2.3σ local excess observed in the $e^+e^- \rightarrow Z(H \rightarrow b\bar{b})$ searches [24], which would be consistent with a scalar mass of ~ 98 GeV (but with a rather coarse mass resolution). Furthermore, recent LHC Run II results [27] for CMS Higgs searches in the diphoton final state show a local excess of $\sim 3\sigma$ in the vicinity of ~ 96 GeV, with a similar upward fluctuation of 2σ in the Run I data at a comparable mass. First Run II results from ATLAS with 80 fb^{-1} in the $\gamma\gamma$ searches below 125 GeV are well compatible with the limit obtained by CMS at ~ 96 GeV (although not showing a relevant excess).

We have briefly reviewed BSM interpretations, explaining simultaneously the LEP and the CMS “excess”. In particular we have reviewed the solution within the NMSSM and the $\mu\nu$ SVM. Within the NMSSM we investigated the case of a mostly singlet-like state with a mass of $\lesssim 96$ GeV. The decays of such a state can be notably affected by suppressed couplings to down- or up-type quarks which can occur in certain parameter regions due to the mixing between the different Higgs states. In particular, an additional Higgs boson h_i of this kind could manifest itself via signatures in the channels $e^+e^- \rightarrow Z(h_i \rightarrow b\bar{b})$ and/or $pp \rightarrow h_i \rightarrow \gamma\gamma$. The presence of such a light Higgs boson could thus explain the “excesses” reported by LEP and CMS in those channels.

Within the $\mu\nu$ SVM we reviewed that this model can accommodate a right-handed ($\mathcal{C}\mathcal{P}$ -even) scalar neutrino with a mass of ~ 96 GeV, where the full Higgs sector is in agreement with the Higgs-boson measurements and exclusion bounds obtained at the LHC, as well as at LEP and the Tevatron. It was demonstrated that the light right-handed sneutrino can explain an excess

of $\gamma\gamma$ events at ~ 96 GeV as reported recently by CMS in their Run I and Run II date. It can simultaneously describe the 2σ excess of $b\bar{b}$ events observed at LEP at a similar mass scale.

A new Higgs boson with a mass of ~ 96 GeV is easily kinematically accessible at future e^+e^- colliders, assuming an energy of at least $\sqrt{s} = 250$ GeV. It was shown in Ref. [47] that a light Higgs boson explaining the LEP “excess” is easily within the reach of the ILC250. Further confirmation of these “excesses” would strengthen the already robust physics case for such a machine.

Acknowledgements

S.H. thanks T. Biekötter, F. Domingo, C. Muñoz, S. Paßehr and G. Weiglein, with whom the NMSSM and μ vSSM results presented here have been obtained. S.H. thanks K. Tackmann for interesting discussions. The work of S.H. was supported in part by the MEINCOP (Spain) under contract FPA2016-78022-P, in part by the Spanish Agencia Estatal de Investigación (AEI), in part by the EU Fondo Europeo de Desarrollo Regional (FEDER) through the project FPA2016-78645-P, in part by the “Spanish Red Consolider MultiDark” FPA2017-90566-REDC, and in part by the AEI through the grant IFT Centro de Excelencia Severo Ochoa SEV-2016-0597.

References

- [1] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716** (2012) 1 [arXiv:1207.7214 [hep-ex]].
- [2] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716** (2012) 30 [arXiv:1207.7235 [hep-ex]].
- [3] G. Aad *et al.* [ATLAS and CMS Collab.], JHEP **1608** (2016) 045 [arXiv:1606.02266 [hep-ex]].
- [4] H. P. Nilles, Phys. Rept. **110** (1984) 1.
- [5] H. E. Haber and G. L. Kane, Phys. Rept. **117** (1985) 75.
- [6] S. Heinemeyer, O. Stål and G. Weiglein, Phys. Lett. B **710** (2012) 201 [arXiv:1112.3026 [hep-ph]].
- [7] P. Bechtle, H. E. Haber, S. Heinemeyer, O. Stål, T. Stefaniak, G. Weiglein and L. Zeune, Eur. Phys. J. C **77** (2017) no.2, 67 [arXiv:1608.00638 [hep-ph]].
- [8] H. Bahl *et al.*, arXiv:1808.07542 [hep-ph].
- [9] U. Ellwanger, C. Hugonie and A. M. Teixeira, Phys. Rept. **496** (2010) 1 [arXiv:0910.1785 [hep-ph]].
- [10] M. Maniatis, Int. J. Mod. Phys. A **25** (2010) 3505 [arXiv:0906.0777 [hep-ph]].
- [11] J. R. Ellis, J. F. Gunion, H. E. Haber, L. Roszkowski and F. Zwirner, Phys. Rev. D **39** (1989) 844.
- [12] D. J. Miller, R. Nevzorov and P. M. Zerwas, Nucl. Phys. B **681** (2004) 3 [hep-ph/0304049].
- [13] S. King, M. Mühlleitner and R. Nevzorov, Nucl. Phys. B **860** (2012) 207 [arXiv:1201.2671 [hep-ph]].
- [14] F. Domingo and G. Weiglein, JHEP **1604** (2016) 095 [arXiv:1509.07283 [hep-ph]].
- [15] D. E. López-Fogliani and C. Muñoz, Phys. Rev. Lett. **97** (2006) 041801 [hep-ph/0508297].
- [16] N. Escudero *et al.*, JHEP **0812** (2008) 099 [arXiv:0810.1507 [hep-ph]].
- [17] C. Muñoz, AIP Conf. Proc. **1200** (2010) 413 [arXiv:0909.5140 [hep-ph]].
- [18] C. Muñoz, PoS DSU **2015** (2016) 034 [arXiv:1608.07912 [hep-ph]].
- [19] P. Ghosh *et al.*, Int. J. Mod. Phys. A **33** (2018) no.18n19, 1850110 [arXiv:1707.02471 [hep-ph]].

- [20] K. Y. Choi, D. E. López-Fogliani, C. Muñoz and R. Ruiz de Austri, JCAP **1003** (2010) 028 [arXiv:0906.3681 [hep-ph]].
- [21] G. A. Gómez-Vargas *et al.*, JCAP **1202** (2012) 001 [arXiv:1110.3305 [astro-ph.HE]].
- [22] A. Albert *et al.* [Fermi-LAT Collab.], JCAP **1410** (2014) no.10, 023 [arXiv:1406.3430 [astro-ph.HE]].
- [23] G. A. Gómez-Vargas *et al.*, JCAP **1703** (2017) no.03, 047 [arXiv:1608.08640 [hep-ph]].
- [24] R. Barate *et al.* [ALEPH and DELPHI and L3 and OPAL Collaborations and LEP Working Group for Higgs boson searches], Phys. Lett. B **565** (2003) 61 [hep-ex/0306033].
- [25] J. Cao *et al.*, Phys. Rev. D **95** (2017) no.11, 116001 [arXiv:1612.08522 [hep-ph]].
- [26] A. Azatov, R. Contino and J. Galloway, JHEP **1204** (2012) 127 Erratum: [JHEP **1304** (2013) 140] [arXiv:1202.3415 [hep-ph]].
- [27] A. M. Sirunyan *et al.* [CMS Collaboration], CMS-PAS-HIG-17-013 [arXiv:1811.08459 [hep-ex]].
- [28] The ATLAS collaboration, ATLAS-CONF-2018-025.
- [29] S. Heinemeyer, Int. J. Mod. Phys. A **33** (2018) no.31, 1844006.
- [30] P. J. Fox and N. Weiner, JHEP **1808** (2018) 025 [arXiv:1710.07649 [hep-ph]].
- [31] F. Richard, arXiv:1712.06410 [hep-ex].
- [32] U. Haisch and A. Malinauskas, JHEP **1803** (2018) 135 [arXiv:1712.06599 [hep-ph]].
- [33] T. Biekötter, S. Heinemeyer and C. Muñoz, Eur. Phys. J. C **78** (2018) no.6, 504 [arXiv:1712.07475 [hep-ph]].
- [34] D. Liu, J. Liu, C. E. M. Wagner and X. P. Wang, JHEP **1806** (2018) 150 [arXiv:1805.01476 [hep-ph]].
- [35] F. Domingo, S. Heinemeyer, S. Paßehr and G. Weiglein, arXiv:1807.06322 [hep-ph].
- [36] W. G. Hollik, S. Liebler, G. Moortgat-Pick, S. Paßehr and G. Weiglein, arXiv:1809.07371 [hep-ph].
- [37] U. Ellwanger, Phys. Lett. B **698** (2011) 293 [arXiv:1012.1201 [hep-ph]].
- [38] R. Benbrik, M. Gomez Bock, S. Heinemeyer, O. Stål, G. Weiglein and L. Zeune, Eur. Phys. J. C **72** (2012) 2171 [arXiv:1207.1096 [hep-ph]].
- [39] P. Bechtle, O. Brein, S. Heinemeyer, G. Weiglein and K. E. Williams, Comput. Phys. Commun. **181** (2010) 138 [arXiv:0811.4169 [hep-ph]].
- [40] P. Bechtle, O. Brein, S. Heinemeyer, G. Weiglein and K. E. Williams, Comput. Phys. Commun. **182** (2011) 2605 [arXiv:1102.1898 [hep-ph]].
- [41] P. Bechtle, O. Brein, S. Heinemeyer, O. Stål, T. Stefaniak, G. Weiglein and K. E. Williams, Eur. Phys. J. C **74** (2014) no.3, 2693 [arXiv:1311.0055 [hep-ph]].
- [42] P. Bechtle, S. Heinemeyer, O. Stål, T. Stefaniak and G. Weiglein, Eur. Phys. J. C **75** (2015) no.9, 421 [arXiv:1507.06706 [hep-ph]].
- [43] <http://higgsbounds.hepforge.org>.
- [44] P. Bechtle, S. Heinemeyer, O. Stål, T. Stefaniak and G. Weiglein, Eur. Phys. J. C **74** (2014) no.2, 2711 [arXiv:1305.1933 [hep-ph]].
- [45] P. Bechtle, S. Heinemeyer, O. Stål, T. Stefaniak and G. Weiglein, JHEP **1411** (2014) 039 [arXiv:1403.1582 [hep-ph]].
- [46] P. Bechtle, D. Derks, S. Heinemeyer, T. Klingl, T. Stefaniak, G. Weiglein, IFT-UAM/CSIC-18-125.
- [47] P. Drechsel, G. Moortgat-Pick and G. Weiglein, arXiv:1801.09662 [hep-ph].