

DEUTSCHES ELEKTRONEN-SYNCHROTRON
Ein Forschungszentrum der Helmholtz-Gemeinschaft



DESY 19-103
KW 19-005
arXiv:1906.08815
June 2019

Electroweak Pseudo-Observables and Z-Boson Form Factors at Two-Loop Accuracy

I. Dubovyk

II. Institut für Theoretische Physik, Universität Hamburg

A. Freitas

*Pittsburgh Particle Physics, Astrophysics and Cosmology Center (PITT PACC),
Department of Physics and Astronomy, University of Pittsburgh, USA*

J. Gluza

Institute of Physics, University of Silesia, Katowice, Poland

and

Faculty of Science, University of Hradec Králové, Czech Republic

T. Riemann

Institute of Physics, University of Silesia, Katowice, Poland

and

Deutsches Elektronen-Synchrotron DESY, Zeuthen

J. Usovitsch

School of Mathematics, Trinity College Dublin, University of Dublin, Ireland

ISSN 0418-9833

PLATANENALLEE 6 - 15738 ZEUTHEN

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your reports and preprints are promptly included in the
HEP literature database
send them to (if possible by air mail):

DESY Zentralbibliothek Notkestraße 85 22607 Hamburg Germany	DESY Bibliothek Platanenallee 6 15738 Zeuthen Germany
---	---

Electroweak pseudo-observables and Z-boson form factors at two-loop accuracy

Ievgen Dubovyk,^a Ayres Freitas,^b Janusz Gluza,^{c,d} Tord Riemann,^{c,e}
Johann Usovitsch^f

^a*II. Institut für Theoretische Physik, Universität Hamburg, 22761 Hamburg, Germany*

^b*Pittsburgh Particle Physics, Astrophysics & Cosmology Center (PITT PACC), Department of Physics & Astronomy, University of Pittsburgh, Pittsburgh, PA 15260, USA*

^c*Institute of Physics, University of Silesia, Katowice, Poland*

^d*Faculty of Science, University of Hradec Králové, Czech Republic*

^e*Deutsches Elektronen-Synchrotron, DESY, 15738 Zeuthen, Germany*

^f*School of Mathematics, Trinity College Dublin, University of Dublin, Dublin 2, Ireland*

E-mail: e.a.dubovyk@gmail.com, afreitas@pitt.edu,
gluza@us.edu.pl, tordriemann@gmail.com, jusovitsch@gmail.com

ABSTRACT: We present Standard Model predictions for the complete set of phenomenologically relevant electroweak precision pseudo-observables related to the Z-boson: the leptonic and bottom-quark effective weak mixing angles $\sin^2 \theta_{\text{eff}}^{\ell}$, $\sin^2 \theta_{\text{eff}}^b$, the Z-boson partial decay widths Γ_f , where f indicates any charged lepton, neutrino and quark flavor (except for the top quark), as well as the total Z decay width Γ_Z , the branching ratios R_{ℓ} , R_c , R_b , and the hadronic cross section σ_{had}^0 . The input parameters are the masses M_Z , M_H and m_t , and the couplings α_s , α . The scheme dependence due to the choice of M_W or its alternative G_{μ} as a last input parameter is also discussed. Recent substantial technical progress in the calculation of Minkowskian massive higher-order Feynman integrals allows the calculation of the complete electroweak two-loop radiative corrections to all the observables mentioned. QCD contributions are included appropriately. Results are provided in terms of simple and convenient parameterization formulae whose coefficients have been determined from the full numerical multi-loop calculation. The size of the missing electroweak three-loop or QCD higher-order corrections is estimated. We briefly comment on the prospects for their calculation. Finally, direct predictions for the $Z\bar{f}f$ vector and axial-vector form-factors are given, including a discussion of separate order-by-order contributions.

Contents

1	Introduction	2
2	Z -boson decay width, branching ratios and cross-sections	4
3	Asymmetries and effective weak mixing angles	7
4	Vector and axial-vector Z -boson form factors F_V^f and F_A^f	11
5	Theoretical error estimates for missing higher order corrections	14
6	Conclusions and outlook	14
	References	17

1 Introduction

In 2018 we celebrated 50 years of the Standard Model of elementary particles. The basics of the model were formulated and experimentally validated in the 1960s/70s. The next decade brought an intensive development of the calculation of quantum field theoretical radiative corrections in that model and in its alternatives. An experimental highlight in this context was the e^+e^- -collider LEP, which enabled us to check the Standard Model at an accuracy of better than the per-cent level, which corresponds to effects from more than one electroweak and two QCD loop orders. This proved, for the first time in a systematic way, the Standard Model as a quantum field theory. LEP 1 was running, from Summer 1989 to 1995, at and around the Z -boson peak. The expectation for the experimental precision of M_Z and Γ_Z was 20 MeV in 1986 [1] and reached finally 2 MeV [2]. This precision tag was extremely important because M_Z is one of the Standard Model input parameters to the commonly used on-mass-shell renormalization scheme. Indeed, the experimental accuracy of M_Z triggered much of the precision loop calculations, including the prediction of the top quark and Higgs masses prior to their discoveries from loop corrections to LEP observables in the Standard Model, see Refs. [3–5] (as well as Refs. [6, 7] for an overview of the current state of the art). Data from the Z peak and the Z resonance curve (the Z line shape) allow to measure a large variety of observables, such as M_Z , Γ_Z , cross-sections for different two-fermion final states and their ratios and angular asymmetries, together with radiation of (sufficiently soft) photons, gluons, etc. From the real observables, the so-called electroweak pseudo-observables (EWPOs) are extracted by means of a de-convolution of initial-state radiation and subtraction of backgrounds. The fine details of relating EWPOs to real cross-sections at LEP 1 precision are described in detail in Ref. [8] and references quoted therein.

On occasion of the 50th anniversary of the Standard Model, many of the related crucial developments were remembered at the conference "SM@50" [9].

This article is devoted to the state-of-the art calculation of the Standard Model (SM) corrections to the $Z\bar{f}f$ -vertex and their inclusion into the predictions for the various EWPOs. We will mostly focus on recent advances in the calculation of the electroweak two-loop terms. QCD contributions, which are known up to four-loop order, have also been taken into account in the results presented here, but we will refer to the literature for further details.

The Z resonance curve can be described theoretically by writing the S-matrix elements as a Laurent series in the center-of-mass energy squared s (also called S-matrix ansatz). This Laurent series ansatz is worked out up to two loops [10–18]. The coefficients of the leading series term contain the Z vertex form factors. Their one-loop corrections were studied in the 1980s; first with massless fermions [19–22], and slightly later with the full mass dependence of the Standard Model [23–26].

After several papers on approximate/partial higher-order corrections, the complete two-loop weak corrections were determined in a series of papers from 2004 to 2018 [27–37]. The correct formulation of the interplay of the 2→2 loop corrections with higher order real QED corrections in the S-matrix approach, also called un-folding of the effective 2→2 Born terms from the realistic 2→n observables, is a topic on its own. It was first studied in Refs. [38–41], but its extension beyond two-loop level will require more work [8, 42, 43]. The numerically relevant two-loop and partial higher-order corrections were included in the final analysis of LEP 1 data [44].¹

The theoretical advances described here go beyond the Standard Model theory used for physics at LEP 1 [45, 46] but will be needed for the FCC-ee Tera-Z project [6, 7, 50–52] whose unique experimental precision calls for perturbative predictions at three electroweak loops together with corresponding QCD terms.

In this work, the following pseudo-EWPOs are discussed: The partial widths Γ_f for Z -boson decay into $f\bar{f}$ final states; the total Z width Γ_Z ; the branching fractions Γ_f/Γ_Z ; the total hadronic Z -pole quark-pair production cross-section σ_{had}^0 ; and the effective weak mixing angles $\sin^2\theta_{\text{eff}}^f$, defined from the ratio of vector- and axial Z -boson couplings. The precise definition of all these pseudo-observables will be given below. Pseudo-observables differ from real observables by removing from the former the effects of initial-state and initial-final QED radiation, as well as non-resonant photon-exchange, box and t -channel contributions [6, 44].

The Standard Model predictions for Z -pole pseudo-observables can be constructed in terms of the following three theoretical building blocks [29]:

$$v_f(s) \equiv v_f^Z(s) - v_f^\gamma(s) \frac{\Sigma_{\gamma Z}(s)}{s + \Sigma_{\gamma\gamma}(s)}, \quad (1.1)$$

$$a_f(s) \equiv a_f^Z(s) - a_f^\gamma(s) \frac{\Sigma_{\gamma Z}(s)}{s + \Sigma_{\gamma\gamma}(s)}, \quad (1.2)$$

$$\Sigma_Z(s) \equiv \Sigma_{ZZ}(s) - \frac{[\Sigma_{\gamma Z}(s)]^2}{s + \Sigma_{\gamma\gamma}(s)}, \quad (1.3)$$

where v_f^Z and a_f^Z are the one-particle irreducible $Zf\bar{f}$ vector- and axial-vector vertex contributions, respectively, whereas v_f^γ and a_f^γ are their counterparts for the $\gamma f\bar{f}$ vertex. The $\Sigma_{V_1 V_2}$ denotes the one-particle irreducible V_1 – V_2 self-energy. At tree level,

$$v_{f(0)}^Z = e \frac{I_f^3 - 2Q_f s_w^2}{2s_w c_w}, \quad v_{f(0)}^\gamma = eQ_f, \quad (1.4)$$

¹The EWPOs at LEP 1 were determined order by order without a Laurent expansion. This was based on the ZFITTER software [45, 46], for both the Standard Model loop calculation and the unfolding of cross-sections. The relevant higher-order corrections to the input W mass [47] and to the leptonic weak mixing angle [27] are implemented in ZFITTER v.6.42. While ZFITTER v.6.44beta [48] also contains the QCD four-loop corrections of [49], they are of no experimental relevance so far.

$$a_{f(0)}^Z = e \frac{I_f^3}{2s_w c_w}, \quad a_{f(0)}^\gamma = 0. \quad (1.5)$$

Here I_f^3 and Q_f are the weak isospin and electric charge (in units of the elementary charge $e > 0$) of the fermion f , respectively. s_w and c_w are the sine and cosine of the weak mixing angle, respectively, and the subscript (0) is used to denote tree-level order.

For the theory calculations, these building blocks must be evaluated at the complex Z pole [11–13, 17], $s_0 \equiv \overline{M}_Z^2 - i\overline{M}_Z\overline{\Gamma}_Z$, where \overline{M}_Z and $\overline{\Gamma}_Z$ are the on-shell mass and width of the Z -boson, respectively. Note that \overline{M}_Z and $\overline{\Gamma}_Z$ differ from the mass M_Z and width Γ_Z reported in publications of LEP, Tevatron and LHC experiments by a fixed factor [38, 53]:

$$\begin{aligned} \overline{M}_Z &= M_Z / \sqrt{1 + \Gamma_Z^2 / M_Z^2}, \\ \overline{\Gamma}_Z &= \Gamma_Z / \sqrt{1 + \Gamma_Z^2 / M_Z^2}. \end{aligned} \quad (1.6)$$

Similar expressions hold for \overline{M}_W and $\overline{\Gamma}_W$ [54, 55].

2 Z -boson decay width, branching ratios and cross-sections

The width of the Z boson, $\overline{\Gamma}_Z$, is related to the imaginary part of the Z self-energy. Using the optical theorem, one can derive the following expression for $\overline{\Gamma}_Z$ [33, 35]:

$$\overline{\Gamma}_Z = \sum_f \overline{\Gamma}_f, \quad f = e, \mu, \tau, \nu_e, \nu_\mu, \nu_\tau, u, c, s, d, b, \quad (2.1)$$

$$\overline{\Gamma}_f = \frac{N_c^f \overline{M}_Z}{12\pi} \left[\mathcal{R}_V^f F_V^f + \mathcal{R}_A^f F_A^f \right]_{s=\overline{M}_Z^2}. \quad (2.2)$$

Here N_c^f is the color factor and $\mathcal{R}_{V,A}$ are radiator functions that capture final-state QCD and QED corrections, see section 7 in Ref. [56], whereas the remaining electroweak and mixed electroweak–QCD corrections are contained in the form factors $F_{V,A}^f$. Up to two-loop accuracy, the form factors can be written as follows [35]:

$$F_V^f = \frac{|v_f|^2}{1 + \text{Re}\{\Sigma_Z' - \frac{i}{2}\overline{M}_Z\overline{\Gamma}_Z\Sigma_Z''\}} \Big|_{s=\overline{M}_Z^2}, \quad (2.3)$$

$$F_A^f = \frac{|a_f|^2}{1 + \text{Re}\{\Sigma_Z' - \frac{i}{2}\overline{M}_Z\overline{\Gamma}_Z\Sigma_Z''\}} \Big|_{s=\overline{M}_Z^2}, \quad (2.4)$$

where Σ_Z' and Σ_Z'' are shorthand expressions for $d\Sigma_Z/ds$ and $d^2\Sigma_Z/ds^2$, respectively.

In addition to the partial widths, certain branching ratios are of phenomenological importance:

$$R_\ell = \overline{\Gamma}_{\text{had}} / \overline{\Gamma}_\ell, \quad R_c = \overline{\Gamma}_c / \overline{\Gamma}_{\text{had}}, \quad R_b = \overline{\Gamma}_b / \overline{\Gamma}_{\text{had}}. \quad (2.5)$$

Here $\bar{\Gamma}_{\text{had}} = \sum_{f=u,c,d,s,b} \bar{\Gamma}_f$. Further, the cross-section for $e^+e^- \rightarrow \text{hadrons}$ at the Z peak can be expressed in terms of partial widths [35],

$$\sigma_{\text{had}} = \sigma_{\text{had}}^0 + \sigma_{\text{had,non-res}}, \quad \sigma_{\text{had}}^0 = \sum_{f=u,c,d,s,b} \frac{12\pi}{M_Z^2} \frac{\bar{\Gamma}_e \bar{\Gamma}_f}{\bar{\Gamma}_Z^2} (1 + \delta X). \quad (2.6)$$

Here $\sigma_{\text{had,non-res}}^0$ accounts for non-resonant photon-exchange, box and t -channel contributions. Furthermore, δX occurs from higher-order terms of the Laurent expansion of the full amplitude around the complex pole s_0 . At two-loop order, δX can be written as $\delta X_{(2)} = -(\text{Im} \Sigma'_{Z(1)})^2 - 2\bar{\Gamma}_Z \bar{M}_Z \text{Im} \Sigma''_{Z(1)}$, where subscripts (n) indicate the loop order. In the limit $m_f \ll M_W$ ($f \neq t$), it is given by

$$\delta X_{(2)} = - \left[\frac{\alpha}{s_W^2 c_W^2} \left(\frac{7}{8} - \frac{5}{3} s_W^2 + \frac{20}{9} s_W^4 \right) \right]^2. \quad (2.7)$$

It is important to note that eq. (2.6) assumes that initial-state photon radiation effects have been removed by means of a de-convolution procedure, see e.g. Ref. [57].

Results for the partial and total Z widths, branching ratios and σ_{had}^0 including the full two-loop corrections have first been published in Ref. [37]. They can be expressed in simple parameterization formulae, which are adequate for most phenomenological applications. Here, we present slightly more complicated formulae that cover an extended numerical range of input parameters:²

$$\begin{aligned} 25 \text{ GeV} < M_H < 225 \text{ GeV}, & & 155 \text{ GeV} < m_t < 192 \text{ GeV}, \\ M_Z &= 91.1876 \pm 0.0084 \text{ GeV}, \\ \alpha_s &= 0.1184 \pm 0.0050, & \Delta\alpha &= 0.0590 \pm 0.0005. \end{aligned} \quad (2.8)$$

Here $\Delta\alpha$ is the shift in the running electromagnetic coupling $\alpha(q^2)$ from $q^2 = 0$ to M_Z^2 , defined by $\alpha(M_Z^2) = \alpha(0)/(1 - \Delta\alpha)$. It can be divided into a leptonic and a hadronic part, $\Delta\alpha = \Delta\alpha_{\text{lept}} + \Delta\alpha_{\text{had}}$. $\Delta\alpha_{\text{lept}}$ has been computed to three-loop order [59], whereas $\Delta\alpha_{\text{had}}$ contains non-perturbative hadronic contributions, which are commonly extracted from data [60–62]. We neglect the light fermion masses m_f , $f \neq t$, everywhere besides in $\Delta\alpha$ and (at leading power) in the radiator functions $\mathcal{R}_{V/A}^f$. The W boson mass M_W can be computed from the Fermi constant G_μ (see eqs. (6)–(8) in the arXiv version of Ref. [47]) and thus is not listed as an independent input parameter. Both G_μ and α , the electromagnetic fine structure constant in the Thomson limit, are known with very small uncertainties, and thus we use their central experimental values [58] without any uncertainty interval.

²This extended input parameter range is useful for determining indirect constraints on various SM parameters from electroweak precision observables (see e.g. section 10 in Ref. [58]), since these indirect bounds often extend over larger intervals than the corresponding direct measurements.

Observable	X_0	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8
$\Gamma_{e,\mu}$ [MeV]	83.983	-0.1202	-0.06919	0.00383	0.0597	0.8037	-0.015	-0.0195	0.0032
Γ_τ [MeV]	83.793	-0.1200	-0.06905	0.00382	0.0596	0.8023	-0.015	-0.0195	0.0032
Γ_ν [MeV]	167.176	-0.1752	-0.1249	0.00595	0.1046	1.253	-0.110	-0.0232	0.0064
Γ_u [MeV]	299.994	-0.6152	-0.2771	0.0174	0.2341	4.051	-0.467	-0.0676	0.017
Γ_c [MeV]	299.918	-0.6152	-0.2771	0.0174	0.2340	4.051	-0.467	-0.0676	0.017
$\Gamma_{d,s}$ [MeV]	382.829	-0.6685	-0.3322	0.0193	0.2792	3.792	-0.18	-0.0706	0.020
Γ_b [MeV]	375.890	-0.6017	-0.3158	0.0190	0.227	-2.174	0.042	-0.027	0.021
Γ_Z [MeV]	2494.75	-4.055	-2.117	0.122	1.746	19.68	-1.63	-0.432	0.12
R_ℓ [10^{-3}]	20751.6	-8.112	-1.174	0.155	0.16	-37.59	-10.9	1.27	0.29
R_c [10^{-5}]	17222.2	-4.049	-0.749	0.0832	1.08	98.956	-15.1	-0.761	0.080
R_b [10^{-5}]	21585.0	4.904	0.9149	-0.0535	-2.676	-292.21	20.0	1.97	-0.11
σ_{had}^0 [pb]	41489.6	0.408	-0.320	0.0424	1.32	60.17	16.3	-2.31	-0.19

Observable	a_9	a_{10}	a_{11}	a_{12}	a_{13}	a_{14}	a_{15}	a_{16}	max. dev.
$\Gamma_{e,\mu}$ [MeV]	-0.0956	-0.0078	-0.0095	0.25	-1.08	0.056	-0.37	286	< 0.0015
Γ_τ [MeV]	-0.0954	-0.0078	-0.0094	0.25	-1.08	0.056	-0.37	285	< 0.0015
Γ_ν [MeV]	-0.187	-0.014	-0.014	0.37	-0.085	0.054	-0.30	503	< 0.002
Γ_u [MeV]	14.26	1.6	-0.046	1.82	-11.1	0.16	-1.0	1253	< 0.006
Γ_c [MeV]	14.26	1.6	-0.046	1.82	-11.1	0.16	-1.0	1252	< 0.006
$\Gamma_{d,s}$ [MeV]	10.20	-2.4	-0.052	0.71	-10.1	0.16	-0.92	1469	< 0.007
Γ_b [MeV]	10.53	-2.4	-0.056	1.2	-10.1	0.15	-0.95	1458	< 0.007
Γ_Z [MeV]	58.61	-4.0	-0.32	8.1	-56.1	1.1	-6.8	9267	< 0.04
R_ℓ [10^{-3}]	732.30	-44	-0.61	5.7	-358	-4.7	37	11649	< 0.12
R_c [10^{-5}]	230.9	125	0.045	36.9	-120	1.2	-6.2	3667	< 0.1
R_b [10^{-5}]	-131.9	-84	-0.27	4.4	71.9	-0.77	-4.4	-1790	< 0.12
σ_{had}^0 [pb]	-579.58	38	0.010	7.5	85.2	9.1	-68	-85957	< 0.15

Table 1. Coefficients for the parameterization formula (2.9) for various observables. Within the ranges $25 \text{ GeV} < M_H < 225 \text{ GeV}$, $155 \text{ GeV} < m_t < 195 \text{ GeV}$, $\alpha_s = 0.1184 \pm 0.0050$, $\Delta\alpha = 0.0590 \pm 0.0005$ and $M_Z = 91.1876 \pm 0.0084 \text{ GeV}$, the formulae approximate the full result with maximal deviations given in the last column.

The fitting formulae for the EWPOs have the form

$$\begin{aligned}
X = & X_0 + a_1 L_H + a_2 L_H^2 + a_3 L_H^4 + a_4 \Delta_H + a_5 \Delta_t + a_6 \Delta_t^2 + a_7 \Delta_t L_H + a_8 \Delta_t L_H^2 \\
& + a_9 \Delta_{\alpha_s} + a_{10} \Delta_{\alpha_s}^2 + a_{11} \Delta_{\alpha_s} \Delta_H + a_{12} \Delta_{\alpha_s} \Delta_t + a_{13} \Delta_\alpha + a_{14} \Delta_\alpha \Delta_H \\
& + a_{15} \Delta_\alpha \Delta_t + a_{16} \Delta_Z,
\end{aligned} \tag{2.9}$$

$$\begin{aligned}
L_H = & \log \frac{M_H}{125.7 \text{ GeV}}, \quad \Delta_H = \frac{M_H}{125.7 \text{ GeV}} - 1, \quad \Delta_t = \left(\frac{m_t}{173.2 \text{ GeV}} \right)^2 - 1, \\
\Delta_{\alpha_s} = & \frac{\alpha_s(M_Z)}{0.1184} - 1, \quad \Delta_\alpha = \frac{\Delta\alpha}{0.059} - 1, \quad \Delta_Z = \frac{M_Z}{91.1876 \text{ GeV}} - 1.
\end{aligned}$$

The coefficients X_0 and a_1, \dots, a_{16} are obtained from fits to a grid of 8750 data points of the full computation. The latter includes

- Complete one-loop corrections [23], which have been re-computed for this work, and full two-loop [33, 35, 37] electroweak corrections;
- Corrections of order $\mathcal{O}(\alpha\alpha_s)$ to vector-boson self-energies [63–67], which have been re-evaluated for this work;
- Non-factorizable $\mathcal{O}(\alpha\alpha_s)$ $Zq\bar{q}$ vertex contributions [68–73], which are not captured in the products $\mathcal{R}_i^f F_i^f$ ($i = V, A$);
- Higher-loop QCD corrections in the limit of a large top Yukawa coupling y_t , of orders $\mathcal{O}(\alpha_t\alpha_s^2)$ [74, 75], $\mathcal{O}(\alpha_t^2\alpha_s)$, $\mathcal{O}(\alpha_t^3)$ [76, 77], and $\mathcal{O}(\alpha_t\alpha_s^3)$ [78–80], where $\alpha_t \equiv y_t^2/(4\pi)$.
- Final-state QED and QCD radiation effects, which enter through the radiator functions $\mathcal{R}_{V,A}$, up to the orders $\mathcal{O}(\alpha^2)$, $\mathcal{O}(\alpha\alpha_s)$ and $\mathcal{O}(\alpha_s^4)$ [56, 81, 82].

Both the Z vertex corrections as well as the prediction of M_W from G_μ have been computed to this same level of perturbation theory.

Numerical values for the coefficients are given in Tab. 1. Some of the numbers for X_0 deviate slightly in the last digit from those in Ref. [37]. This is due to the larger grid of input parameters used here, which can exert a pull on the fit parameters. The differences are well within the accuracy quoted in the last column of Tab. 1.

3 Asymmetries and effective weak mixing angles

The effective weak mixing angle for the $Zf\bar{f}$ vertex is defined, from the theory side, as

$$\sin^2 \theta_{\text{eff}}^f \equiv \frac{1}{4|Q_f|} \left(1 - \text{Re} \left\{ \frac{v_f(\overline{M_Z^2})}{a_f(\overline{M_Z^2})} \right\} \right). \tag{3.1}$$

Here \overline{M}_W^2 and \overline{M}_Z^2 are the real parts of the complex pole of the W and Z propagators, respectively. They are related to the masses commonly reported by experiments at LEP, Tevatron, LHC according to eq (1.6). Moreover, Q_f denotes the electric charge of the fermion f .

The effective weak mixing angles can be extracted from a range of asymmetries [8], defined from effective Born two-particle cross-sections, including the left-right asymmetry

$$A_{\text{LR}} = \frac{\sigma_{e_L} - \sigma_{e_R}}{\sigma_{e_L} + \sigma_{e_R}} = A_e + A_{\text{LR}}^{\text{non-res}} \quad (3.2)$$

and the forward-backward asymmetry

$$A_{\text{FB}}^f = \frac{\sigma_{\cos\theta>0} - \sigma_{\cos\theta<0}}{\sigma_{\cos\theta>0} + \sigma_{\cos\theta<0}} = \frac{3}{4}A_e A_f + A_{\text{FB}}^{f,\text{non-res}}, \quad (3.3)$$

where

$$A_f \equiv \frac{2 \operatorname{Re}\{v_f/a_f\}}{1 + (\operatorname{Re}\{v_f/a_f\})^2} = \frac{1 - 4|Q_f| \sin^2 \theta_{\text{eff}}^f}{1 - 4|Q_f| \sin^2 \theta_{\text{eff}}^f + 8(|Q_f| \sin^2 \theta_{\text{eff}}^f)^2}. \quad (3.4)$$

Here σ_{e_L} and σ_{e_R} are the cross-sections for $e^+e^- \rightarrow f\bar{f}$ for left- and right-handed polarized electron beams, respectively, whereas $\sigma_{\cos\theta>0}$ and $\sigma_{\cos\theta<0}$ denote the cross-section for f restricted to the forward and backward hemisphere, respectively. Furthermore, $A_X^{\text{non-res}}$ accounts for the non-resonant photon-exchange, box and t -channel contributions.

The most precisely measured effective weak mixing angles are the leptonic effective weak mixing angle $\sin^2 \theta_{\text{eff}}^\ell$ (extracted from A_{LR}) and the bottom-quark one, $\sin^2 \theta_{\text{eff}}^b$ (extracted from A_{FB}^b) [57].

Standard Model predictions for $\sin^2 \theta_{\text{eff}}^\ell$ including the full two-loop corrections have been presented originally in Ref. [28, 29, 31]. We reproduced by an independent calculation the contribution of the bosonic electroweak two-loop corrections using the methods of Ref. [37]. The corrections can be expressed in terms of a weak form factor $\Delta\kappa_\ell^{(\alpha^2, \text{bos})}$, where

$$\Delta\kappa_f = \left(1 - \frac{\overline{M}_W}{\overline{M}_Z}\right)^{-1} \sin^2 \theta_{\text{eff}}^f - 1. \quad (3.5)$$

The comparison with Ref. [28] is shown in Tab. 2, which demonstrates that the two calculations agree to an accuracy of $\mathcal{O}(10^{-7})$, which implies an accuracy of better than 10^{-7} for $\sin^2 \theta_{\text{eff}}^\ell$. The full two-loop corrections for $\sin^2 \theta_{\text{eff}}^b$ have been presented first in Ref. [36].

M_H [GeV]	Result of Ref. [28]	Our result
100	-0.733989×10^{-4}	-0.733955×10^{-4}
200	-0.469470×10^{-4}	-0.471273×10^{-4}

Table 2. Comparison of numerical results for $\Delta\kappa_\ell^{(\alpha^2, \text{bos})}$ from Ref. [28] with our calculation of Z vertex corrections from Ref. [37]. We use $\overline{M}_Z = 91.1876$ GeV and $\overline{M}_W = 80.385$ GeV.

In the following, we present simple parameterization formulae for $\sin^2\theta_{\text{eff}}^\ell$ and $\sin^2\theta_{\text{eff}}^b$, which cover the extended range of input parameters of eq. (2.8). The parameterization formula

$$\begin{aligned} \sin^2\theta_{\text{eff}}^f = & s_0 + d_1 L_H + d_2 L_H^2 + d_3 L_H^4 + d_4 \Delta_\alpha + d_5 \Delta_t + d_6 \Delta_t^2 + d_7 \Delta_t L_H \\ & + d_8 \Delta_{\alpha_s} + d_9 \Delta_{\alpha_s} \Delta_t + d_{10} \Delta_Z \end{aligned} \quad (3.6)$$

with

$$\begin{aligned} L_H = & \log \frac{M_H}{125.7 \text{ GeV}}, \quad \Delta_t = \left(\frac{m_t}{173.2 \text{ GeV}} \right)^2 - 1, \\ \Delta_{\alpha_s} = & \frac{\alpha_s(M_Z)}{0.1184} - 1, \quad \Delta_\alpha = \frac{\Delta\alpha}{0.059} - 1, \quad \Delta_Z = \frac{M_Z}{91.1876 \text{ GeV}} - 1 \end{aligned}$$

provides a good description of the full result in the parameter region (2.8). Values for the coefficients are obtained by fitting (3.6) to a grid of 8750 data points.

Table 3 shows the result of a fit to a calculation that includes all known corrections:

- Complete one- and two-loop electroweak corrections, (see Refs. [21, 23, 27, 28, 30–32, 36] for the original references);
- Corrections of order $\mathcal{O}(\alpha\alpha_s)$ to vector-boson self-energies [63–67], which we have re-evaluated for this work;
- Non-factorizable $\mathcal{O}(\alpha\alpha_s)$ $Zb\bar{b}$ vertex contributions [68–73], which do not cancel in the ratio v_b/a_b ;
- Higher-loop corrections in the limit of a large top Yukawa coupling y_t , of orders $\mathcal{O}(\alpha_t\alpha_s^2)$ [74, 75], $\mathcal{O}(\alpha_t^2\alpha_s)$, $\mathcal{O}(\alpha_t^3)$ [76, 77], and $\mathcal{O}(\alpha_t\alpha_s^3)$ [78–80] where $\alpha_t \equiv y_t^2/(4\pi)$.

As indicated by the last column in the table, the largest deviation of the fit formulae from the full result is $\mathcal{O}(\text{few} \times 10^{-6})$, while for most of the parameter region in (2.8) the agreement is better than 10^{-6} . The careful reader may realize that the parameterization for $\sin^2\theta_{\text{eff}}^b$ in Table 3 deviates slightly from Eqs. (20,22) in [36]. The difference is due to the larger grid of data points used here. A fit formula is,

Observable	s_0	d_1	d_2	d_3	d_4	d_5
$\sin^2 \theta_{\text{eff}}^\ell \times 10^4$	2314.64	4.616	0.539	-0.0737	206	-25.71
$\sin^2 \theta_{\text{eff}}^b \times 10^4$	2327.04	4.638	0.558	-0.0700	207	-9.554

Observable	d_6	d_7	d_8	d_9	d_{10}	max. dev.
$\sin^2 \theta_{\text{eff}}^\ell \times 10^4$	4.00	0.288	3.88	-6.49	-6560	< 0.056
$\sin^2 \theta_{\text{eff}}^b \times 10^4$	3.83	0.179	2.41	-8.24	-6630	< 0.025

Table 3. Coefficients for the parameterization formula (3.6) for the leptonic and bottom-quark effective weak mixing angles. Within the ranges given in eq. (2.8), the formula deviates from the full result up to the maximal amount given in the last column.

Observable	max. dev.	EXP now	FCC-ee	CEPC	GigaZ
Γ_Z [MeV]	0.04	2.3	0.1	0.5	0.8
$\sin^2 \theta_{\text{eff}}^\ell \times 10^4$	0.056	1.6	0.06	0.23	0.1
$\sin^2 \theta_{\text{eff}}^b \times 10^4$	0.025	160	9	9	15

Table 4. Goodness of fit for some chosen EWPOs, compared with the envisaged precision measurements for Γ_Z and $\sin^2 \theta_{\text{eff}}^\ell$ (statistical errors), and $\sin^2 \theta_{\text{eff}}^b$ (systematic errors) at the collider projects FCC-ee Tera-Z [83], CEPC [84] and ILC/GigaZ [85]. The values of maximal deviations are taken from Tabs. 1 and 3. The entry “EXP now” gives the present experimental precision, as known since LEP 1 [44].

obviously, not able to reproduce the data points in a grid perfectly. The fitting aims to find the best average agreement between the data points (which are generated with our full numerical calculation) and the fit formula. A larger grid therefore can lead to some shifts of the coefficients. As a consequence, the formula in [36] will probably be more accurate for input values within the ranges in Tab. 1 there. On the other hand, while the formula here may be a little less accurate within these ranges, it covers a much larger range of input values.

It should also be noted that the fit formula for $\sin^2 \theta_{\text{eff}}^\ell$ in Ref. [28] does not include the $\mathcal{O}(\alpha_t \alpha_s^3)$ corrections from Refs. [78–80], but they are included in the formula presented here.

In Tab. 4 it is shown that the technical accuracy of our fit formulae is adequate for the expected experimental precision of several future e^+e^- colliders, although it will get modified by anticipated future three-loop electroweak corrections.

4 Vector and axial-vector Z -boson form factors F_V^f and F_A^f

The pseudo-observables discussed in the previous sections aim to be closely related to actual observables, such as cross-sections, branching ratios, or asymmetries. On the other hand, for some purposes it is also useful to have numerical results for the underlying vertex corrections themselves [34], for example: (i) Inclusion of selected corrections from Beyond Standard Model (BSM) physics, (ii) Estimations of magnitudes of selected single terms, (iii) Partial cross-checks with other calculations. For such purposes, the form factors F_V^f and F_A^f introduced in eq. (2.2) are needed explicitly.

Tables 5 and 6 show the numerical contributions of different orders of perturbation theory to F_V^f and F_A^f . Here the form factors are always understood to include the appropriate (on-shell) counterterms to render them UV-finite. The non-factorizable $\mathcal{O}(\alpha\alpha_s)$ corrections are shown separately in the table. Strictly speaking they are not part of the form factors F_V^f and F_A^f , but they finally contribute to observables, such as the decay width, and we incorporate them in the tables and the fit formula below. In Tab. 5 the form factors are computed using the following input values:

$$M_Z = 91.1876 \text{ GeV}, \quad \Gamma_Z = 2.4952 \text{ GeV}, \quad \Rightarrow \quad \overline{M}_Z = 91.1535 \text{ GeV} \quad (4.1a)$$

$$M_W = 80.385 \text{ GeV}, \quad \Gamma_W = 2.085 \text{ GeV}, \quad \Rightarrow \quad \overline{M}_W = 80.358 \text{ GeV} \quad (4.1b)$$

$$\begin{aligned} M_H &= 125.1 \text{ GeV}, & m_t &= 173.2 \text{ GeV}, \\ m_b^{\overline{\text{MS}}} &= 4.2 \text{ GeV}, & \Delta\alpha &= 0.059, & \alpha_s &= 0.1184 \end{aligned} \quad (4.1c)$$

For Tab. 6, on the other hand, the Fermi constant G_μ is used as an input instead of (4.1b), and \overline{M}_W is computed from

$$\overline{M}_W^2 \left(1 - \frac{\overline{M}_W^2}{\overline{M}_Z^2} \right) = \frac{\pi\alpha}{\sqrt{2}G_\mu} (1 + \Delta r), \quad (4.2)$$

where Δr has been evaluated to the same orders as given in each column of the table. More details about the calculation of Δr can be found in Ref. [47]. As before, the dependence of the Standard Model prediction on various input parameters can be expressed in terms of the simple parameterization formula eq. (2.9).

Table 7 shows the numerical values for the coefficients obtained by fitting this formula to the currently most precise computation, including the same corrections as in section 2, except for the final-state QED and QCD radiation effects, i.e.

$$\begin{aligned} F_V^f &= |v_{f(0)}|^2 + F_{V(\alpha)}^f + F_{V(\alpha\alpha_s)}^f + F_{V(\alpha^2)}^f + F_{V(\alpha_t\alpha_s^2)}^f + F_{V(\alpha_t^2\alpha_s)}^f + F_{V(\alpha_t^3)}^f + F_{V(\alpha_t\alpha_s^3)}^f, \\ F_A^f &= |a_{f(0)}|^2 + F_{A(\alpha)}^f + F_{A(\alpha\alpha_s)}^f + F_{A(\alpha^2)}^f + F_{A(\alpha_t\alpha_s^2)}^f + F_{A(\alpha_t^2\alpha_s)}^f + F_{A(\alpha_t^3)}^f + F_{A(\alpha_t\alpha_s^3)}^f. \end{aligned} \quad (4.3)$$

Note that G_μ (rather than M_W) has been used as one input in Tab. 7.

Form fact.	Born	$\mathcal{O}(\alpha)$	$\mathcal{O}(\alpha\alpha_s)$	$\mathcal{O}(\alpha\alpha_s)$ non-fact.	$\mathcal{O}(\alpha_t\alpha_s^2, \alpha_t\alpha_s^3,$ $\alpha_t^2\alpha_s, \alpha_t^3)$	$\mathcal{O}(N_f^2\alpha^2)$	$\mathcal{O}(N_f\alpha^2)$	$\mathcal{O}(\alpha_{\text{bos}}^2)$
$F_V^\ell [10^{-5}]$	39.07	-24.86	2.41	-	0.35	1.47	2.37	0.27
$F_A^\ell [10^{-5}]$	3309.44	118.59	9.46	-	1.22	8.60	2.60	0.45
$F_{V,A}^\nu [10^{-5}]$	3309.44	127.56	9.46	-	1.22	8.60	3.83	0.39
$F_V^{u,c} [10^{-5}]$	544.88	-44.80	7.29	-0.39	1.02	-1.67	3.25	0.33
$F_A^{u,c} [10^{-5}]$	3309.44	120.79	9.46	-0.98	1.22	8.60	3.27	0.44
$F_V^{d,s} [10^{-5}]$	1635.01	5.84	9.64	-0.80	1.32	0.71	3.45	0.37
$F_A^{d,s} [10^{-5}]$	3309.44	123.78	9.46	-1.14	1.22	8.60	3.11	0.42
$F_V^b [10^{-5}]$	1635.01	-26.16	9.64	3.13	1.32	0.71	1.77	1.05
$F_A^b [10^{-5}]$	3309.44	78.26	9.46	4.45	1.22	8.60	0.13	1.18

Table 5. Contributions of different perturbative orders to the Z vertex form factors. A fixed value of M_W has been used as input, instead of G_μ . N_f^n refers to corrections with n closed fermions loops, whereas α_{bos}^2 denotes corrections without closed fermions loops. Furthermore, $\alpha_t = y_t/(4\pi)$ where y_t is the top Yukawa coupling.

Form fact.	Born	$\mathcal{O}(\alpha)$	$\mathcal{O}(\alpha\alpha_s)$	$\mathcal{O}(\alpha\alpha_s)$ non-fact.	$\mathcal{O}(\alpha_t\alpha_s^2, \alpha_t\alpha_s^3,$ $\alpha_t^2\alpha_s, \alpha_t^3)$	$\mathcal{O}(N_f^2\alpha^2, N_f\alpha^2)$	$\mathcal{O}(\alpha_{\text{bos}}^2)$
$F_V^\ell [10^{-5}]$	77.63	-59.86	0.31	-	-0.09	1.88	0.24
$F_A^\ell [10^{-5}]$	3426.43	19.32	-1.12	-	-0.92	1.62	0.27
$F_{V,A}^\nu [10^{-5}]$	3426.43	28.36	-1.16	-	-0.93	2.81	0.22
$F_V^{u,c} [10^{-5}]$	644.45	-129.87	-1.36	-0.40	-0.73	-6.25	0.19
$F_A^{u,c} [10^{-5}]$	3426.43	21.54	-1.13	-0.99	-0.92	2.28	0.27
$F_V^{d,s} [10^{-5}]$	1760.71	-100.64	-1.85	-0.81	-1.01	-6.23	0.19
$F_A^{d,s} [10^{-5}]$	3426.43	24.56	-1.15	-1.15	-0.93	2.11	0.25
$F_V^b [10^{-5}]$	1760.71	-133.08	-1.58	3.15	-0.96	-7.70	0.86
$F_A^b [10^{-5}]$	3426.43	-21.45	-0.85	4.47	-0.88	-0.64	1.01

Table 6. Same as Tab. 5, but with M_W calculated from G_μ .

The form factor results presented here can be easily augmented to include the effects of some new physics model:

$$F_{V,\text{SM+NP}}^f \approx F_{V,\text{SM}}^f + 2\text{Re}\{v_{f(0)}v_{f,\text{NP}}\}, \quad (4.4)$$

$$F_{A,\text{SM+NP}}^f \approx F_{A,\text{SM}}^f + 2\text{Re}\{a_{f(0)}a_{f,\text{NP}}\}. \quad (4.5)$$

Here ‘‘SM’’ denotes the SM contributions discussed in the present paper, while ‘‘NP’’ stands for the new physics correction on top of the SM. Since the existing experimental constraints imply that any possible new physics effect is small, it is sufficient to use the tree-level couplings $v_{f(0)}$ and $a_{f(0)}$ in the interference terms and neglect the $|v_{f,\text{NP}}|^2$ and $|a_{f,\text{NP}}|^2$ terms.

Form fact.	X_0	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8
$F_V^\ell [10^{-5}]$	20.11	-1.317	-0.2614	0.0333	0.276	7.472	1.55	-0.326	0.0012
$F_A^\ell [10^{-5}]$	3445.60	-3.640	-2.592	0.124	2.186	25.67	-2.17	-0.480	0.13
$F_{V,A}^\nu [10^{-5}]$	3455.73	-3.618	-2.580	0.123	2.161	25.89	-2.27	-0.480	0.13
$F_V^{u,c} [10^{-5}]$	506.025	-4.483	-1.062	0.105	0.912	23.20	-1.54	-0.409	0.085
$F_A^{u,c} [10^{-5}]$	3447.47	-3.630	-2.588	0.124	2.178	25.80	-2.24	-0.480	0.13
$F_V^{d,s} [10^{-5}]$	1650.36	-5.243	-1.834	0.134	1.540	29.20	-2.78	-0.458	0.13
$F_A^{d,s} [10^{-5}]$	3450.12	-3.612	-2.581	0.123	2.167	25.64	-2.21	-0.476	0.13
$F_V^b [10^{-5}]$	1621.40	-4.888	-1.756	0.134	1.27	-2.87	-1.5	-0.21	0.13
$F_A^b [10^{-5}]$	3408.08	-3.110	-2.458	0.120	1.754	-21.04	-0.60	-0.15	0.14

Form fact.	a_9	a_{10}	a_{11}	a_{12}	a_{13}	a_{14}	a_{15}	a_{16}	max. dev.
$F_V^\ell [10^{-5}]$	-0.13	-0.037	-0.11	2.78	-43.15	1.2	-8.9	1396	< 0.02
$F_A^\ell [10^{-5}]$	-3.87	-0.28	-0.28	7.6	-1.9	1.1	-6.3	6912	< 0.04
$F_{V,A}^\nu [10^{-5}]$	-3.93	-0.29	-0.28	7.6	-1.7	1.1	-6.3	6939	< 0.04
$F_V^{u,c} [10^{-5}]$	-3.89	-0.16	-0.22	6.5	-145.6	0.98	-6.9	5648	< 0.03
$F_A^{u,c} [10^{-5}]$	-4.86	-0.27	-0.27	7.5	-1.9	1.1	-6.3	6918	< 0.04
$F_V^{d,s} [10^{-5}]$	-5.56	-0.24	-0.28	8.1	-132	0.95	-6.0	7492	< 0.04
$F_A^{d,s} [10^{-5}]$	-5.03	-0.27	-0.27	7.5	-1.6	1.1	-6.2	6923	< 0.04
$F_V^b [10^{-5}]$	-1.3	-0.27	-0.32	12.5	-131	0.95	-5.7	7451	< 0.04
$F_A^b [10^{-5}]$	0.87	-0.31	-0.31	13.5	-2.4	1.1	-7.2	6918	< 0.04

Table 7. Coefficients for the parameterization formula (2.9) for various form factors (X). Within the ranges $25 \text{ GeV} < M_H < 225 \text{ GeV}$, $155 \text{ GeV} < m_t < 195 \text{ GeV}$, $\alpha_s = 0.1184 \pm 0.0050$, $\Delta\alpha = 0.0590 \pm 0.0005$ and $M_Z = 91.1876 \pm 0.0084 \text{ GeV}$, the formula approximates the full result with average and maximal deviations given in the last two columns.

5 Theoretical error estimates for missing higher order corrections

The main theory uncertainty of the results presented in this paper stems from unknown three- and four-loop corrections. The leading missing orders are $\mathcal{O}(\alpha^3)$, $\mathcal{O}(\alpha^2\alpha_s)$, $\mathcal{O}(\alpha\alpha_s^2)$, and $\mathcal{O}(\alpha\alpha_s^3)$. Partial results for these contributions, in the limit of a large top Yukawa coupling y_t , have already been computed [74–80]. Therefore, when evaluating the impact of theory uncertainties, it is always implied that we refer to these contributions *beyond* the leading- y_t limit.

There are a number of different methods for assessing theory uncertainties from unknown higher orders, none of which is fully reliable. Rather, they should be taken as an order-of-magnitude estimate of the size of these terms. A convenient and widely applicable method is based on the assumption that the first few orders of the perturbation series approximately follow a geometric series [35, 37, 86]. In this way one obtains as an ansatz

$$\begin{aligned}
 \mathcal{O}(\alpha^3) - \mathcal{O}(\alpha_t^3) &\sim \frac{\mathcal{O}(\alpha^2) - \mathcal{O}(\alpha_t^2)}{\mathcal{O}(\alpha)} \mathcal{O}(\alpha^2), \\
 \mathcal{O}(\alpha^2\alpha_s) - \mathcal{O}(\alpha_t^2\alpha_s) &\sim \frac{\mathcal{O}(\alpha^2) - \mathcal{O}(\alpha_t^2)}{\mathcal{O}(\alpha)} \mathcal{O}(\alpha\alpha_s), \\
 \mathcal{O}(\alpha\alpha_s^2) - \mathcal{O}(\alpha_t\alpha_s^2) &\sim \frac{\mathcal{O}(\alpha\alpha_s) - \mathcal{O}(\alpha_t\alpha_s)}{\mathcal{O}(\alpha)} \mathcal{O}(\alpha\alpha_s), \\
 \mathcal{O}(\alpha\alpha_s^3) - \mathcal{O}(\alpha_t\alpha_s^3) &\sim \frac{\mathcal{O}(\alpha\alpha_s) - \mathcal{O}(\alpha_t\alpha_s)}{\mathcal{O}(\alpha)} \mathcal{O}(\alpha\alpha_s^2),
 \end{aligned} \tag{5.1}$$

where $\alpha_t = y_t^2/(4\pi)$. Since we are only interested in the missing higher orders beyond the leading large- y_t limit, the same leading large- y_t approximations have been subtracted in the numerators on the right-hand sides.

The contribution of these estimates to the overall theory error evaluation is shown in Tab. 8 for various pseudo-observables, and in Tab. 9 for the Z -boson form factors. Note that the error estimate for $\sin^2\theta_{\text{eff}}^\ell$ is slightly improved compared to Refs. [28, 29] due to the inclusion of $\mathcal{O}(\alpha_t\alpha_s^3)$ corrections from Refs. [78–80].

Nevertheless, we would also like to remind the reader that any estimate of the theory error from missing higher orders is not a precise prediction. Therefore it is generally desirable to ensure that the theory error is sub-dominant in any phenomenological analysis. Comparing the numbers in Tab. 8 to current measurement results [57, 58], this is clearly seen to be the case.

6 Conclusions and outlook

In this study, we have presented some phenomenologically useful applications of the recently completed electroweak two-loop calculation of Z -boson vertex corrections

Observable	$\alpha\alpha_s^2$	$\alpha\alpha_s^3$	$\alpha^2\alpha_s$	α^3	Total
$\Gamma_{e,\mu,\tau}$ [MeV]	0.008	0.001	0.010	0.013	0.018
Γ_ν [MeV]	0.008	0.001	0.008	0.011	0.016
$\Gamma_{u,c}$ [MeV]	0.025	0.004	0.08	0.07	0.11
$\Gamma_{d,s}$ [MeV]	0.016	0.003	0.06	0.05	0.08
Γ_b [MeV]	0.11	0.02	0.13	0.06	0.18
Γ_Z [MeV]	0.23	0.035	0.21	0.20	0.4
R_ℓ [10^{-3}]	2.5	0.4	3.6	3.9	6
R_c [10^{-5}]	1.6	0.3	3.4	3.0	5
R_b [10^{-5}]	5.5	0.9	6.4	3.7	10
σ_{had}^0 [pb]	0.2	0.03	4.2	3.7	6
$\sin^2 \theta_{\text{eff}}^\ell$ [10^{-5}]	—	0.3	3.0	3.1	4.3
$\sin^2 \theta_{\text{eff}}^b$ [10^{-5}]	0.7	0.4	4.3	3.2	5.3

Table 8. *Leading unknown higher-order corrections and their estimated order of magnitude for various pseudo-observables. The different orders always correspond to missing higher orders beyond the known approximations in the limit of a large top Yukawa coupling. The total theory error is obtained by adding the individual orders in quadrature.*

Observable	$\alpha\alpha_s^2$	$\alpha\alpha_s^3$	$\alpha^2\alpha_s$	α^3	Total
F_V^ℓ [10^{-5}]	0.03	0.004	0.2	0.4	0.5
F_A^ℓ [10^{-5}]	0.17	0.03	0.3	0.3	0.4
$F_{V,A}^\nu$ [10^{-5}]	0.16	0.02	0.3	0.5	0.6
$F_V^{u,c}$ [10^{-5}]	0.09	0.01	0.4	0.2	0.5
$F_A^{u,c}$ [10^{-5}]	0.17	0.03	0.3	0.4	0.5
$F_V^{d,s}$ [10^{-5}]	0.2	0.03	0.6	0.8	1.1
$F_A^{d,s}$ [10^{-3}]	0.3	0.04	0.4	0.5	0.7
F_V^b [10^{-5}]	0.2	0.03	0.8	0.7	1.1
F_A^b [10^{-5}]	0.3	0.04	0.1	0.1	0.3

Table 9. *Same as Tab. 8, but for various form factors.*

[36, 37]. The work collects multi-year efforts of several groups for predictions of the EWPOs related to the Z peak up to electroweak full two-loop accuracy, supplemented by leading QCD higher-order terms. We have determined the two-loop electroweak contributions with a net relative numerical accuracy of about four digits. This ensures that these two-loop results will be known with sufficient accuracy even when adding the next perturbative order, as it might be needed for applications at the next generation of e^+e^- colliders.

For practical applications, the results for the EWPOs, as well as for the Z -boson vertex form factors, have been presented in terms of simple parameterization formulae, whose coefficients have been fitted to the full numerical computation. It is planned to include these fitting formulae into a new version of the weak library DIZET of ZFITTER [45, 46, 48, 87–89].³ The accuracy of the fitting formulae is less than our full numerical two-loop calculation, but more than sufficient for present-day purposes. For the future FCC-ee Tera- Z project, it may be necessary to provide more precise formulae by including more terms with higher powers of the input parameters.

Finally, we would like to make a few comments on the prospects for the calculation of electroweak three-loop corrections, which will be necessary for the level of precision foreseen for FCC-ee and similar e^+e^- collider proposals. This electroweak third order, by itself, will be needed with only about two digits accuracy [90]. The generation of the amplitudes for $\mathcal{O}(10^4-10^5)$ diagrams as well as the evaluation of the Lorentz and Dirac algebra are routine tasks performed by computer algebra programs, and they should be straightforward with increased computing power in the future. Potential specific problems related to the treatment of γ_5 at three-loop level have to be controlled [91].

The most challenging problem will certainly be the stable numerical computation of three-loop Feynman integrals with several different internal mass scales. At two loops, we did not perform any reduction of the Feynman integrals to a smaller number of masters and thus had to calculate about 1000 previously unknown numerical integrals. In the next perturbative order, it may be advantageous to perform such a reduction to masters, given the ever increasing performance of programs like KIRA [92, 93], Reduze 2 [94], FIRE [95, 95], and LiteRED [96].

For the calculation of the (master) integrals themselves, it is desirable to have a procedure to automatically isolate and treat the ultra-violet and infra-red singularities. Although there is rapid progress in several analytical approaches to complicated loop integrals [6, 7], one has to expect that the more complicated ones will have to be done numerically. An additional complication is that for integrals with physical cuts, their stable numerical evaluation becomes more challenging. There are two kinds of software packages available that address these problems, based on either sector decomposition (SD) as realized in the SecDec project [97–102] and the FIESTA project

³Private communication by L. Kalinovskaya for the ZFITTER/DIZET support team.

[103–106], or based on Mellin-Barnes (MB) transformations, as implemented in the AMBRE project [107–120]. Sector decomposition is typically advantageous for integrals with many different mass scales, while the MB approach is more efficient for integrals with fewer independent parameters. Both methods certainly have room for crucial improvements. Several other numerical integration methods, as reviewed e.g. in Refs. [6, 7], are useful for certain classes of multi-loop integrals, even though they are less general than the SD and MB approaches. Overall, numerical loop integration techniques are well positioned to meet the necessary future precision demands.

An electroweak three-loop result for the Z -peak EWPOs must be accompanied by improved calculations of the corrections needed to translate EWPOs to real observables. These include initial-state and final-state QED corrections and their interference as well as higher-order terms of the Laurent series expansion about the Z resonance pole [6]. The latter will, for example, involve massive two-loop box diagrams. A complete accounting of the required correction terms is still lacking.

To summarize, we have completed the electroweak two-loop predictions for the EWPOs of the Z resonance and collect here an extensive set of new fitting formulae for them. On a longer time scale, the calculation of the next perturbative order for the calculation of the EWPOs will be necessary and, with proper investments, realistically accessible.

Acknowledgments

The authors are grateful to V. Yermolchyk for checks and useful comments. The work of *I.D.* was supported by a research grant of Deutscher Akademischer Austauschdienst (DAAD) and by Deutsches Elektronensynchrotron DESY and CERN. The work of *J.G.* was supported by the Polish National Science Centre (NCN) under the Grant Agreement 2017/25/B/ST2/01987 and by the program International mobilities for research activities of the University of Hradec Králové, CZ.02.2.69/0.0/0.0/16_027/0008487. The work of *A.F.* was supported in part by the National Science Foundation under grant no. PHY-1820760. The work of *T.R.* is funded by Deutsche Rentenversicherung Bund. He is supported in part by a 2015 Alexander von Humboldt Honorary Research Scholarship of the Foundation for Polish Sciences (FNP) and by the Polish National Science Centre (NCN) under the Grant Agreement 2017/25/B/ST2/01987. *J.U.* received funding from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme under grant agreement no. 647356 (CutLoops). The project was partly supported by the COST (European Cooperation in Science and Technology) Action CA16201 PARTICLEFACE. We finally acknowledge kind support for a working meeting by Prof. Peter Uwer at Humboldt-Universität zu Berlin.

References

- [1] A. Blondel, F. Dydak, A. Para, W. Blum, Measurement of the Z^0 Mass and Width at LEP, in [3].
- [2] G. Arduini, et al., LEP-1 operation, 1989 - 1995, Conf. Proc. C960610 (1996) 286–288.
<http://accelconf.web.cern.ch/AccelConf/e96/PAPERS/ORALS/M0002A.PDF>.
- [3] J. Ellis, R. Peccei (eds.), Physics at LEP 1, Yellow Report CERN 86-02 (1986),
<http://cds.cern.ch/record/166310/files/CERN-86-02-V-1.pdf>.
- [4] G. Altarelli, R. Kleiss, C. Verzegnassi (eds.), Z physics at LEP 1, Yellow Report CERN 89-08 (1989): vol. 1: Standard physics, vol. 2: Higgs search and new physics, vol. 3: Event generators and software,
<http://dx.doi.org/10.5170/CERN-1989-008-V-1>,
<http://dx.doi.org/10.5170/CERN-1989-008-V-2>,
<http://dx.doi.org/10.5170/CERN-1989-008-V-3>.
- [5] D. Bardin, W. Hollik, G. Passarino (eds.), Reports of the working group on precision calculations for the Z resonance, Yellow Report CERN 95-03 (1995), parts I to III, 410 p., <http://cds.cern.ch/record/280836/files/CERN-95-03.pdf>.
[doi:10.5170/CERN-1995-003](https://doi.org/10.5170/CERN-1995-003).
- [6] A. Blondel, J. Gluza, S. Jadach, P. Janot, T. Riemann (eds.), Standard Model Theory for the FCC-ee: The Tera- Z , report on the mini workshop on precision EW and QCD calculations for the FCC studies: methods and techniques, CERN, Geneva, Switzerland, January 12-13, 2018; to appear as CERN Yellow Report 2019. [arXiv:1809.01830](https://arxiv.org/abs/1809.01830).
- [7] A. Blondel, J. Gluza, S. Jadach, P. Janot, T. Riemann (eds.), Theory report on the 11th FCC-ee workshop: Theory and Experiments, 8-11 January 2019, CERN, Geneva, Switzerland. [arXiv:1905.05078](https://arxiv.org/abs/1905.05078).
- [8] A. Freitas, J. Gluza, S. Jadach, T. Riemann, *Loops, matrix elements, EWPOs*, section C.2 in [6].
- [9] B. Lynn, G.D. Starkman (orgs.), The Standard Model at 50 Years: A Celebratory Symposium, hosted by the Case Western Reserve University, Department of Physics, Cleveland, Ohio, June 1-4, 2018. Proceedings in preparation.
<https://indico.cern.ch/event/704471/timetable/> .
- [10] M. Consoli, A. Sirlin, The role of the one loop electroweak effects in $e^+e^- \rightarrow \mu^+\mu^-$, in [3].
- [11] A. Sirlin, Theoretical considerations concerning the Z^0 mass, Phys. Rev. Lett. 67 (1991) 2127–2130. [doi:10.1103/PhysRevLett.67.2127](https://doi.org/10.1103/PhysRevLett.67.2127).
- [12] S. Willenbrock, G. Valencia, On the definition of the Z boson mass, Phys. Lett. B259 (1991) 373–376. [doi:10.1016/0370-2693\(91\)90843-F](https://doi.org/10.1016/0370-2693(91)90843-F).

- [13] R. G. Stuart, Gauge invariance, analyticity and physical observables at the Z^0 resonance, Phys. Lett. B262 (1991) 113–119.
[doi:10.1016/0370-2693\(91\)90653-8](https://doi.org/10.1016/0370-2693(91)90653-8).
- [14] A. Sirlin, Observations concerning mass renormalization in the electroweak theory, Phys. Lett. B267 (1991) 240–242,
<https://lib-extopc.kek.jp/preprints/PDF/1991/9111/9111179.pdf>.
[doi:10.1016/0370-2693\(91\)91254-S](https://doi.org/10.1016/0370-2693(91)91254-S).
- [15] R. G. Stuart, General renormalization of the gauge invariant perturbation expansion near the Z^0 resonance, Phys. Lett. B272 (1991) 353–358,
<https://lib-extopc.kek.jp/preprints/PDF/1991/9109/9109014.pdf>.
[doi:10.1016/0370-2693\(91\)91842-J](https://doi.org/10.1016/0370-2693(91)91842-J).
- [16] R. G. Stuart, The structure of the Z^0 resonance and the physical properties of the Z^0 boson, Phys. Rev. Lett. 70 (1993) 3193–3196.
[doi:10.1103/PhysRevLett.70.3193](https://doi.org/10.1103/PhysRevLett.70.3193).
- [17] H. G. J. Veltman, Mass and width of unstable gauge bosons, Z. Phys. C62 (1994) 35–52. [doi:10.1007/BF01559523](https://doi.org/10.1007/BF01559523).
- [18] M. Passera, A. Sirlin, Radiative corrections to W and quark propagators in the resonance region, Phys. Rev. D58 (1998) 113010. [arXiv:hep-ph/9804309](https://arxiv.org/abs/hep-ph/9804309),
[doi:10.1103/PhysRevD.58.113010](https://doi.org/10.1103/PhysRevD.58.113010).
- [19] D. Bardin, P. Khristova, O. Fedorenko, On the lowest order electroweak corrections to spin 1/2 fermion scattering. 1. The one loop diagram, Nucl. Phys. B175 (1980) 435. [doi:10.1016/0550-3213\(80\)90021-8](https://doi.org/10.1016/0550-3213(80)90021-8).
- [20] D. Bardin, P. Khristova, O. Fedorenko, On the lowest order electroweak corrections to spin 1/2 fermion scattering. 2. The one loop amplitudes, Nucl. Phys. B197 (1982) 1. [doi:10.1016/0550-3213\(82\)90152-3](https://doi.org/10.1016/0550-3213(82)90152-3).
- [21] W. J. Marciano, A. Sirlin, Radiative corrections to neutrino induced neutral current phenomena in the $SU(2)_L \times U(1)$ theory, Phys. Rev. D22 (1980) 2695, Erratum: Phys. Rev. D31 (1985) 213.
[doi:10.1103/PhysRevD.31.213](https://doi.org/10.1103/PhysRevD.31.213), [10.1103/PhysRevD.22.2695](https://doi.org/10.1103/PhysRevD.22.2695).
- [22] W. J. Marciano, A. Sirlin, Testing the Standard Model by precise determinations of W^\pm and Z masses, Phys. Rev. D29 (1984) 945, Erratum: Phys. Rev. D31 (1985) 213. [doi:10.1103/PhysRevD.29.945](https://doi.org/10.1103/PhysRevD.29.945), [10.1103/PhysRevD.31.213](https://doi.org/10.1103/PhysRevD.31.213).
- [23] A. Akhundov, D. Bardin, T. Riemann, Electroweak one loop corrections to the decay of the neutral vector boson, Nucl. Phys. B276 (1986) 1, Dubna preprint JINR-E2-85-617 (Aug 1985). [doi:10.1016/0550-3213\(86\)90014-3](https://doi.org/10.1016/0550-3213(86)90014-3).
- [24] J. Bernabeu, A. Pich, A. Santamaria, $\Gamma(Z \rightarrow b\bar{b})$: A signature of hard mass terms for a heavy top, Phys. Lett. B200 (1988) 569–574.
[doi:10.1016/0370-2693\(88\)90173-6](https://doi.org/10.1016/0370-2693(88)90173-6).
- [25] F. Jegerlehner, Precision tests of electroweak interaction parameters, in: R. Manka,

- M. Zralek (eds.), proceedings of the 11th Int. School of Theoretical Physics *Testing the standard model*, Szczyrk, Poland, Sep 18-22, 1987 (Singapore, World Scientific, 1988), pp. 33-108, Bielefeld preprint BI-TP-87/16,
<https://lib-extopc.kek.jp/preprints/PDF/1988/8801/8801263.pdf>.
- [26] W. Beenakker, W. Hollik, The width of the Z boson, *Z. Phys. C*40 (1988) 141.
[doi:10.1007/BF01559728](https://doi.org/10.1007/BF01559728).
- [27] M. Awramik, M. Czakon, A. Freitas, G. Weiglein, Complete two-loop electroweak fermionic corrections to $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ and indirect determination of the Higgs boson mass, *Phys. Rev. Lett.* 93 (2004) 201805. [arXiv:hep-ph/0407317](https://arxiv.org/abs/hep-ph/0407317),
[doi:10.1103/PhysRevLett.93.201805](https://doi.org/10.1103/PhysRevLett.93.201805).
- [28] M. Awramik, M. Czakon, A. Freitas, Bosonic corrections to the effective weak mixing angle at $O(\alpha^2)$, *Phys. Lett. B*642 (2006) 563–566. [arXiv:hep-ph/0605339](https://arxiv.org/abs/hep-ph/0605339),
[doi:10.1016/j.physletb.2006.07.035](https://doi.org/10.1016/j.physletb.2006.07.035).
- [29] M. Awramik, M. Czakon, A. Freitas, Electroweak two-loop corrections to the effective weak mixing angle, *JHEP* 0611 (2006) 048. [arXiv:hep-ph/0608099](https://arxiv.org/abs/hep-ph/0608099),
[doi:10.1088/1126-6708/2006/11/048](https://doi.org/10.1088/1126-6708/2006/11/048).
- [30] W. Hollik, U. Meier, S. Uccirati, The effective electroweak mixing angle $\sin^2 \theta^{\text{eff}}$ with two-loop fermionic contributions, *Nucl. Phys. B*731 (2005) 213–224.
[arXiv:hep-ph/0507158](https://arxiv.org/abs/hep-ph/0507158), [doi:10.1016/j.nuclphysb.2005.10.015](https://doi.org/10.1016/j.nuclphysb.2005.10.015).
- [31] W. Hollik, U. Meier, S. Uccirati, The effective electroweak mixing angle $\sin^2 \theta^{\text{eff}}$ with two-loop bosonic contributions, *Nucl. Phys. B*765 (2007) 154–165.
[arXiv:hep-ph/0610312](https://arxiv.org/abs/hep-ph/0610312), [doi:10.1016/j.nuclphysb.2006.12.001](https://doi.org/10.1016/j.nuclphysb.2006.12.001).
- [32] M. Awramik, M. Czakon, A. Freitas, B. Kniehl, Two-loop electroweak fermionic corrections to $\sin^2 \theta_{\text{eff}}^{\text{bb}}$, *Nucl. Phys. B*813 (2009) 174–187. [arXiv:0811.1364](https://arxiv.org/abs/0811.1364),
[doi:10.1016/j.nuclphysb.2008.12.031](https://doi.org/10.1016/j.nuclphysb.2008.12.031).
- [33] A. Freitas, Two-loop fermionic electroweak corrections to the Z -boson width and production rate, *Phys. Lett. B*730 (2014) 50–52. [arXiv:1310.2256](https://arxiv.org/abs/1310.2256),
[doi:10.1016/j.physletb.2014.01.017](https://doi.org/10.1016/j.physletb.2014.01.017).
- [34] A. Freitas, Electroweak precision tests in the LHC era and Z -decay form factors at two-loop level, *PoS LL2014* (2014) 050, <https://pos.sissa.it/211/050/pdf>,
<https://doi.org/10.22323/1.211.0050>. [arXiv:1406.6980](https://arxiv.org/abs/1406.6980).
- [35] A. Freitas, Higher-order electroweak corrections to the partial widths and branching ratios of the Z boson, *JHEP* 04 (2014) 070. [arXiv:1401.2447](https://arxiv.org/abs/1401.2447),
[doi:10.1007/JHEP04\(2014\)070](https://doi.org/10.1007/JHEP04(2014)070).
- [36] I. Dubovyk, A. Freitas, J. Gluza, T. Riemann, J. Usovitsch, The two-loop electroweak bosonic corrections to $\sin^2 \theta_{\text{eff}}^{\text{b}}$, *Phys. Lett. B*762 (2016) 184–189.
[arXiv:1607.08375](https://arxiv.org/abs/1607.08375), [doi:10.1016/j.physletb.2016.09.012](https://doi.org/10.1016/j.physletb.2016.09.012).
- [37] I. Dubovyk, A. Freitas, J. Gluza, T. Riemann, J. Usovitsch, Complete electroweak

- two-loop corrections to Z boson production and decay, Phys. Lett. B783 (2018) 86–94. [arXiv:1804.10236](#), [doi:10.1016/j.physletb.2018.06.037](#).
- [38] A. Leike, T. Riemann, J. Rose, S matrix approach to the Z line shape, Phys. Lett. B273 (1991) 513–518. [arXiv:hep-ph/9508390](#), [doi:10.1016/0370-2693\(91\)90307-C](#).
- [39] T. Riemann, Cross-section asymmetries around the Z peak, Phys. Lett. B293 (1992) 451–456. [arXiv:hep-ph/9506382](#), [doi:10.1016/0370-2693\(92\)90911-M](#).
- [40] S. Kirsch, T. Riemann, SMATASY: A program for the model independent description of the Z resonance, Comput. Phys. Commun. 88 (1995) 89–108. [arXiv:hep-ph/9408365](#), [doi:10.1016/0010-4655\(95\)00016-9](#).
- [41] T. Riemann, S-matrix Approach to the Z Resonance, Acta Phys. Polon. B46 (11) (2015) 2235. [arXiv:1610.04501](#), [doi:10.5506/APhysPolB.46.2235](#).
- [42] A. Freitas, J. Gluza, S. Jadach, *QED deconvolution and pseudo-observables at FCC-ee precision*, section C.3 in [6].
- [43] S. Jadach, M. Skrzypek, QED challenges at FCC-ee precision measurements. [arXiv:1903.09895](#).
- [44] ALEPH collab., DELPHI collab., L3 collab., OPAL collab., SLD collab., LEP Electroweak Working Group, SLD Electroweak Group, SLD Heavy Flavour Group, S. Schael, et al., Precision electroweak measurements on the Z resonance, Phys. Rept. 427 (2006) 257–454. [arXiv:hep-ex/0509008](#), [doi:10.1016/j.physrep.2005.12.006](#).
- [45] D. Bardin, M. Bilenky, P. Christova, M. Jack, L. Kalinovskaya, A. Olchevski, S. Riemann, T. Riemann, ZFITTER 6.21: A semi-analytical program for fermion pair production in e^+e^- annihilation, Comput. Phys. Commun. 133 (2001) 229–395. [arXiv:hep-ph/9908433](#), [doi:10.1016/S0010-4655\(00\)00152-1](#).
- [46] A. Arbuzov, M. Awramik, M. Czakon, A. Freitas, M. Grünewald, K. Mönig, S. Riemann, T. Riemann, ZFITTER: A semi-analytical program for fermion pair production in e^+e^- annihilation, from version 6.21 to version 6.42, Comput. Phys. Commun. 174 (2006) 728–758. [arXiv:hep-ph/0507146](#), [doi:10.1016/j.cpc.2005.12.009](#).
- [47] M. Awramik, M. Czakon, A. Freitas, G. Weiglein, Precise prediction for the W boson mass in the Standard Model, Phys. Rev. D69 (2004) 053006. [arXiv:hep-ph/0311148](#), [doi:10.1103/PhysRevD.69.053006](#).
- [48] A. Akhundov, A. Arbuzov, S. Riemann, T. Riemann, The ZFITTER project, Phys. Part. Nucl. 45 (3) (2014) 529–549. [arXiv:1302.1395](#), [doi:10.1134/S1063779614030022](#).
- [49] P. Baikov, K. Chetyrkin, J. Kühn, J. Rittinger, Complete $O(\alpha_s^4)$ QCD corrections to hadronic Z decays, Phys. Rev. Lett. 108 (2012) 222003. [arXiv:1201.5804v3](#), [doi:10.1103/PhysRevLett.108.222003](#).

- [50] I. Dubovyk, J. Gluza, A. Freitas, K. Grzanka, S. Jadach, T. Riemann, J. Usovitsch, Precision calculations for FCC, poster presented by S. Jadach at the FCC weak 2018, 9-13 April 2018, Amsterdam.
https://indico.cern.ch/event/656491/contributions/2947663/attachments/1622685/2582801/Poster-FCC-Amsterdam_SJadach_et_al.pdf.
- [51] A. Abada, et al., Future Circular Collider: Vol. 1 “Physics opportunities”,
<http://inspirehep.net/record/1713706/files/CERN-ACC-2018-0056.pdf>.
- [52] I. Dubovyk, A. Freitas, J. Gluza, K. Grzanka, S. Jadach, T. Riemann, J. Usovitsch, Theory status of Z-boson physics, chapter B of [6].
- [53] D. Bardin, A. Leike, T. Riemann, M. Sachwitz, Energy dependent width effects in e^+e^- annihilation near the Z boson pole, Phys. Lett. B206 (1988) 539–542.
[doi:10.1016/0370-2693\(88\)91625-5](https://doi.org/10.1016/0370-2693(88)91625-5).
- [54] D. Bardin, S. Riemann, T. Riemann, Electroweak one loop corrections to the decay of the charged vector boson, Z. Phys. C32 (1986) 121,
<https://lib-extopc.kek.jp/preprints/PDF/1986/8607/8607019.pdf>.
[doi:10.1007/BF01441360](https://doi.org/10.1007/BF01441360).
- [55] A. Denner, T. Sack, The W boson width, Z. Phys. C46 (1990) 653–663.
[doi:10.1007/BF01560267](https://doi.org/10.1007/BF01560267).
- [56] K. G. Chetyrkin, J. H. Kühn, A. Kwiatkowski, QCD corrections to the e^+e^- cross-section and the Z boson decay rate, in [5]. See also: QCD corrections to the e^+e^- cross-section and the Z boson decay rate: Concepts and results, LBL-36678-REV and Phys. Rept. 277 (1996) 189-281. [arXiv:hep-ph/9503396](https://arxiv.org/abs/hep-ph/9503396),
[doi:10.1016/S0370-1573\(96\)00012-9](https://doi.org/10.1016/S0370-1573(96)00012-9).
- [57] ALEPH collab., DELPHI collab., L3 collab., OPAL collab., LEP Electroweak Working Group, S. Schael, et al., Electroweak Measurements in Electron-Positron Collisions at W-Boson-Pair Energies at LEP, Phys. Rept. 532 (2013) 119–244.
[arXiv:1302.3415](https://arxiv.org/abs/1302.3415), [doi:10.1016/j.physrep.2013.07.004](https://doi.org/10.1016/j.physrep.2013.07.004).
- [58] C. Patrignani, et al., Review of Particle Physics, Chin. Phys. C40 (10) (2016) 100001. [doi:10.1088/1674-1137/40/10/100001](https://doi.org/10.1088/1674-1137/40/10/100001).
- [59] M. Steinhauser, Leptonic contribution to the effective electromagnetic coupling constant up to three loops, Phys. Lett. B429 (1998) 158–161.
[arXiv:hep-ph/9803313](https://arxiv.org/abs/hep-ph/9803313).
- [60] M. Davier, A. Hoecker, B. Malaescu, Z. Zhang, Reevaluation of the hadronic vacuum polarisation contributions to the Standard Model predictions of the muon $g - 2$ and $\alpha(m_Z^2)$ using newest hadronic cross-section data, Eur. Phys. J. C77 (12) (2017) 827. [arXiv:1706.09436](https://arxiv.org/abs/1706.09436), [doi:10.1140/epjc/s10052-017-5161-6](https://doi.org/10.1140/epjc/s10052-017-5161-6).
- [61] F. Jegerlehner, Variations on Photon Vacuum Polarization. [arXiv:1711.06089](https://arxiv.org/abs/1711.06089).
- [62] A. Keshavarzi, D. Nomura, T. Teubner, The muon $g - 2$ and $\alpha(M_Z^2)$: a new

- data-based analysis, Phys. Rev. D97 (11) (2018) 114025. [arXiv:1802.02995](#),
[doi:10.1103/PhysRevD.97.114025](#).
- [63] A. Djouadi, C. Verzegnassi, Virtual very heavy top effects in LEP/SLC precision measurements, Phys. Lett. B195 (1987) 265–271.
[doi:10.1016/0370-2693\(87\)91206-8](#).
- [64] A. Djouadi, $O(\alpha\alpha_s)$ vacuum polarization functions of the standard model gauge bosons, Nuovo Cim. A100 (1988) 357. [doi:10.1007/BF02812964](#).
- [65] B. A. Kniehl, Two loop corrections to the vacuum polarizations in perturbative QCD, Nucl. Phys. B347 (1990) 86–104. [doi:10.1016/0550-3213\(90\)90552-0](#).
- [66] B. A. Kniehl, A. Sirlin, Dispersion relations for vacuum polarization functions in electroweak physics, Nucl. Phys. B371 (1992) 141–148.
[doi:10.1016/0550-3213\(92\)90232-Z](#).
- [67] A. Djouadi, P. Gambino, Electroweak gauge boson selfenergies: Complete QCD corrections, Phys. Rev. D49 (1994) 3499–3511, Erratum: Phys. Rev. D53 (1996) 4111. [arXiv:hep-ph/9309298](#),
[doi:10.1103/PhysRevD.49.3499](#), [10.1103/PhysRevD.53.4111](#).
- [68] A. Czarnecki, J. H. Kühn, Nonfactorizable QCD and electroweak corrections to the hadronic Z boson decay rate, Phys. Rev. Lett. 77 (1996) 3955–3958.
[arXiv:hep-ph/9608366](#), [doi:10.1103/PhysRevLett.77.3955](#).
- [69] R. Harlander, T. Seidensticker, M. Steinhauser, Complete corrections of order $\alpha\alpha_s$ to the decay of the Z boson into bottom quarks, Phys. Lett. B426 (1998) 125–132.
[arXiv:hep-ph/9712228](#), [doi:10.1016/S0370-2693\(98\)00220-2](#).
- [70] J. Fleischer, O. Tarasov, F. Jegerlehner, P. Raczka, Two loop $O(\alpha_s G_\mu m_t^2)$ corrections to the partial decay width of the Z^0 into $b\bar{b}$ final states in the large top mass limit, Phys. Lett. B293 (1992) 437–444.
[doi:10.1016/0370-2693\(92\)90909-N](#).
- [71] G. Buchalla, A. J. Buras, QCD corrections to the $\bar{s}dZ$ vertex for arbitrary top quark mass, Nucl. Phys. B398 (1993) 285–300.
[doi:10.1016/0550-3213\(93\)90110-B](#).
- [72] G. Degrandi, Current algebra approach to heavy top effects in $Z \rightarrow b + \bar{b}$, Nucl. Phys. B407 (1993) 271–289. [arXiv:hep-ph/9302288](#),
[doi:10.1016/0550-3213\(93\)90058-W](#).
- [73] K. Chetyrkin, A. Kwiatkowski, M. Steinhauser, Leading top mass corrections of order $O(\alpha\alpha_s m_t^2/m_W^2)$ to partial decay rate $\Gamma(Z \rightarrow b\bar{b})$, Mod. Phys. Lett. A8 (1993) 2785–2792. [doi:10.1142/S0217732393003172](#).
- [74] L. Avdeev, J. Fleischer, S. Mikhailov, O. Tarasov, $O(\alpha\alpha_s^2)$ correction to the electroweak ρ parameter, Phys. Lett. B336 (1994) 560–566, Erratum: Phys. Lett. B349 (1995) 597. [arXiv:hep-ph/9406363](#),
[doi:10.1016/0370-2693\(94\)90573-8](#), [10.1016/0370-2693\(95\)00269-Q](#).

- [75] K. Chetyrkin, J. H. Kühn, M. Steinhauser, Corrections of order $O(G_F M_t^2 \alpha_s^2)$ to the ρ parameter, Phys. Lett. B351 (1995) 331–338, doi:10.1016/0370-2693(95)00380-4. arXiv:hep-ph/9502291, doi:10.1016/0370-2693(95)00380-4.
- [76] J. J. van der Bij, K. G. Chetyrkin, M. Faisst, G. Jikia, T. Seidensticker, Three loop leading top mass contributions to the ρ parameter, Phys. Lett. B498 (2001) 156–162. arXiv:hep-ph/0011373, doi:10.1016/S0370-2693(01)00002-8.
- [77] M. Faisst, J. H. Kühn, T. Seidensticker, O. Veretin, Three loop top quark contributions to the ρ parameter, Nucl. Phys. B665 (2003) 649–662. arXiv:hep-ph/0302275, doi:10.1016/S0550-3213(03)00450-4.
- [78] Y. Schröder, M. Steinhauser, Four-loop singlet contribution to the ρ parameter, Phys. Lett. B622 (2005) 124–130. arXiv:hep-ph/0504055, doi:10.1016/j.physletb.2005.06.085.
- [79] K. G. Chetyrkin, M. Faisst, J. H. Kühn, P. Maierhöfer, C. Sturm, Four-loop QCD corrections to the ρ parameter, Phys. Rev. Lett. 97 (2006) 102003. arXiv:hep-ph/0605201, doi:10.1103/PhysRevLett.97.102003.
- [80] R. Boughezal, M. Czakon, Single scale tadpoles and $O(G_F m_t^2 \alpha_s^3)$ corrections to the ρ parameter, Nucl. Phys. B755 (2006) 221–238. arXiv:hep-ph/0606232, doi:10.1016/j.nuclphysb.2006.08.007.
- [81] P. Baikov, K. Chetyrkin, J. Kühn, Order α_s^4 QCD corrections to Z and τ decays, Phys. Rev. Lett. 101 (2008) 012002. arXiv:0801.1821, doi:10.1103/PhysRevLett.101.012002.
- [82] A. Kataev, Higher order $O(\alpha^2)$ and $O(\alpha\alpha_s)$ corrections to $\sigma_{tot}(e^+e^- \rightarrow hadrons)$ and Z-boson decay rate, Phys. Lett. B287 (1992) 209–212. doi:10.1016/0370-2693(92)91901-K.
- [83] A. Abada, et al., FCC Physics Opportunities, Eur. Phys. J. C79 (6) (2019) 474. doi:10.1140/epjc/s10052-019-6904-3.
- [84] CEPC Study Group, J. Guimares da Costa, Y. Gao, S. Jin, J. Qian, C. Tully, C. Young, L. Wang, M. Ruan, H. Zhu, Q. Ouyang et al., CEPC Conceptual Design Report: Volume 2 - Physics & Detector. http://cds.cern.ch/record/2678417. arXiv:1811.10545.
- [85] R. Hawkings, K. Monig, Electroweak and CP violation physics at a linear collider Z factory, Eur. Phys. J. direct 1 (1) (1999) 8. arXiv:hep-ex/9910022, doi:10.1007/s1010599c0008.
- [86] A. Freitas, Numerical multi-loop integrals and applications, Prog. Part. Nucl. Phys. 90 (2016) 201–240. arXiv:1604.00406, doi:10.1016/j.pnpnp.2016.06.004.
- [87] A. Akhundov, D. Bardin, M. Bilenky, P. Christova, L. Kalinovskaya, S. Riemann, T. Riemann, M. Sachwitz, H. Vogt, Fortran program ZFITTER/DIZET v. 6.42, see webpage http://sanc.jinr.ru/users/zfitter/. DIZET support team (June 2019): A. Arbuzov, L. Kalinovskaya, S. Riemann, T. Riemann, V. Yermolchyk.

- [88] D. Bardin, M. Bilenky, G. Mitselmakher, T. Riemann, M. Sachwitz, A realistic approach to the standard Z peak, *Z. Phys. C* 44 (1989) 493, <https://lib-extopc.kek.jp/preprints/PDF/1989/8906/8906215.pdf>, [doi:10.1007/BF01415565](https://doi.org/10.1007/BF01415565).
- [89] D. Y. Bardin, M. S. Bilenky, T. Riemann, M. Sachwitz, H. Vogt, P. C. Christova, DIZET: A program package for the calculation of electroweak one loop corrections for the process $e^+e^- \rightarrow f^+f^-$ around the Z^0 peak, *Comput. Phys. Commun.* 59 (1990) 303–312. [doi:10.1016/0010-4655\(90\)90179-5](https://doi.org/10.1016/0010-4655(90)90179-5).
- [90] I. Dubovyk, A. Freitas, J. Gluza, K. Grzanka, S. Jadach, T. Riemann, J. Usovitsch, *Theory status of Z-boson physics*, section B.1 in [6].
- [91] P. Marquard, D. Stöckinger, *Towards 3- and 4-loop form factors*, section D.1 in [6].
- [92] P. Maierhöfer, J. Usovitsch, P. Uwer, Kira - a Feynman integral reduction program, *Comput. Phys. Commun.* 230 (2018) 99–112. [arXiv:1705.05610](https://arxiv.org/abs/1705.05610), [doi:10.1016/j.cpc.2018.04.012](https://doi.org/10.1016/j.cpc.2018.04.012).
- [93] P. Maierhöfer, J. Usovitsch, Recent developments in Kira, section C.8 in [7].
- [94] A. von Manteuffel, C. Studerus, Reduze 2 - Distributed Feynman Integral Reduction. [arXiv:1201.4330](https://arxiv.org/abs/1201.4330).
- [95] A. V. Smirnov, FIRE5: a C++ implementation of Feynman Integral REduction, *Comput. Phys. Commun.* 189 (2015) 182–191. [arXiv:1408.2372](https://arxiv.org/abs/1408.2372), [doi:10.1016/j.cpc.2014.11.024](https://doi.org/10.1016/j.cpc.2014.11.024).
- [96] R. N. Lee, LiteRed 1.4: a powerful tool for reduction of multiloop integrals, *J. Phys. Conf. Ser.* 523 (2014) 012059. [arXiv:1310.1145](https://arxiv.org/abs/1310.1145), [doi:10.1088/1742-6596/523/1/012059](https://doi.org/10.1088/1742-6596/523/1/012059).
- [97] J. Carter, G. Heinrich, SecDec: A general program for sector decomposition, *Comput. Phys. Commun.* 182 (2011) 1566–1581. [arXiv:1011.5493](https://arxiv.org/abs/1011.5493), [doi:10.1016/j.cpc.2011.03.026](https://doi.org/10.1016/j.cpc.2011.03.026).
- [98] S. Borowka, J. Carter, G. Heinrich, Numerical evaluation of multi-loop integrals for arbitrary kinematics with SecDec 2.0, *Comput. Phys. Commun.* 184 (2013) 396–408. [arXiv:1204.4152](https://arxiv.org/abs/1204.4152), [doi:10.1016/j.cpc.2012.09.020](https://doi.org/10.1016/j.cpc.2012.09.020).
- [99] S. Borowka, G. Heinrich, S. P. Jones, M. Kerner, J. Schlenk, T. Zirke, SecDec-3.0: numerical evaluation of multi-scale integrals beyond one loop, *Comput. Phys. Commun.* 196 (2015) 470–491. [arXiv:1502.06595](https://arxiv.org/abs/1502.06595), [doi:10.1016/j.cpc.2015.05.022](https://doi.org/10.1016/j.cpc.2015.05.022).
- [100] S. Borowka, G. Heinrich, S. Jahn, S. P. Jones, M. Kerner, J. Schlenk, T. Zirke, pySecDec: a toolbox for the numerical evaluation of multi-scale integrals, *Comput. Phys. Commun.* 222 (2018) 313–326. [arXiv:1703.09692](https://arxiv.org/abs/1703.09692), [doi:10.1016/j.cpc.2017.09.015](https://doi.org/10.1016/j.cpc.2017.09.015).
- [101] S. Borowka, G. Heinrich, S. Jahn, S. P. Jones, M. Kerner, J. Schlenk, A GPU

- compatible quasi-Monte Carlo integrator interfaced to pySecDec, *Comp. Phys. Comm.*, online. [arXiv:1811.11720](https://arxiv.org/abs/1811.11720), [doi:10.1016/j.cpc.2019.02.015](https://doi.org/10.1016/j.cpc.2019.02.015).
- [102] S. Jahn, SecDec: a toolbox for the numerical evaluation of multi-scale integrals, *PoS RADCOR2017* (2018) 017. [arXiv:1802.07946](https://arxiv.org/abs/1802.07946), [doi:10.22323/1.290.0017](https://doi.org/10.22323/1.290.0017).
- [103] A. V. Smirnov, M. N. Tentyukov, Feynman Integral Evaluation by a Sector decomposition Approach (FIESTA), *Comput. Phys. Commun.* 180 (2009) 735–746. [arXiv:0807.4129](https://arxiv.org/abs/0807.4129), [doi:10.1016/j.cpc.2008.11.006](https://doi.org/10.1016/j.cpc.2008.11.006).
- [104] A. V. Smirnov, V. A. Smirnov, M. Tentyukov, FIESTA 2: Parallelizeable multiloop numerical calculations, *Comput. Phys. Commun.* 182 (2011) 790–803. [arXiv:0912.0158](https://arxiv.org/abs/0912.0158), [doi:10.1016/j.cpc.2010.11.025](https://doi.org/10.1016/j.cpc.2010.11.025).
- [105] A. V. Smirnov, FIESTA 3: cluster-parallelizable multiloop numerical calculations in physical regions, *Comput. Phys. Commun.* 185 (2014) 2090–2100. [arXiv:1312.3186](https://arxiv.org/abs/1312.3186), [doi:10.1016/j.cpc.2014.03.015](https://doi.org/10.1016/j.cpc.2014.03.015).
- [106] A. V. Smirnov, FIESTA 4: Optimized Feynman integral calculations with GPU support, *Comput. Phys. Commun.* 204 (2016) 189–199. [arXiv:1511.03614](https://arxiv.org/abs/1511.03614), [doi:10.1016/j.cpc.2016.03.013](https://doi.org/10.1016/j.cpc.2016.03.013).
- [107] J. Gluza, K. Kajda, T. Riemann, AMBRE: A Mathematica package for the construction of Mellin-Barnes representations for Feynman integrals, *Comput. Phys. Commun.* 177 (2007) 879–893. [arXiv:0704.2423](https://arxiv.org/abs/0704.2423), [doi:10.1016/j.cpc.2007.07.001](https://doi.org/10.1016/j.cpc.2007.07.001).
- [108] J. Gluza, K. Kajda, T. Riemann, V. Yundin, New results for loop integrals: AMBRE, CSectors, hexagon, *PoS ACAT08* (2008) 124, <https://pos.sissa.it/070/124/pdf>. [arXiv:0902.4830](https://arxiv.org/abs/0902.4830), [doi:10.22323/1.070.0124](https://doi.org/10.22323/1.070.0124).
- [109] J. Gluza, K. Kajda, T. Riemann, V. Yundin, Numerical Evaluation of Tensor Feynman Integrals in Euclidean Kinematics, *Eur. Phys. J. C* 71 (2011) 1516. [arXiv:1010.1667](https://arxiv.org/abs/1010.1667), [doi:10.1140/epjc/s10052-010-1516-y](https://doi.org/10.1140/epjc/s10052-010-1516-y).
- [110] J. Gluza, K. Kajda, T. Riemann, V. Yundin, News on Ambre and CSectors, *Nucl. Phys. Proc. Suppl.* 205-206 (2010) 147–151. [arXiv:1006.4728](https://arxiv.org/abs/1006.4728), [doi:10.1016/j.nuclphysbps.2010.08.034](https://doi.org/10.1016/j.nuclphysbps.2010.08.034).
- [111] J. Blümlein, I. Dubovyk, J. Gluza, M. Ochman, C. Raab, T. Riemann, C. Schneider, Non-planar Feynman integrals, Mellin-Barnes representations, multiple sums, *PoS LL2014* (2014) 052. <https://pos.sissa.it/211/052/pdf>. [arXiv:1407.7832](https://arxiv.org/abs/1407.7832), [doi:10.22323/1.211.0052](https://doi.org/10.22323/1.211.0052).
- [112] I. Dubovyk, J. Gluza, T. Riemann, Non-planar Feynman diagrams and Mellin-Barnes representations with AMBRE 3.0, *J. Phys. Conf. Ser.* 608 (1) (2015) 012070. <https://iopscience.iop.org/article/10.1088/1742-6596/608/1/012070/pdf>. [doi:10.1088/1742-6596/608/1/012070](https://doi.org/10.1088/1742-6596/608/1/012070).

- [113] I. Dubovyk, J. Gluza, T. Riemann, J. Usovitsch, Numerical integration of massive two-loop Mellin-Barnes integrals in Minkowskian regions, PoS LL2016 (2016) 034. <https://pos.sissa.it/260/034/pdf>. [arXiv:1607.07538](https://arxiv.org/abs/1607.07538), [doi:10.22323/1.260.0034](https://doi.org/10.22323/1.260.0034).
- [114] I. Dubovyk, J. Gluza, T. Jelinski, T. Riemann, J. Usovitsch, New prospects for the numerical calculation of Mellin-Barnes integrals in Minkowskian kinematics, Acta Phys. Polon. B48 (2017) 995. [arXiv:1704.02288](https://arxiv.org/abs/1704.02288), [doi:10.5506/APhysPolB.48.995](https://doi.org/10.5506/APhysPolB.48.995).
- [115] I. Dubovyk, J. Gluza, T. Riemann, AMBRE - Construction of Mellin-Barnes integrals for two- and three-loop Z-boson decays, section E.4 of [6].
- [116] I. Dubovyk, Mellin-Barnes representations for multiloop Feynman integrals with applications to 2-loop electroweak Z boson studies, PhD thesis, Universität Hamburg, to be submitted.
- [117] J. Usovitsch, I. Dubovyk, T. Riemann, MBnumerics: Numerical integration of Mellin-Barnes integrals in physical regions, PoS LL2018 (2018) 046. <https://pos.sissa.it/303/046/pdf>. [arXiv:1810.04580](https://arxiv.org/abs/1810.04580), [doi:10.22323/1.303.0046](https://doi.org/10.22323/1.303.0046).
- [118] J. Usovitsch, et al., MBnumerics: Numerical integration of Mellin-Barnes integrals in physical regions. Talk held by J. Usovitsch at LL2018, April 29 to May 4, 2018, St. Goar, Germany. Slides: <https://indico.desy.de/indico/event/16613/session/4/contribution/22/material/slides/0.pdf>.
- [119] J. Usovitsch, I. Dubovyk, T. Riemann, The MBnumerics project, section E.2 of [6].
- [120] J. Usovitsch, Numerical evaluation of Mellin-Barnes integrals in Minkowskian regions and their application to two-loop bosonic electroweak contributions to the weak mixing angle of the $Z\bar{b}b$ -vertex, Ph.D. thesis, DESY (2018). [doi:10.3204/PUBDB-2018-05160](https://doi.org/10.3204/PUBDB-2018-05160).