

DEUTSCHES ELEKTRONEN-SYNCHROTRON
Ein Forschungszentrum der Helmholtz-Gemeinschaft



DESY 21-010
IFT-UAM/CSIC-20-143
FTUAM-20-21
arXiv:2102.00012
January 2021

An Even Lighter QCD Axion

L. Di Luzio, P. Quilez, A. Ringwald

Deutsches Elektronen-Synchrotron DESY, Hamburg

B. Gavela

*Departamento de Física Teórica, Universidad Autónoma de Madrid,
Cantoblanco, Madrid, Spain*

and

*Instituto de Física Teórica, IFT-UAM/CSIC,
Cantoblanco, Madrid, Spain*

ISSN 0418-9833

NOTKESTRASSE 85 - 22607 HAMBURG

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your reports and preprints are promptly included in the
HEP literature database
send them to (if possible by air mail):

DESY Zentralbibliothek Notkestraße 85 22607 Hamburg Germany	DESY Bibliothek Platanenallee 6 15738 Zeuthen Germany
---	---

An even lighter QCD axion

Luca Di Luzio^a, Belen Gavela^{b,c}, Pablo Quilez^a, Andreas Ringwald^a

^a*Deutsches Elektronen-Synchrotron DESY,
Notkestraße 85, D-22607 Hamburg, Germany*

^b*Departamento de Física Teórica, Universidad Autónoma de Madrid,
Cantoblanco, 28049, Madrid, Spain*

^c*Instituto de Física Teórica, IFT-UAM/CSIC,
Cantoblanco, 28049, Madrid, Spain*

Abstract

We explore whether the axion which solves the strong CP problem can naturally be much lighter than the canonical QCD axion. The $Z_{\mathcal{N}}$ symmetry proposed by Hook, with \mathcal{N} mirror and degenerate worlds coexisting in Nature and linked by the axion field, is considered in terms of generic effective axion couplings. We show that the total potential is safely approximated by a single cosine in the large \mathcal{N} limit, and we determine the analytical formula for the exponentially suppressed axion mass. The resulting universal enhancement of all axion interactions relative to those of the canonical QCD axion has a strong impact on the prospects of axion-like particle experiments such as ALPS II, IAXO and many others: experiments searching for generic axion-like particles have in fact discovery potential to solve the strong CP problem. The finite density axion potential is also analyzed and we show that the $Z_{\mathcal{N}}$ asymmetric background of high-density stellar environments sets already significant model-independent constraints: $3 \leq \mathcal{N} \lesssim 47$ for an axion scale $f_a \lesssim 2.4 \times 10^{15}$ GeV, with tantalizing discovery prospects for any value of f_a and down to $\mathcal{N} \sim 9$ with future neutron star and gravitational wave data, down to the ultra-light mass region. In addition, two specific ultraviolet $Z_{\mathcal{N}}$ completions are developed: a composite axion one and a KSVZ-like model with improved Peccei-Quinn quality.

E-mail: luca.diluzio@desy.de, belen.gavela@uam.es,
pablo.quilez@desy.de, andreas.ringwald@desy.de

Contents

1	Introduction	3
2	Down-tuning the axion mass	5
2.1	The Z_2 case	5
2.2	$Z_{\mathcal{N}}$ axion	7
2.3	Axion potential in the large \mathcal{N} limit	10
3	UV completions and alternative scenarios	13
3.1	KSVZ $Z_{\mathcal{N}}$ axion	13
3.2	Composite $Z_{\mathcal{N}}$ axion	16
3.3	Ultra-light QCD axions	17
3.4	A heavier-than-QCD axion	17
4	Experimental probes of down-tuned axions	18
4.1	Axion coupling to photons	18
4.2	Finite density constraints on f_a	19
5	Conclusions	23
A	Holomorphicity properties of $Z_{\mathcal{N}}$ axion potential	24
B	Fourier series of the $Z_{\mathcal{N}}$ axion potential	25
C	Analytical axion mass dependence from hypergeometric functions	26

1 Introduction

The axion experimental program is in a blooming phase, with several new experiments and detection concepts promising the exploration of regions of parameter space thought to be unreachable until a decade ago. Many of those experiments are simply prototypes, awaiting the jump to become ‘big-experiments’, or, in the case of more consolidated techniques, they are still far from saturating their full physics potential. Nonetheless, they sometimes reach sensitivities which go well-beyond astrophysical limits, albeit often still far from the customary QCD axion window.

On the other hand, since axion couplings are inherently ultraviolet (UV) dependent, such early stage experiments already provide valuable probes of the QCD axion parameter space. Imagine for definiteness that ALPS II would detect a signal in 2021, would it be possible to interpret that as an axion that solves the strong CP problem? Since the strong CP problem is one of the strongest motivations for new physics, if an axion-like particle (ALP) will be ever discovered, there or elsewhere, it would be compelling to explore whether it had something to do with the strong CP problem. This work explores whether wide regions in the ALP parameter space, well outside the traditional QCD axion band, may correspond to solutions of the strong CP problem. This is a question of profound theoretical and experimental relevance.

In axion solutions to the strong CP problem¹ both the axion mass and the couplings to ordinary matter scale as $1/f_a$, where f_a is the axion decay constant, denoting the scale of new physics. The precise relation between mass and decay constant depends on the characteristics of the strong interacting sector of the theory. When QCD is the only confining group to which the axion a couples, in which case we denote the axion mass as m_a^{QCD} , they are necessarily linked by the relation [3, 4]

$$m_a^{\text{QCD}} = \frac{\sqrt{\chi_{\text{QCD}}}}{f_a} \simeq m_\pi f_\pi \frac{\sqrt{m_u m_d}}{m_u + m_d} \frac{1}{f_a}, \quad (1.1)$$

where χ_{QCD} , m_π , f_π , m_u and m_d denote respectively the QCD topological susceptibility, the pion mass, its decay constant, and the up and down quark masses. Equation (1.1) is completely model-independent as far as QCD is the only source of the axion mass, and it defines the ‘‘canonical QCD axion’’, also often called ‘‘invisible axion’’. For this axion the $aG_{\mu\nu}\tilde{G}^{\mu\nu}$ coupling to the gluon strength $G_{\mu\nu}$ is directly responsible for the axion mass, since the only source of explicit breaking of the global axial Peccei-Quinn (PQ) symmetry $U(1)_{\text{PQ}}$ is its QCD anomaly. The strength of other axion couplings to Standard Model (SM) fields is instead model-dependent: it varies with the matter content of the UV complete axion model.

In recent years there have been many attempts to enlarge the canonical QCD axion window, by considering UV completions of the axion effective Lagrangian which departed from the minimal DFSZ [5, 6] and KSVZ [7, 8] constructions. Most approaches actually focussed on the possibility of modifying the Wilson coefficient of specific axion-SM effective operators [9–15]. That is, the size of the coupling coefficients, at fixed f_a , is modified. This has for example allowed to populate new regions of the parameter space by moving vertically the axion band in the axion mass versus coupling plane, see Fig. 1 left. The results are then ‘‘channel specific’’: different couplings c are modified differently.

¹That is, via a global chiral $U(1)$ symmetry, exact although hidden (aka spontaneously broken) at the classical level and explicitly broken by instanton effects at the quantum level [1, 2].




Figure 1: Different approaches to enlarge the parameter space of axions that solve the strong CP problem. The canonical QCD axion relation is represented by the lower black line in the $\{m_a, c/f_a\}$ parameter space, where c denotes a generic effective axion coupling. Vertical displacements, possible within pure QCD axion models (i.e. $m_a = m_a^{\text{QCD}}$), are depicted on the left. Horizontal displacements (via enlarged strong gauge sectors) are illustrated on the right for the case of a lighter than usual axion to be explored here.

The parameter space of solutions can be alternatively changed by varying the axion mass at fixed f_a . This corresponds to horizontal displacements of the canonical axion band in the parameter space, see right panel in Fig. 1. It always requires that the magnitude of the relation between the axion mass m_a and $1/f_a$ departs from that in Eq. (1.1): the confining sector of the SM must be enlarged beyond QCD. New instanton sources give then additional contributions to the right-hand side of Eq. (1.1). The practical consequence is a universal modification of the parameter space of *all* axion couplings at a given m_a , at variance with the vertical displacement scenarios. This feature could *a priori* allow for the two mechanisms in Fig. 1 to be distinguished.²

Examples of horizontal enlargement of the parameter space towards the right of the canonical QCD axion band are heavy axion models that solve the strong CP problem at low scales (e.g. $f_a \sim \text{TeV}$) [18–34]. The present work explores instead left horizontal shifts: true axions that solve the strong CP problem with $m_a \ll m_a^{\text{QCD}}$. This avenue is more challenging, since it requires a new source of PQ breaking aligned with QCD, whose contribution to the axion mass needs to *almost* cancel that from QCD without relying on fine-tunings.

A possible mechanism to achieve this lighter-than-usual true axion in a technically natural way was recently put forth by Hook [35], in terms of a discrete $Z_{\mathcal{N}}$ symmetry. \mathcal{N} mirror and degenerate worlds would coexist in Nature, linked by an axion field³ which implements non-linearly the $Z_{\mathcal{N}}$ symmetry. One of those worlds is our SM one. All the confining sectors contribute now to the right-hand side of Eq. (1.1), conspiring by symmetry to suppress the axion mass without spoiling the solution to the strong CP problem.

²For instance, via the measurement of the axion coupling to the neutron electric dipole moment (nEDM) operator at CASPER-electric [16, 17], in case the axion would also account for dark matter (DM). The axion-to-nEDM coupling directly follows from the m_a - f_a relation and so it is unmodified in standard approaches to axion coupling enhancements (left panel in Fig. 1) that still rely on Eq. (1.1).

³This setup for $\mathcal{N} = 2$ had previously led instead to an *enhancement* of the axion mass [36], because the axion field was assumed to be invariant under the Z_2 transformation.

The direct consequence is, for fixed f_a , a \mathcal{N} -dependent reduced axion mass in spite of all confining scales being equal to Λ_{QCD} . In other words, for a given value of m_a it follows a universal enhancement of all axion interactions relative to those of the canonical QCD axion. In this paper, we expand on the mathematical properties of the implementation of the $Z_{\mathcal{N}}$ symmetry and determine the analytic form of the exponential suppression of the axion mass and its potential in the large \mathcal{N} limit. The phenomenological analysis of the number of possible mirror worlds \mathcal{N} will be next carried out with present and projected data.

The study will also explore the $Z_{\mathcal{N}}$ axion potential at finite density, to confront present constraints and prospects from very dense stellar objects and gravitational waves. It has been recently pointed out in [37, 38] that a generic reduced-mass axion leads to strong effects on those systems, raising the effective mass in the dense media. In the scenario considered here, a stellar background made only of SM matter is by nature $Z_{\mathcal{N}}$ -asymmetric: we will show analytically how such an asymmetric background breaks the cancellations which guaranteed an exponentially suppressed axion mass for the $Z_{\mathcal{N}}$ symmetric vacuum potential. Limits on the number of possible worlds will be obtained in turn.

The theoretical framework to be used throughout the work described above is that of effective axion couplings. Nevertheless, two concrete UV completions of the $Z_{\mathcal{N}}$ scenario under consideration will be developed as well: a model à la KSVZ [7, 39], and a composite model à la Choi-Kim [40, 41]. The status of the Peccei-Quinn (PQ) quality problem will be also addressed.

An important remark is that we will consider in this paper experiments that can test the solution to the strong CP problem without further assumptions. Indeed, it is most relevant to get a clear panorama on the strong CP problem by itself, given its fundamental character. In particular, we *will not* discuss axion or ALP experiments that *do* rely on the assumption that the DM of the Universe may be constituted by axions. The cosmological evolution of the axion field in the $Z_{\mathcal{N}}$ scenario under discussion and its contribution to the DM relic abundance departs drastically from the standard case, and it is discussed in a companion paper [42].

The structure of the present paper can be easily inferred from the Table of Contents.

2 Down-tuning the axion mass

In Ref. [35] it was shown how to naturally down-tune the axion mass from its natural QCD value in Eq. (1.1), exploiting the analyticity structure of the QCD axion potential in the presence of a $Z_{\mathcal{N}}$ symmetry. For pedagogical purposes, before turning to the generic $Z_{\mathcal{N}}$ case we analyze the (unsuccessful) case of a Z_2 symmetry: the SM plus one degenerate mirror world linked by an axion which realizes the symmetry non-linearly.

2.1 The Z_2 case

Consider the SM plus a complete copy SM' , related via a Z_2 symmetry which exchanges each SM field with its mirror counterpart, while the axion field is shifted by π :

$$Z_2 : \quad \text{SM} \longleftrightarrow \text{SM}' \quad (2.1)$$

$$a \longrightarrow a + \pi f_a. \quad (2.2)$$




Figure 2: Z_2 axion potential. The mirror contribution to the axion potential $V_{SM'}(a/f_a)$ (in **green**) partially cancels that of the SM, $V_{SM}(a/f_a)$ (in **blue**), leading to a total shallower potential $V_2(a/f_a)$ (in **orange**). The total potential has a maximum in $a/f_a = 0$ and thus this Z_2 axion does not solve the SM strong CP problem.

The Lagrangian, including the anomalous effective couplings of the axion to SM fields, then reads

$$\mathcal{L} = \mathcal{L}_{SM} + \mathcal{L}_{SM'} + \frac{\alpha_s}{8\pi} \left(\frac{a}{f_a} - \theta \right) G\tilde{G} + \frac{\alpha_s}{8\pi} \left(\frac{a}{f_a} - \theta + \pi \right) G'\tilde{G}' + \dots, \quad (2.3)$$

where θ parametrizes the anomalous QCD coupling, α_s is the QCD fine-structure constant, the Lorentz indices of the field strength $G_{\mu\nu}$ have been obviated, and the dots stand for possible Z_2 -symmetric portals between the two mirror worlds (see Sect. 2.2.1). Without loss of generality, we can perform a uniform shift in a such that the θ term in Eq. (2.3) is set to zero. Therefore, the effective θ -parameter of the SM corresponds to $\theta_{\text{eff}} \equiv \langle a \rangle / f_a$, where $\langle a \rangle$ denotes the vacuum expectation value (vev) of the axion field.

In the case of an exact Z_2 symmetry, all couplings and masses of the mirror world and the SM would coincide with the exception of the effective θ -parameter. It is this difference (namely the π shift in the effective θ -parameters of the SM and its mirror) the one responsible for displaced contributions to the total axion potential, with destructive interference effects. Were the QCD axion potential to be a simple cosine, the total potential would vanish because the two contributions (from QCD and mirror QCD) would have exactly the same size but opposite sign, i.e. $\propto \cos(a/f_a)$ and $\propto \cos(a/f_a + \pi) = -\cos(a/f_a)$ respectively. However, for the true chiral axion potential [43–45] the exact cancellation disappears and a residual potential –and thus a non-zero axion mass– remains, which at leading chiral order reads (keeping only two flavours)

$$V_2(a) = -\frac{m_\pi^2 f_\pi^2}{m_u + m_d} \left\{ \sqrt{m_u^2 + m_d^2 + 2m_u m_d \cos\left(\frac{a}{f_a}\right)} + \sqrt{m_u^2 + m_d^2 - 2m_u m_d \cos\left(\frac{a}{f_a}\right)} \right\}. \quad (2.4)$$

This Z_2 -symmetric world would not solve the strong CP problem, though, because $a/f_a = 0$ is a maximum of the axion potential, as illustrated in Fig. 2. Indeed, as already pointed out in Ref. [35], $a/f_a = 0$ is a minimum of the potential only for odd values of \mathcal{N} , while it is a maximum for \mathcal{N} . Therefore, the simplest viable axion model that solves the strong CP problem with a reduced axion mass incorporates a Z_3 symmetry.

2.2 $Z_{\mathcal{N}}$ axion

We consider now \mathcal{N} copies of the SM that are interchanged under a $Z_{\mathcal{N}}$ symmetry which is non-linearly realized by the axion field:

$$Z_{\mathcal{N}} : \text{SM}_k \longrightarrow \text{SM}_{k+1 \pmod{\mathcal{N}}} \quad (2.5)$$

$$a \longrightarrow a + \frac{2\pi k}{\mathcal{N}} f_a, \quad (2.6)$$

with $k = 0, \dots, \mathcal{N} - 1$. One of those worlds will be our SM one. The most general Lagrangian implementing this symmetry describes \mathcal{N} mirror worlds whose couplings take exactly the same values as in the SM, with the exception of the effective θ -parameter: for each copy the effective θ value is shifted by $2\pi/\mathcal{N}$ with respect to that in the neighbour k sector,

$$\mathcal{L} = \sum_{k=0}^{\mathcal{N}-1} \left[\mathcal{L}_{\text{SM}_k} + \frac{\alpha_s}{8\pi} \left(\theta_a + \frac{2\pi k}{\mathcal{N}} \right) G_k \tilde{G}_k \right] + \dots \quad (2.7)$$

where $\mathcal{L}_{\text{SM}_k}$ denotes exact copies of the SM total Lagrangian excluding the strong anomalous coupling, and the dots stand for $Z_{\mathcal{N}}$ -symmetric portal couplings that may connect those different sectors (to be discussed in Sect. 2.2.1). In this equation $\theta_a \equiv a/f_a$ is the angular axion field defined in the interval $[-\pi, \pi)$, and a universal (equal for all k sectors) bare theta parameter has been set to zero via an overall shift of the axion field. The SM is identified from now on with the $k = 0$ sector: to ease the notation, the label $k = 0$ on SM quantities will be often dropped below. Each QCD_k sector contributes to the θ_a potential, which in the 2-flavour leading order chiral expansion reads

$$V_{\mathcal{N}}(\theta_a) = -A \sum_{k=0}^{\mathcal{N}-1} \sqrt{1 + z^2 + 2z \cos \left(\theta_a + \frac{2\pi k}{\mathcal{N}} \right)}, \quad (2.8)$$

where

$$z \equiv m_u/m_d \approx 0.48, \quad A \equiv \Sigma_0 m_d \approx \chi_0 (1+z)/z, \quad (2.9)$$

and

$$\Sigma_0 \equiv -\langle \bar{u}u \rangle = -\langle \bar{d}d \rangle = m_\pi^2 f_\pi^2 / (m_u + m_d) \quad (2.10)$$

denotes the chiral condensate [44], while $\chi_0 \approx (75 \text{ MeV})^4$ is the zero temperature QCD topological susceptibility [45, 46]. Alternatively, the total $Z_{\mathcal{N}}$ axion potential can be written as

$$V_{\mathcal{N}}(\theta_a) = -m_\pi^2 f_\pi^2 \sum_{k=0}^{\mathcal{N}-1} \sqrt{1 - \beta \sin^2 \left(\frac{\theta_a}{2} + \frac{\pi k}{\mathcal{N}} \right)}, \quad (2.11)$$

where $\beta \equiv 4m_u m_d / (m_u + m_d)^2 = 4z / (1+z)^2$.

For any \mathcal{N} , $\theta_a = 0$ is an extrema of the axion potential. Indeed, using the property $\sin(2\pi(\mathcal{N} - k)/\mathcal{N}) = -\sin(2\pi k/\mathcal{N})$ it is straightforward to see that

$$\left. \frac{\partial V_{\mathcal{N}}(\theta_a)}{\partial \theta_a} \right|_{\theta_a=0} = \frac{m_\pi^2 f_\pi^2 \beta}{f_a} \frac{1}{4} \sum_{k=0}^{\mathcal{N}-1} \frac{\sin\left(\frac{2\pi k}{\mathcal{N}}\right)}{\sqrt{1 - \beta \sin^2\left(\frac{\pi k}{\mathcal{N}}\right)}} = 0. \quad (2.12)$$

The same holds for any $\theta_a = 2\pi n/\mathcal{N}$ with $n \in \mathbb{Z}$, because of the periodicity of the potential. For \mathcal{N} odd the potential $V(\theta_a)$ has \mathcal{N} minima located at

$$\theta_a = \{\pm 2\pi\ell/\mathcal{N}\} \quad \text{for } \ell = 0, 1, \dots, \frac{\mathcal{N}-1}{2}, \quad (2.13)$$

which includes the origin $\theta_a = 0$, while for \mathcal{N} even the origin becomes a maximum. This result –valid for any \mathcal{N} – can be shown for instance using the exact Fourier series expansion of the potential in Eqs. (2.8)-(2.11) (see final part of Appendix C). It follows that \mathcal{N} odd is required in order to solve the SM strong CP problem (albeit with a $1/\mathcal{N}$ tuning in the cosmological evolution [35, 42]). The $k \neq 0$ worlds have instead non-zero effective θ -parameters: $\theta_k \equiv 2\pi k/\mathcal{N}$ for $\langle \theta_a \rangle = 0$, see Eq. (2.7). A typical shape of the axion potential for $\mathcal{N} = 3$ is illustrated in Fig. 3.




Figure 3: Z_3 axion potential. The contributions from the $\mathcal{N} = 3$ worlds partially cancel each other, leading to an exponentially small total potential $V_{\mathcal{N}=3}(\theta_a)$ (in blue) that exhibits a minimum in $\theta_a = 0$.

The different effective θ_k values translate into slightly different masses for the pion mass in each mirror world, $m_\pi(\theta_k)$. At quadratic order in m_π a reduction factor of up to $\sim \sqrt{3}$ results [45],

$$m_\pi^2(\theta_k) = m_\pi^2 \sqrt{1 - \frac{4m_u m_d}{(m_u + m_d)^2} \sin^2\left(\frac{\pi k}{\mathcal{N}}\right)}. \quad (2.14)$$

Interestingly, nuclear physics would be drastically different in the different mirror copies. In particular, a new scalar pion (π_k) to nucleon (N_k) coupling is generated in all worlds but the SM one (see e.g. Refs. [47, 48]):

$$\mathcal{L}_{\chi\text{PT}} \supset c_+ \sum_{k=0}^{\mathcal{N}-1} \frac{m_u m_d \sin \theta_k}{[m_u^2 + m_d^2 + 2m_u m_d \cos \theta_k]^{1/2}} \frac{\pi_k^a}{f_\pi} \bar{N}_k \tau^a N_k, \quad (2.15)$$

where c_+ is an $\mathcal{O}(1)$ low-energy constant of the baryon chiral Lagrangian. Its impact on the cosmological histories of the mirror worlds is discussed in Ref. [42] for the $Z_{\mathcal{N}}$ scenario under discussion.

Overall, for our world to be that with vanishing effective θ , the ~ 10 orders of magnitude tuning required by the SM strong CP problem has been traded by a $1/\mathcal{N}$ adjustment, while \mathcal{N} could a priori be as low as $\mathcal{N} = 3$.⁴

2.2.1 Renormalizable portals to the SM

Renormalizable portals between the SM and its mirror copies (left implicit in Eq. (2.7)) are allowed by the $Z_{\mathcal{N}}$ symmetry. In the following, we classify for completeness the portal operators connecting the different k sectors.

Higgs portals

The most general $Z_{\mathcal{N}}$ symmetric scalar potential for the Higgs doublets H_k of the different mirror worlds includes terms of the form

$$\mathcal{V}(H_k) \supset \sum_{i=1}^{(\mathcal{N}-1)/2} \kappa_i \sum_{k=0}^{\mathcal{N}-1} \left(|H_k|^2 - \frac{v^2}{2} \right) \left(|H_{k+i}|^2 - \frac{v^2}{2} \right) \Big|_{(\text{mod } \mathcal{N})}, \quad (2.16)$$

where v denotes the Higgs vev and κ_i are dimensionless parameters. Note that the $Z_{\mathcal{N}}$ -symmetric mixings between different worlds may include next-neighbour, next-to-next neighbour etc. interactions. All $\kappa_{i \geq 1}$ terms provide renormalizable portals between the mirror Higgs copies ($H_{k \neq 0}$) and the SM Higgs ($H_{k=0}$).

Kinetic mixing

Terms mixing the $U(1)_Y^k$ hypercharge field strengths of mirror worlds are *a priori* also allowed by the $Z_{\mathcal{N}}$ symmetry,

$$\mathcal{L} \supset \sum_{i=1}^{(\mathcal{N}-1)/2} \epsilon_i \sum_{k=0}^{\mathcal{N}-1} F_k^{\mu\nu} F_{\mu\nu, k+i} \Big|_{(\text{mod } \mathcal{N})}, \quad (2.17)$$

where $F_k^{\mu\nu}$ denote here the k -hypercharge field strengths and ϵ_i are free dimensionless parameters.

The above renormalizable portals are subject to strong cosmological constraints, as discussed in Ref. [42]. This can suggest a ‘naturalness’ issue for the Higgs and the kinetic

⁴Although we work in the exact $Z_{\mathcal{N}}$ limit, cosmological considerations require the temperature of the SM thermal bath to be higher than that of the other $k \neq 0$ sectors [49–51]. Mechanisms to achieve these different temperatures will be discussed in Ref. [42].

portal couplings, as they cannot be forbidden in terms of internal symmetries. Nevertheless, such small couplings may be technically natural because of an enhanced Poincaré symmetry [52, 53]: in the limit where non-renormalizable interactions are neglected, the $\kappa_{i \neq 0}$ and $\epsilon_{i \neq 0} \rightarrow 0$ limit corresponds to an enhanced $\mathcal{P}^{\mathcal{N}}$ symmetry (namely an independent space-time Poincaré transformation \mathcal{P} in each sector). Those couplings are then protected from receiving radiative corrections other than those induced by the explicit $\mathcal{P}^{\mathcal{N}}$ breaking due to gravitational and axion-mediated interactions, which are presumably small. In addition, other terms in the scalar potential which depend on the details of the UV completion of the $Z_{\mathcal{N}}$ axion scenario may be present and strongly constrained; an example is given below in Sect. 3.1.

2.3 Axion potential in the large \mathcal{N} limit

It is non-trivial to sum the series which defines the axion potential, Eq. (2.11). However, the presence of the $Z_{\mathcal{N}}$ symmetry allows for the application of powerful mathematical tools related to its Fourier decomposition and holomorphicity properties, that lead to simplified expressions in the large \mathcal{N} limit.

2.3.1 Holomorphicity bounds and convergence of Riemannian sums

As first noticed in Ref. [35], the fact that the potential in Eq. (2.11) corresponds to a Riemann sum allows one to express it as an integral plus subleading terms,

$$V_{\mathcal{N}}(\theta_a) = \sum_{k=0}^{\mathcal{N}-1} V\left(\theta_a + \frac{2\pi k}{\mathcal{N}}\right) = \frac{\mathcal{N}}{2\pi} \int_0^{2\pi} V(x) dx + \mathcal{O}(\mathcal{N}^0), \quad (2.18)$$

where the definition of each single-world potential, $V\left(\theta_a + \frac{2\pi k}{\mathcal{N}}\right)$, can be read off Eq. (2.8). Most importantly, the integral does not depend on the field θ_a and the amplitude of the axion potential is thus solely contained in the subleading terms. The latter are nothing but the error E committed in approximating the Riemann sum by an integral,

$$E_{\mathcal{N}}(V) = \int_0^{2\pi} V(x) dx - \frac{2\pi}{\mathcal{N}} \sum_{k=0}^{\mathcal{N}-1} V\left(\theta_a + \frac{2\pi k}{\mathcal{N}}\right). \quad (2.19)$$

Powerful theorems exist that describe the fast convergence of this approximation. It can be shown, applying complex analysis, that if some conditions are satisfied the convergence of the rectangular rule is exponential (see e.g. Section 3 in Ref. [54]). More precisely, if $V(\theta_a)$ is a 2π -periodic function and it can be extended to a holomorphic function $V(w)$ in a rectangle from 0 to 2π and from $-ib$ to $+ib$, then the error of the rectangular rule is constrained as

$$|E_{\mathcal{N}}(V)| \leq \frac{4\pi M}{e^{\mathcal{N}b} - 1}, \quad (2.20)$$

where M is an upper limit on $V(w)$ in the rectangular region defined above. As a consequence, the axion mass will be exponentially suppressed for large \mathcal{N} . More in detail, let us apply the theorem to the second derivative of the potential,

$$V''(\theta_a) = -\frac{m_{\pi}^2 f_{\pi}^2}{2} \frac{z}{1+z} \frac{2(1+z^2) \cos(\theta_a) + z[3 + \cos(\theta_a/2)]}{[1+z^2+2z \cos(\theta_a)]^{3/2}}, \quad (2.21)$$

which can be extended in the complex plane to a holomorphic function until the expression under the square root vanishes. Indeed, this function has branch points in⁵

$$w_{cut} = \pi \pm i \log z. \quad (2.22)$$

Naively, it is tempting to apply the theorem assuming $b = \log z$ in Eq. (2.20). This is not possible though, since $V''(w)$ is not bounded in the rectangular region, due to a divergence in the branch point. As we show in Appendix A, it is possible to optimize the bound obtained above on the axion mass ($V''(\theta_a)/f_a^2$) by allowing a departure from $\log z$, $b = \log z + \Delta b$, which leads to

$$\Delta b = \frac{3}{2} \frac{1}{\mathcal{N}}, \quad (2.23)$$

where the factor 3/2 stems from the order of the divergence of Eq. (2.21) in the branch point w_{cut} . Implementing this result in Eq. (2.20), it follows that

$$m_a^2 f_a^2 \leq |E_{\mathcal{N}}(V'')| \leq \pi m_\pi^2 f_\pi^2 \sqrt{\frac{1-z}{1+z}} \left(\frac{2}{3}\right)^{3/2} \mathcal{N}^{3/2} \frac{1}{e^{-3/2} z^{-\mathcal{N}} - 1}. \quad (2.24)$$

In Fig. 4 we compare this analytical bound with the numerical result: our analytical bound captures the correct dependence on \mathcal{N} of the $Z_{\mathcal{N}}$ axion mass,

$$m_a^2 f_a^2 \propto m_\pi^2 f_\pi^2 \sqrt{\frac{1-z}{1+z}} \mathcal{N}^{3/2} z^{\mathcal{N}}, \quad (2.25)$$

although it misses the overall constant factor. The overall factor will be analytically determined in the following. Nevertheless, the discussion above has the two-fold interest of determining the correct exponential suppression *and* of being very general, as it only relies on the holomorphicity structure of the potential, and not on the specific form it takes. As a consequence, *the exponential suppression of the axion mass is not spoiled when considering the subleading chiral corrections to Eq. (2.11).*

2.3.2 Fourier expansion: axion mass from hypergeometric functions

It is possible to gain further physical insight on the origin of the cancellations in the potential by constructing its Fourier series expansion. As shown in Appendix B, the Fourier series of any scalar potential respecting the $Z_{\mathcal{N}}$ shift symmetry only receives contributions from modes that are multiples of \mathcal{N} . Moreover, if the potential can be written as a sum of shifted contributions, as it is the case for the $Z_{\mathcal{N}}$ axion under discussion –see Eq. (2.18)– then the Fourier series of the total potential $V_{\mathcal{N}}(\theta_a)$ can be easily obtained in terms of the Fourier series of a single $V(\theta_a)$ term, leading to

$$V_{\mathcal{N}}(\theta_a) = 2\mathcal{N} \sum_{t=1}^{\infty} \hat{V}(t\mathcal{N}) \cos(t\mathcal{N}\theta_a), \quad (2.26)$$

where $\hat{V}(n)$ denotes the coefficient of the Fourier series for the single-world potential $V(\theta_a)$,

$$\hat{V}(n) = -\frac{m_\pi^2 f_\pi^2}{1+z} \int_0^{2\pi} \cos(nt) \sqrt{1+z^2+2z \cos(t)} dt. \quad (2.27)$$

⁵This result coincides with that in Ref. [35], which defines $a = \log(c + \sqrt{c^2 - 1})$, for $c = \frac{(m_u + m_d)^2}{2m_u m_d} - 1$. Note that the variable a can be simplified as $a = \log(m_d/m_u) = -\log z$.

It is convenient to express this integral in terms of the Gauss hypergeometric function (see Appendix C and Ref. [55] for conventions and relevant properties),

$$\hat{V}(n) = (-1)^{n+1} \frac{m_\pi^2 f_\pi^2}{1+z} z^n \frac{\Gamma(n-1/2)}{\Gamma(-1/2) n!} {}_2F_1 \left(\begin{matrix} -1/2, n-1/2 \\ n+1 \end{matrix} \middle| z^2 \right). \quad (2.28)$$

As shown in Appendix C, in the large \mathcal{N} limit this expression further simplifies to

$$\hat{V}(n) \simeq (-1)^n \frac{m_\pi^2 f_\pi^2}{2\sqrt{\pi}} \sqrt{\frac{1-z}{1+z}} n^{-3/2} z^n, \quad (2.29)$$

leading to the following expression for the total potential

$$\begin{aligned} V_{\mathcal{N}}(\theta_a) &\simeq \frac{m_\pi^2 f_\pi^2}{\sqrt{\pi}} \sqrt{\frac{1-z}{1+z}} \mathcal{N}^{-1/2} \sum_{t=1}^{\infty} (-1)^{t\mathcal{N}} t^{-3/2} z^{t\mathcal{N}} \cos(t\mathcal{N}\theta_a) \\ &\simeq \frac{m_\pi^2 f_\pi^2}{\sqrt{\pi}} \sqrt{\frac{1-z}{1+z}} \mathcal{N}^{-1/2} (-1)^{\mathcal{N}} z^{\mathcal{N}} \cos(\mathcal{N}\theta_a), \end{aligned} \quad (2.30)$$

where in the second line we have kept only the first mode in the expansion, as the higher modes are exponentially suppressed with respect to it. *The total potential is thus safely approximated by a single cosine.* It trivially follows from Eq. (2.30) that $\theta_a = 0$ is a minimum of the total potential for \mathcal{N} odd, and a maximum for \mathcal{N} even. Here and all through this work purely constant terms in the potential are obviated, as they have no impact on the axion mass.

Eq. (2.30) can be rewritten as

$$V_{\mathcal{N}}(\theta_a) \simeq -\frac{m_a^2 f_a^2}{\mathcal{N}^2} \cos(\mathcal{N}\theta_a), \quad (2.31)$$

where the $Z_{\mathcal{N}}$ axion mass m_a in the large \mathcal{N} limit is finally given by a compact and analytical formula,

$$m_a^2 f_a^2 \simeq \frac{m_\pi^2 f_\pi^2}{\sqrt{\pi}} \sqrt{\frac{1-z}{1+z}} \mathcal{N}^{3/2} z^{\mathcal{N}}. \quad (2.32)$$

The overall coefficient is thus determined, in addition to exhibiting the $z^{\mathcal{N}}$ exponential suppression of the potential and the specific \mathcal{N} dependence previously argued in Eq. (2.25) from holomorphicity arguments. In summary, in the large \mathcal{N} limit the axion mass is reduced with respect to that of the QCD axion by a factor

$$\left(\frac{m_a}{m_a^{\text{QCD}}} \right)^2 \simeq \frac{1}{\sqrt{\pi}} \sqrt{1-z^2} (1+z) \mathcal{N}^{3/2} z^{\mathcal{N}-1}, \quad (2.33)$$

where m_a^{QCD} denotes the mass of the canonical QCD axion as given in Eq. (1.1).⁶ This ratio is illustrated in Fig. 4, which compares the numerical behaviour with: *a)* the analytical dependence previously proposed in Ref. [35]; *b)* that from the holomorphicity bound in Eq. (2.24); *c)* the full analytical result in Eq. (2.33). Our analytical results improve on previous ones by Hook on a number of aspects: *i)* the explicit determination of the exponential behavior controlled by $z^{\mathcal{N}} \sim 2^{-\mathcal{N}}$; *ii)* the improved \mathcal{N} dependence from the factor $\mathcal{N}^{3/2}$; *iii)* the z -dependence of the axion mass in $\sqrt{\frac{1-z}{1+z}}$; *iv)* the determination of the prefactor $1/\sqrt{\pi}$.

In practice, the large \mathcal{N} results in Eqs. (2.30)-(2.33) turn out to be an excellent approximation already for $\mathcal{N} = 3$.

⁶Note that, although $\mathcal{N} = 1$ denotes the SM world, m_a^{QCD} does not correspond to $\mathcal{N} = 1$ in Eq. (2.32), because the latter is only valid in the large \mathcal{N} limit.




Figure 4: Comparison between different evaluations of the axion mass as a function of \mathcal{N} . Our large- \mathcal{N} analytical result in Eq. (2.33) (green curve) provides a remarkably good approximation to the numerical evaluation (dots).

3 UV completions and alternative scenarios

Up to this point, the analysis has been largely independent from the precise UV completion of the $Z_{\mathcal{N}}$ axion scenario. For the sake of illustration, in this section we provide two UV completions of the axion effective Lagrangian in Eq. (2.7). We also briefly discuss an alternative implementation of the $Z_{\mathcal{N}}$ symmetry in which the resulting axion is heavier than usual (rather than lighter).

3.1 KSVZ $Z_{\mathcal{N}}$ axion

Consider \mathcal{N} copies of vector-like Dirac fermions \mathcal{Q}_k ($k = 0, \dots, \mathcal{N} - 1$) transforming in the fundamental representation of QCD_k , together with a gauge singlet complex scalar \mathcal{S} . The action of the $Z_{\mathcal{N}}$ symmetry on these fields is postulated to be

$$Z_{\mathcal{N}} : \mathcal{Q}_k \rightarrow \mathcal{Q}_{k+1 \pmod{\mathcal{N}}}, \quad (3.1)$$

$$\mathcal{S} \rightarrow e^{2\pi i/\mathcal{N}} \mathcal{S}, \quad (3.2)$$

while the SM Lagrangian and its copies obey Eq. (2.5) under $Z_{\mathcal{N}}$. The most general Lagrangian containing the new degrees of freedom then reads

$$\mathcal{L}_{\text{UV}} = |\partial_{\mu} \mathcal{S}|^2 + \sum_{k=0}^{\mathcal{N}-1} \left[\bar{\mathcal{Q}}_k i \not{D} \mathcal{Q}_k + y e^{2\pi i k/\mathcal{N}} \mathcal{S} \bar{\mathcal{Q}}_k P_R \mathcal{Q}_k + \text{h.c.} \right] - \mathcal{V}(\mathcal{S}, H_k), \quad (3.3)$$

where $P_R \equiv (1 + \gamma_5)/2$. It exhibits an accidental $\text{U}(1)_{\text{PQ}}$ symmetry

$$\text{U}(1)_{\text{PQ}} : \mathcal{Q}_k \rightarrow e^{-i\gamma_5 \frac{\alpha}{2}} \mathcal{Q}_k, \quad (3.4)$$

$$\mathcal{S} \rightarrow e^{i\alpha} \mathcal{S}, \quad (3.5)$$

that is spontaneously broken by the vev of \mathcal{S} , v_S , via a proper ‘mexican-hat’ potential $\mathcal{V}(\mathcal{S}, H_k)$, whose structure is discussed below. Decomposing the \mathcal{S} field in a polar basis,

$$\mathcal{S} = \frac{1}{\sqrt{2}}(v_S + \rho)e^{i\frac{\alpha}{v_S}}, \quad (3.6)$$

in terms of canonically normalized radial (ρ) and axion modes, the latter can be rotated away from the Yukawa term in Eq. (3.3) via an axion-dependent axial transformation

$$\mathcal{Q}_k \rightarrow e^{-i\gamma_5\left(\frac{a}{2v_S} + \frac{\pi k}{N}\right)} \mathcal{Q}_k. \quad (3.7)$$

The heavy quarks, with real mass⁷ $m_{\mathcal{Q}_k} = \frac{y_S v_S}{\sqrt{2}}$, can next be integrated out in order to obtain the low-energy axion effective field theory. Because the transformation in Eq. (3.7) is QCD_k anomalous, with anomaly factor $2N_k = 1$, the resulting axion effective Lagrangian is given by

$$\delta\mathcal{L}_{\text{UV}} = \sum_{k=0}^{N-1} \frac{\alpha_s}{8\pi} \left(\frac{a}{v_S} + \frac{2\pi k}{N} \right) G_k \tilde{G}_k, \quad (3.8)$$

which yields precisely Eq. (2.7), after the identification $v_S = f_a$.

Furthermore, the presence of the singlet scalar \mathcal{S} introduces new scalar portals between the SM and its mirror worlds, in addition to the generic ones in Eq. (2.16). The scalar potential in the latter equation should thus be enlarged by

$$\mathcal{V}(H_k) \longrightarrow \mathcal{V}(\mathcal{S}, H_k) = \mathcal{V}(H_k) + \delta\mathcal{V}, \quad (3.9)$$

with

$$\delta\mathcal{V} = \lambda_S \left(|\mathcal{S}|^2 - \frac{f_a^2}{2} \right)^2 + \kappa_S \left(|\mathcal{S}|^2 - \frac{f_a^2}{2} \right) \sum_{k=0}^{N-1} \left(|H_k|^2 - \frac{v^2}{2} \right). \quad (3.10)$$

Note that, because the Higgs vev v is the same in all k sectors due to the unbroken Z_N symmetry, the required hierarchy of scales is obtained with a single fine-tuning between v and f_a , as for elementary canonical QCD axions.

It is also possible to choose the representations of the \mathcal{Q}_k fields to transform non-trivially under the electroweak_k gauge groups, so that they could e.g. mix with SM_k quarks in a Z_N invariant way and decay efficiently in the early Universe, thus avoiding possible issues with colored/charged stable relics in the SM sector [9, 11]. Depending on the \mathcal{Q}_k quantum numbers, this would change in turn the value of the electromagnetic-to-QCD anomaly ratio of the PQ current, usually denoted as E/N , which enters the axion-photon coupling.

3.1.1 Peccei-Quinn quality

The threat posed on traditional QCD axion models by quantum non-perturbative gravitational corrections [56–65] may also affect the models discussed here, as f_a is not very far from the Planck scale. These contributions are usually parametrized via effective operators, suppressed by powers of the Planck mass, that could explicitly violate the PQ symmetry and thus spoil the solution to the strong CP problem [56–59].⁸

⁷Note that we crucially removed also the k -dependent phases from the Yukawas, in order to properly integrate out the heavy \mathcal{Q}_k fields.

⁸UV sources of PQ breaking can be avoided in some invisible axion constructions within a variety of extra assumptions or frameworks [66–78], or be arguably negligible under certain conditions [79].




Figure 5: Parameter space in the $\{\mathcal{N}, f_a\}$ plane that is free from the PQ quality problem, within the KSVZ-like UV completion of the reduced-mass $Z_{\mathcal{N}}$ axion, for the PQ-breaking parameter values indicated in the text. The PQ protected region has a sizable overlap with the regions of parameter space where the $Z_{\mathcal{N}}$ axion can account for the total DM relic density, see Ref. [42].

In the context of the KSVZ $Z_{\mathcal{N}}$ axion model above, the exponentially small axion mass could seem to worsen this threat, increasing the sensitivity to explicit PQ-breaking effective operators. Interestingly, promoting the *in built* $Z_{\mathcal{N}}$ symmetry to a gauge symmetry leads to an accidental $U(1)_{\text{PQ}}$ invariance, that for large \mathcal{N} is efficiently protected from those extra sources of explicit breaking. Indeed, the lowest-dimensional PQ-violating operator in the scalar potential compatible with the $Z_{\mathcal{N}}$ symmetry is $\mathcal{S}^{\mathcal{N}}$, leading to an explicitly PQ-breaking contribution to the potential of the form

$$V_{\text{PQ-break.}} = c \frac{\mathcal{S}^{\mathcal{N}}}{M_{\text{Pl}}^{\mathcal{N}-4}} + \text{h.c.} \supset \frac{|c|}{2^{\mathcal{N}/2-1}} \frac{f_a^{\mathcal{N}}}{M_{\text{Pl}}^{\mathcal{N}-4}} \cos(\mathcal{N}\theta_a + \delta), \quad (3.11)$$

where $M_{\text{Pl}} = 1.22 \times 10^{19}$ GeV is the Planck mass and c is a dimensionless coefficient with phase $\delta \equiv \text{Arg } c$. Considering now $V_{\mathcal{N}}(\theta_a) + V_{\text{PQ-break.}}$, expanding for small θ_a the axion potential $V_{\mathcal{N}}(\theta_a) \approx V_{\mathcal{N}}(0) + \frac{1}{2} m_a^2 f_a^2 \theta_a^2$, and solving the tadpole equation, the induced effective θ parameter in the SM sector reads

$$\begin{aligned} \langle \theta_a \rangle &\simeq \frac{|c| \mathcal{N} f_a^{\mathcal{N}} M_{\text{Pl}}^4 \sin \delta}{2^{\mathcal{N}/2-1} m_a^2 f_a^2 M_{\text{Pl}}^{\mathcal{N}} - |c| \mathcal{N}^2 f_a^{\mathcal{N}} M_{\text{Pl}}^4 \cos \delta} \\ &\simeq 2\sqrt{\pi} |c| \sin \delta \sqrt{\frac{1+z}{1-z}} \frac{M_{\text{Pl}}^4}{m_{\pi}^2 f_{\pi}^2} \frac{1}{\sqrt{\mathcal{N}}} \left(\frac{f_a}{\sqrt{2} z M_{\text{Pl}}} \right)^{\mathcal{N}}, \end{aligned} \quad (3.12)$$

where m_a^2 from Eq. (2.32) has been used, and in the last step we neglected the second term in the denominator in the first line of Eq. (3.12): this is always justified in the $\langle \theta_a \rangle \lesssim 10^{-10}$ regime.

In summary, unlike the customary ad-hoc $Z_{\mathcal{N}}$ protection mechanism for the standard KSVZ axion, in the $Z_{\mathcal{N}}$ axion scenario under discussion the discrete symmetry is already present by construction. Note that the scaling with \mathcal{N} is slightly different as compared to the standard KSVZ axion, due to the enhancement factor $1/z^{\mathcal{N}}$. But eventually the

$(f_a/M_{\text{Pl}})^{\mathcal{N}}$ suppression dominates and provides an efficient protection mechanism, even though the axion mass is exponentially suppressed. For the sake of an estimate, Fig. 5 shows the regions in the $\{\mathcal{N}, f_a\}$ plane that saturate the nEDM bound for $|c| = 1$ and $\sin \delta = 1$.

3.2 Composite $Z_{\mathcal{N}}$ axion

It is also possible to construct a UV completion of the $Z_{\mathcal{N}}$ scenario which corresponds to a dynamical (composite) axion *à la* Kim-Choi [40, 41], without extending its exotic fermionic content. In the original version of that model, the SM fields are not charged under the PQ symmetry while two exotic *massless* quarks, ψ and χ , transform under an extra confining “axi-color” group $\text{SU}(\tilde{N})_a$ and one of them, ψ , is also a triplet of QCD. Upon confinement of the axi-color group at a large scale $\Lambda_a \sim f_a \gg \Lambda_{\text{QCD}}$, pseudo-Goldstone bosons composed of the exotic quarks emerge. All but one of them are coloured under QCD and become safely heavy. The light remaining one is the composite axion, whose mass obeys the usual formula for QCD axions Eq. (1.1).

We implement the Kim-Choi idea in the framework of our $Z_{\mathcal{N}}$ framework without increasing the number of massless exotic fermions representations. The fermion ψ is simply extended to be now a triplet under *all* QCD_k mirror sectors, see Table 1. The axion field will thus be unique and will couple to all anomalous terms.

	$\text{SU}(\tilde{N})_a$	$\text{SU}(3)_{c,0}$	\dots	$\text{SU}(3)_{c,k}$	\dots	$\text{SU}(3)_{c,\mathcal{N}}$
ψ	$\mathbf{3}$	$\mathbf{3}$	\dots	$\mathbf{3}$	\dots	$\mathbf{3}$
χ	$\mathbf{1}$	$\mathbf{1}$	\dots	$\mathbf{1}$	\dots	$\mathbf{1}$

(3.13)

Table 1: Exotic fermionic sector of the $Z_{\mathcal{N}}$ composite axion model.

Upon $\text{SU}(\tilde{N})_a$ confinement at the large scale of order f_a , the QCD_k couplings α_s^k can be neglected, and therefore a large global flavor symmetry arises in the exotic fermionic sector: $\text{SU}(3^{\mathcal{N}} + 1)_L \times \text{SU}(3^{\mathcal{N}} + 1)_R \times \text{U}(1)_V$.⁹ This symmetry is spontaneously broken down to $\text{SU}(3^{\mathcal{N}} + 1)_{L+R} \times \text{U}(1)_V$ by the exotic fermion condensates. Among the resulting Goldstone bosons, the QCD_k singlet corresponds to the composite axion. Its associated PQ current reads (with $f_{\text{PQ}} \equiv \tilde{N} f_a$)

$$j_{\text{PQ}}^\mu = \bar{\psi} \gamma^\mu \gamma^5 \psi - 3^{\mathcal{N}} \bar{\chi} \gamma^\mu \gamma^5 \chi \equiv f_{\text{PQ}} \partial^\mu a, \quad (3.14)$$

which corresponds to the only element of the Cartan sub-algebra of $\text{SU}(3^{\mathcal{N}} + 1)$ that has a vanishing anomaly coefficient with respect to $\text{SU}(\tilde{N})_a$, but a non-vanishing one with respect to all the QCD_k gauge groups.

Without further elements the model would be viable, but all mirror worlds would have the same θ -parameter: a heavier than usual axion would result. A simple $Z_{\mathcal{N}}$ implementation which leads instead to relatively shifted potentials, and thus to a reduced axion mass, is to have a relative phase between the argument of the determinant of the quark mass matrix of the mirror worlds,

$$\arg(\det(Y_u Y_d))_{k+1} = \arg(\det(Y_u Y_d))_k + \frac{2\pi}{\mathcal{N}}, \quad (3.15)$$

⁹The $\text{U}(1)_A$ of the exotic sector is explicitly broken by the $\text{SU}(\tilde{N})_a$ anomaly.

where Y_u and Y_d denote the Yukawa matrices for the up and down quark sectors, respectively. One of the many possible $Z_{\mathcal{N}}$ charge assignments for the quarks that yield Eq. (3.15) is that in which only the right-handed up quarks would transform as¹⁰

$$Z_{\mathcal{N}} : U_R^k \rightarrow e^{i2\pi/(3\mathcal{N})} U_R^{k+1}, \quad (3.16)$$

corresponding to a Yukawa quark Lagrangian of the form

$$\mathcal{L}_Y = - \sum_{k=0}^{\mathcal{N}-1} \{ e^{i2\pi k/(3\mathcal{N})} \bar{Q}_L Y_u \tilde{H} U_R + \bar{Q}_L Y_d H D_R \}_k + \text{h.c.} \quad (3.17)$$

The resulting low-energy axion effective field theory is then the desired one as in Eq. (2.7).

In this $Z_{\mathcal{N}}$ composite axion model only the exotic fermions are charged under the PQ symmetry, while the $Z_{\mathcal{N}}$ charges are carried solely by SM quarks. This means that the $Z_{\mathcal{N}}$ and PQ symmetries are not directly linked. As a consequence, gauging the in-built $Z_{\mathcal{N}}$ symmetry would not soften the PQ quality problem, contrary to the case of the KSVZ $Z_{\mathcal{N}}$ axion model discussed earlier above. Our $Z_{\mathcal{N}}$ composite axion model is then subject to the usual PQ quality threat. Standard softening solutions often applied to composite axion models could be explored, e.g. those based on a chiral gauging of the global symmetry of the coset space or on introducing a moose structure [66, 69, 73, 80].

3.3 Ultra-light QCD axions

The term ultra-light axions usually refers to the mass range $m_a \in [10^{-33}, 10^{-10}]$ eV (with the extrema of the interval corresponding respectively to an axion Compton wavelength of the size of the Hubble horizon and to the Schwarzschild radius of a stellar mass black hole). As a theoretical motivation for ultra-light axions, the so-called string Axiverse [81] is often invoked, according to which a plenitude of ultra-light axions populating mass regions down to the Hubble scale 10^{-33} eV is a generic prediction of String Theory, although without a direct reference to the solution of the strong CP problem.¹¹ On the other hand, according to the usual QCD mass vs. f_a relation, Eq. (1.1), axion masses below the peV correspond to axion decay constants larger than the Planck mass, and hence they are never entertained within canonical QCD axion models. The $Z_{\mathcal{N}}$ axion framework discussed in the present work allows in contrast to populate the sub-peV axion mass region while keeping sub-Planckian axion decay constants, with the advantage of providing as well a direct solution to the strong CP problem. As shown in Sec. 4.2, the tantalizing prospects for testing the $Z_{\mathcal{N}}$ scenario, through observational data on very dense stellar objects and gravitational waves, can sweep through the discovery region of the ultra-light axion range.

3.4 A heavier-than-QCD axion

A remark is in order regarding the $Z_{\mathcal{N}}$ charge of the axion in the different sectors. If the implementation of the $Z_{\mathcal{N}}$ symmetry would be such that the \mathcal{N} world replicas are

¹⁰Note that a factor of 1/3 in the phase takes into account that there are 3 fermion families.

¹¹See e.g. Ref. [82] for an ultra-light scalar field whose mass is protected by a discrete $Z_{\mathcal{N}}$ symmetry but does not solve the strong CP problem.

degenerate but the axion field is exactly the same in all of them, that is, if Eqs. (2.5)–(2.6) were replaced by

$$Z_{\mathcal{N}} : \quad \text{SM}_k \longrightarrow \text{SM}_{k+1 \pmod{\mathcal{N}}} \quad (3.18)$$

$$a \longrightarrow a, \quad (3.19)$$

the potentials of the different mirror worlds would not be relatively shifted but exactly superpose. The axion would then be a factor $\sqrt{\mathcal{N}}$ heavier than the usual QCD axion in Eq. (1.1). This scenario was proposed in Ref. [36] for a Z_2 symmetry with just one mirror world degenerate with the SM, but its generalization to \mathcal{N} copies is trivial. Such a *heavier-than-QCD* axion solution is viable, and it would transform the ALP arena to the *right* of the canonical QCD axion band into solutions to the SM strong CP problem. The axion $Z_{\mathcal{N}}$ charge assignment explored throughout this work, Eq. (2.6), results instead in *lighter-than-QCD* axions, that is, solutions located to the left of the QCD axion band. Note that this option induces a comparatively much larger impact: a natural *exponential suppression* of the axion mass $\propto z^{\mathcal{N}}$ as the byproduct of the cancellations between the mirror potentials, Eq. (2.32), instead of the mild $\sqrt{\mathcal{N}}$ enhancement just discussed.

All in all, to explore the right-hand side region of the QCD axion band for solutions to the strong CP problem, other heavy axion scenarios proposed in the literature seem more efficient and appealing (e.g. those with mirror worlds much heavier than the SM, or scenarios with novel confining scales much larger than Λ_{QCD} , as mentioned in Sect. 1).

4 Experimental probes of down-tuned axions

The $Z_{\mathcal{N}}$ axion with reduced mass can provide a solution to the SM strong CP problem, independently of whether it accounts or not for the DM content of the Universe. It is hence interesting to get a perspective on the experimental panorama that *does not* rely on the supplementary assumption that the axion may be the DM particle: all experimental bounds and prospects below will be independent of that hypothesis. On the other hand, Ref. [42] will focus on experimental probes that do rely on it.

4.1 Axion coupling to photons

From an experimental point of view, a highly relevant axion coupling is that to photons, defined via the Lagrangian term $\delta\mathcal{L} = \frac{1}{4}g_{a\gamma}aF\tilde{F}$ as [45, 103]

$$g_{a\gamma} = \frac{\alpha}{2\pi f_a}(E/N - 1.92(4)), \quad (4.1)$$

where E and N denote model-dependent anomalous electromagnetic and strong contributions, respectively. Fig. 6 shows the parameter space of the reference $Z_{\mathcal{N}}$ axion model (with $E/N = 0$) in the coupling vs. mass plane. Predictions for the axion photon coupling are obtained by rescaling the $Z_{\mathcal{N}}$ axion mass in Eq. (2.32) for different values of \mathcal{N} . Present axion limits and projected sensitivities are displayed as filled and transparent areas, respectively.

The yellow band depicts the canonical QCD axion solution, which obeys the well-known relation in Eq. (1.1). The oblique lines indicate instead the $Z_{\mathcal{N}}$ lighter axion solutions to the strong CP problem, as a function of the number of mirror worlds \mathcal{N} , see Eq. (2.32). Note that *the overall effect of a reduced mass axion is simply a shift towards*




Figure 6: Limits on the axion-photon coupling as a function of the axion mass. Laboratory constrains [83–90] and astrophysical bounds [91–101] are shown in **blue** and **green**, respectively. Projected sensitivities appear in translucent colors delimited by dashed lines. The **orange** oblique lines represent the theoretical prediction for the $Z_{\mathcal{N}}$ axion photon couplings assuming $E/N = 0$ for different (odd) number of worlds \mathcal{N} . These lines are solid for the regions of the parameter space in which the KSVZ UV completion of the $Z_{\mathcal{N}}$ axion is free from PQ quality problem and dashed otherwise. The secondary vertical axis shows the corresponding axion decay constant f_a if $E/N = 0$ is assumed. Supplementary constraints in case the axion is assumed to account for DM can be found in Ref. [42]. Axion limits adapted from Ref. [102].

the left of the parameter space: each of those oblique lines can be considered to be the center of a displaced yellow band. It is particularly enticing that experiments set *a priori* to only hunt for ALPs may in fact be targeting solutions to the strong CP problem.

4.2 Finite density constraints on f_a

This subsection summarizes the model-independent constraints on f_a for the $Z_{\mathcal{N}}$ scenario under discussion. The result of the analysis is illustrated in Fig. 8. Interestingly, apart from the usual constraints stemming from the SN1987A [104] and black hole superradiance measurements [105–108] (depicted in purple), novel bounds apply to the exceptionally light $Z_{\mathcal{N}}$ axion due to finite density effects. Indeed, it has been recently pointed out in Refs. [37, 38] that finite density media may have a strong impact on the physics of very light axions or ALPs. In those media, the minimum of the total potential

may be shifted to π . This has a number of phenomenological consequences that span from the modification of the nuclear processes in stellar objects due to $\theta \sim \mathcal{O}(1)$, to modifications in the orbital decay of binary systems (and subsequently in the emitted gravitational waves).

For the scenario considered here, the important point is that a background made only of ordinary matter breaks the $Z_{\mathcal{N}}$ symmetry. This hampers the symmetry-induced cancellations in the potential which led to a reduced-mass axion in vacuum: the effective axion mass will be larger within a dense medium.

We will first elaborate on the $Z_{\mathcal{N}}$ axion potential in a nuclear medium. Following Refs. [37, 109], one can compute the finite density effects on the axion potential by considering the quark condensates in a medium made of non-relativistic neutrons and protons [110, 111]. Applying the Hellmann-Feynman theorem, the quark condensate at a finite density n_N of a given nucleon N can be expressed as

$$\langle \bar{q}q \rangle_{n_N} = \langle \bar{q}q \rangle_0 \left(1 - \frac{\sigma_N n_N}{m_\pi^2 f_\pi^2} \right), \quad (4.2)$$

where $\langle \bar{q}q \rangle_0 = \frac{1}{2} (\langle \bar{u}u \rangle + \langle \bar{d}d \rangle) \equiv -\Sigma_0$ is the quark condensate in vacuum –see Eq. (2.10)– and σ_N is defined by

$$\sigma_N \equiv m_q \frac{\partial M_N}{\partial m_q}, \quad (4.3)$$

where $m_q \equiv \frac{1}{2}(m_u + m_d)$ and M_N is the mass of the nucleon N . Because the $Z_{\mathcal{N}}$ potential




Figure 7: Example of the in-medium potential dependence as a function of the nuclear density for $\mathcal{N} = 5$. For large densities (light green) the total potential develops a minimum in $\theta_a \sim \pi$.

is proportional to the quark condensate, see Eq. (2.8), we can simply obtain the potential within a SM nuclear medium $V_{\mathcal{N}}^{f.d.}(\theta_a, n_N)$ by weighting the SM (i.e. $k = 0$) contribution in the vacuum potential by the factor in Eq. (4.2), that is,

$$V_{\mathcal{N}}^{f.d.}(\theta_a, n_N) \simeq \left(1 - \frac{\sigma_N n_N}{m_\pi^2 f_\pi^2} \right) V(\theta_a) + \sum_{k=1}^{\mathcal{N}-1} V(\theta_a + 2\pi k/\mathcal{N}) \quad (4.4)$$

$$= -\frac{\sigma_N n_N}{m_\pi^2 f_\pi^2} V(\theta_a) + \sum_{k=0}^{\mathcal{N}-1} V(\theta_a + 2\pi k/\mathcal{N}) \xrightarrow{\mathcal{N} \gg 1} -\frac{\sigma_N n_N}{m_\pi^2 f_\pi^2} V(\theta_a).$$

In the last step of these expressions the large \mathcal{N} limit has been taken, which allowed us to neglect the term corresponding to the exponentially reduced axion potential in vacuum (see Eq. 2.30). This shows that, if the nucleon density is large enough, the $Z_{\mathcal{N}}$ asymmetric background spoils the cancellations among the mirror world contributions to the potential, in such a way that the total potential in matter is proportional to *minus* the SM one in vacuum $V(\theta_a)$. Therefore, the minimum of the potential is located at $\theta_a = \pi$. More precisely,

$$V_{\mathcal{N}}^{f.d.}(\theta_a, n_N) \xrightarrow{\mathcal{N} \gg 1} \frac{m_\pi^2 f_\pi^2}{1+z} \left[\frac{\sigma_N n_N}{m_\pi^2 f_\pi^2} \sqrt{1+z^2+2z \cos(\theta_a)} - \frac{\mathcal{N}^{-1/2} z^\mathcal{N}}{\sqrt{\pi}} \sqrt{1-z^2} \cos(\mathcal{N}\theta_a) \right], \quad (4.5)$$

which requires

$$\frac{\sigma_N n_N}{m_\pi^2 f_\pi^2} \gg z^\mathcal{N} \quad (4.6)$$

for the minimum to sit at $\theta_a = \pi$. This is illustrated in Fig. 7.

A large value of the θ parameter inside dense stellar objects is rich in physical consequences, which translates into strong constraints for the $Z_{\mathcal{N}}$ scenario. As it was pointed out in Ref. [37], $\theta \sim \mathcal{O}(1)$ inside the solar core is excluded due to the increased proton-neutron mass difference (which would prohibit the neutrino line corresponding to the Be^7 - Li^7 mass difference observed by Borexino [112]). Similarly, for $\theta \sim \pi$ in nearby neutron stars (NS), Co^{56} would be lighter than Fe^{56} [47, 48] and therefore Fe^{56} could have been depleted due to its β -decay to Co^{56} . The presence of iron in the surface of neutron stars and its implications in terms of the allowed θ values could be explored through dedicated X-ray measurements [113]. The corresponding current and projected constraints that were derived in Ref. [37] (within the simplifying assumption $z = 1$) are translated here to the $Z_{\mathcal{N}}$ scenario and further generalized for any z .

A conservative criterion consistent with $\theta = \pi$ inside the medium is to impose that the axion mass at $\theta_a = 0$ becomes tachyonic, i.e. $-m_T^2 > 0$ where m_T^2 is defined by

$$-m_T^2 \equiv \left. \frac{d^2 V_{\mathcal{N}}^{f.d.}}{d^2 a} \right|_{\theta_a=0} = \frac{m_\pi^2 f_\pi^2}{f_a^2} \left[\frac{1}{\sqrt{\pi}} \sqrt{\frac{1-z}{1+z}} \mathcal{N}^{3/2} z^\mathcal{N} - \frac{\sigma_N n_N}{m_\pi^2 f_\pi^2} \frac{z}{(1+z)^2} \right]. \quad (4.7)$$

Requiring this quantity to be positive, it directly follows a limit on the number of worlds allowed by the stellar bounds above:

$$\mathcal{N} \lesssim 47, \quad (4.8)$$

where the most recent estimation of σ_N has been used.¹² This bound does not apply for the whole range of f_a , though, because the argument only makes physical sense as long as the reduced Compton wavelength of the axion is smaller than the stellar object, $r_{\text{core}} \gtrsim 1/m_a^{f.d.}$, where $m_a^{f.d.} \sim 1/f_a$ is the effective axion mass in the medium,

$$(m_a^{f.d.})^2 = \left. \frac{d^2 V_{\mathcal{N}}^{f.d.}}{d^2 a} \right|_{\theta_a=\pi} = \frac{m_\pi^2 f_\pi^2}{f_a^2} \left[\frac{\sigma_N n_N}{m_\pi^2 f_\pi^2} \frac{z}{1-z^2} - \frac{1}{\sqrt{\pi}} \sqrt{\frac{1-z}{1+z}} \mathcal{N}^{3/2} z^\mathcal{N} \right]. \quad (4.9)$$

¹²We employ here $\sigma_N \simeq 59 \text{ MeV}$ which is in agreement with recent determinations based on Roy-Steiner equations $\sigma_N = 59.1(3.5) \text{ MeV}$ [114] and ChPT estimates $\sigma_N = 59(7) \text{ MeV}$ [115].

For the case of the sun, $r_{core} \sim 139.000$ km implies $f_a \lesssim 2.4 \times 10^{15}$ GeV for the observational constraints to apply. Finally, the area in parameter space excluded is depicted in dark blue in Fig. 8. Analogously, the future sensitivity prospects from neutron star data are depicted in shaded blue.¹³




Figure 8: Model-independent constraints on the axion scale f_a versus axion mass, from astrophysical data. Regions presently excluded are depicted in solid colors, while the translucent regions circumscribed by dotted lines are projections. The **orange** oblique lines indicate the theoretical prediction of the reduced-mass $Z_{\mathcal{N}}$ QCD axion scenario, as a function of \mathcal{N} : they are solid where the KSVZ $Z_{\mathcal{N}}$ axion is free from the PQ quality problem, and dashed otherwise. Additional constraints which apply if the axion is assumed to account as well for DM are discussed in Ref. [42].

Even stronger bounds may be established by relaxing the requirement stemming from Eq. (4.7). Indeed, as it can be seen in Fig. 8, long before the mass in $\theta_a = 0$ becomes tachyonic, the absolute minimum of the potential corresponds to $\theta \sim \mathcal{O}(1)$. Therefore one could constrain larger masses or smaller \mathcal{N} values in the $Z_{\mathcal{N}}$ scenario than those obtained above. This would require, however, a dedicated analysis to ensure that the axion field would fall into the absolute minimum, so as to overcome the potential barrier; this development lies beyond the scope of the present work.

The fact that the position of the minimum of the axion potential depends on the nuclear density of the medium not only modifies the effective θ -parameter inside stellar

¹³Our results are analogous to those in Eq. (1.7) of Ref. [37], with their generic parameter ϵ identified as $\epsilon = m_a^2/m_a^2(\mathcal{N} = 1) \simeq \pi^{-1/2} \sqrt{1 - z^2} (1 + z) \mathcal{N}^{3/2} z^{\mathcal{N}-1}$. Note that the location of the QCD axion line, as well as our projected exclusion regions for neutron stars and gravitational waves, are shifted towards the left by a factor of five with respect to those in Refs. [37, 38].

objects but may also source a long-range force between them. This has been studied in Refs. [37, 38]. This new long-range force sourced by the axion can be constrained by the measurement of double pulsar or neutron star (NS) - pulsar binaries [116–118]. Moreover, the existence of these axionic long-range forces would modify the gravitational wave signal emitted by NS-NS mergers or black hole-NS mergers which will be probed in the future by LIGO/VIRGO and Advanced LIGO [38, 119]. The projected constraints from Ref. [38] are shown in green in Fig. 8. It is striking that the whole ultra-light DM region, including the so-called “fuzzy dark matter” region ($m_a \sim 10^{-22}$ eV) [120], will be within observational reach in the next decades, for a wide range of \mathcal{N} values.

5 Conclusions

An axion which solves the strong CP problem may be much lighter than the canonical QCD axion, down to the range of ultra-light axions, provided Nature has a $Z_{\mathcal{N}}$ symmetry implemented via \mathcal{N} degenerate world copies, one of which is our SM. The axion field realizes the symmetry non-linearly, which leads to exponential cancellations among the contributions from each mirror copy to the total axion potential. For large \mathcal{N} , we have shown that the total axion potential is given by a single cosine and we determined analytically the –exponentially suppressed– dependence of the axion mass on the number of mirror worlds, using the properties of hypergeometric functions and the Fourier expansion. In practice, the formula in Eq. (2.32) gives an excellent approximation even down to $\mathcal{N} = 3$. We have also improved the holomorphicity bounds previously obtained.

We compared next the predictions of the theory with present and future data from experiments which do not rely on the additional assumption that an axion abundance may explain the DM content of the Universe. It is particularly enticing that experiments set *a priori* to hunt only for ALPs may in fact be targeting solutions to the strong CP problem. For instance, ALPS II is shown to be able to probe the $Z_{\mathcal{N}}$ scenario here discussed down to $\mathcal{N} \sim 25$ for a large enough axion-photon coupling, while IAXO and BabyIAXO may test the whole \mathcal{N} landscape for values of that coupling even smaller, see Fig. 6. In turn, Fermi SN data can only reach $\mathcal{N} \gtrsim 43$ but are sensitive to smaller values of the coupling.

Highly dense stellar bodies allow one to set even stronger bounds in wide regions of the parameter space. These exciting limits have an added value: they avoid model-dependent assumptions about the axion couplings to SM particles, because they rely exclusively on the anomalous axion-gluon interaction needed to solve the strong CP problem. A dense medium of ordinary matter is a background that breaks the $Z_{\mathcal{N}}$ symmetry. This hampers the symmetry-induced cancellations in the total axion potential: the axion becomes heavier inside dense media *and* the minimum of the potential is located at $\theta_a = \pi$. From present solar data we obtain the bound $\mathcal{N} \lesssim 47$ provided $f_a \lesssim 2.4 \times 10^{15}$ GeV, while larger \mathcal{N} values are allowed for smaller f_a . Furthermore, we showed that projected neutron star and pulsar data should allow to test the scenario down to $\mathcal{N} \sim 9$ –and possibly even below– for the whole range of f_a , see Fig. 8. Furthermore, gravitational wave data from NS-NS and BH-NS mergers by LIGO/VIRGO and Advanced LIGO will allow to probe all values of \mathcal{N} for the remaining f_a range, up to the Planck scale and including the ultra-light axion mass range.

These analytical and phenomenological results have been derived within the model-independent framework of effective couplings. Nevertheless, for the sake of illustration,

we have developed two examples of UV completed models. One is a $Z_{\mathcal{N}}$ KSVZ model, which is shown to enjoy an improved PQ quality behaviour: its $Z_{\mathcal{N}}$ and PQ symmetries are linked and in consequence gauging $Z_{\mathcal{N}}$ alleviates much the PQ quality problem. The other UV completion considered in this paper is a $Z_{\mathcal{N}}$ version of the composite axion *à la* Kim-Choi. While this model is viable, its PQ quality is not improved with respect to the usual situation, because its $Z_{\mathcal{N}}$ and PQ symmetries are independent.

This work is intended to be a proof-of-concept that a much-lighter-than usual axion is a viable solution to the strong CP problem, with spectacular prospects of being tested in near future data. It also pinpoints that experiments searching for generic ALPs have in fact discovery potential to solve the strong CP problem.

The down-tuned axion considered here could also explain the DM content of the Universe in certain regions of the parameter space. The impact of such a light axion on the cosmological history is significant and it will be discussed in a separate paper [42].

Acknowledgments

We thank Gonzalo Alonso-Álvarez, Victor Engueta, Mary K. Gaillard, Yann Gouttenoire, Benjamin Grinstein, Lam Hui, David B. Kaplan, D. Marsh, V. Mehta, Philip Sørensen and Neal Weiner for illuminating discussions. M.B.G. and P.Q. are indebted for hospitality to the Theory Department of Columbia University in New York, where the initial stage of their work took place. The work of L.D.L. is supported by the Marie Skłodowska-Curie Individual Fellowship grant AXIONRUSH (GA 840791). L.D.L., P.Q. and A.R. acknowledge support by the Deutsche Forschungsgemeinschaft under Germany’s Excellence Strategy - EXC 2121 Quantum Universe - 390833306. M.B.G. acknowledges support from the “Spanish Agencia Estatal de Investigación” (AEI) and the EU “Fondo Europeo de Desarrollo Regional” (FEDER) through the projects FPA2016-78645-P and PID2019-108892RB-I00/AEI/10.13039/501100011033. M.B.G. and P. Q. acknowledge support from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreements 690575 (RISE InvisiblesPlus) and 674896 (ITN ELUSIVES), as well as from the Spanish Research Agency (Agencia Estatal de Investigación) through the grant IFT Centro de Excelencia Severo Ochoa SEV-2016-0597. This project has received funding/support from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860881-HIDDeN.

A Holomorphicity properties of $Z_{\mathcal{N}}$ axion potential

In order to determine the parameter b in Eq. (2.20), which controls the exponential suppression of the axion mass, it is necessary to study the region in the complex plane where the extension of the potential $V(w)$ is holomorphic. As the plots in Fig. 9 illustrate, both the potential and its second derivative have branch cuts starting in $w_{cut} = \pi \pm i \log z$. However, the second derivative $V''(w)$ diverges at the branch point and thus b cannot be extended all the way to $\log z$. In order to optimize the bound on the axion mass we allow b to depart from $\log z$, $b = |\log z + \Delta b|$. Taking into account that $V''(w)$ for small Δb can be approximated by

$$V''(\pi + i(\log z + \Delta b)) \simeq -m_{\pi}^2 f_{\pi}^2 \sqrt{\frac{1-z}{1+z}} \left[\frac{1}{4} \frac{1}{(\Delta b)^{3/2}} + \mathcal{O}(\Delta b^{-1/2}) \right], \quad (\text{A.1})$$

the procedure amounts to minimize the function $B(\Delta b)$ that determines the bound $|E_{\mathcal{N}}(V)| \leq B(\Delta b)$ (see Eq. (2.20),

$$B(\Delta b) \equiv \frac{4\pi M(\Delta b)}{e^{\mathcal{N}|\log z + \Delta b|} - 1} = \pi m_{\pi}^2 f_{\pi}^2 \sqrt{\frac{1-z}{1+z}} \frac{1}{(\Delta b)^{3/2}} \frac{1}{e^{\mathcal{N}|\log z - \Delta b|} - 1}. \quad (\text{A.2})$$

The requirement $\frac{dB(\Delta b)}{d(\Delta b)} = 0$ shows that the bound is optimized for

$$\Delta b = \frac{3}{2} \frac{1}{\mathcal{N}}, \quad (\text{A.3})$$

where the factor $3/2$ comes from the order of the divergence in Eq. (A.1).




Figure 9: Representation of the complex functions $V(w)$ (left) and $V''(w)$ (right). Colors represent the phase of the corresponding complex function and the brightness represents the modulus. The singularities can be clearly spotted: branch cuts starting from $w_{cut} = \pi \pm i \log z$ in both functions and divergences in those same points for $V''(w)$.

B Fourier series of the $Z_{\mathcal{N}}$ axion potential

We show here that the coefficients of the Fourier series of any $Z_{\mathcal{N}}$ symmetric potential, such as the $Z_{\mathcal{N}}$ axion potential in Eq. (2.8), vanish unless the corresponding Fourier mode is a multiple of \mathcal{N} . Moreover it will be shown that, when the potential is expressed as

$$V_{\mathcal{N}}(\theta_a) = \sum_{k=0}^{\mathcal{N}-1} V\left(\theta_a + \frac{2\pi k}{\mathcal{N}}\right), \quad (\text{B.1})$$

the non-vanishing coefficients of the Fourier series can be expressed in terms of the Fourier transformation of a single term in the sum Eq. (B.1).

Let us denote by $\hat{V}_{\mathcal{N}}(n)$ the coefficients of the Fourier series of the total potential,

$$V_{\mathcal{N}}(\theta_a) \equiv \sum_{n=-\infty}^{\infty} e^{in\theta_a} \hat{V}_{\mathcal{N}}(n), \quad (\text{B.2})$$

and by $\hat{V}_{2\pi k/\mathcal{N}}(n)$ the coefficients of the Fourier series of each of the terms in the sum in Eq. (B.1),

$$V\left(\theta_a + \frac{2\pi k}{\mathcal{N}}\right) \equiv \sum_{n=-\infty}^{\infty} \hat{V}_{2\pi k/\mathcal{N}}(n) e^{in\theta_a}. \quad (\text{B.3})$$

We will stick to the notation that omits the subindex for the first world ($k = 0$), $\hat{V}_0(n) \equiv \hat{V}(n)$,

$$V(\theta_a) = \sum_{n=-\infty}^{\infty} \hat{V}(n) e^{in\theta_a} \quad \text{with} \quad \hat{V}(n) = \frac{1}{2\pi} \int_0^{2\pi} V(x) e^{-inx} dx. \quad (\text{B.4})$$

Each term in the sum in Eq. (B.1) and Eq. (B.3) corresponds to the function in the first term but with its argument shifted by $\frac{2\pi k}{\mathcal{N}}$. The shift property of the Fourier series relates the Fourier coefficients of the shifted functions $\hat{V}_{2\pi k/\mathcal{N}}(n)$ to that of the original function,

$$\hat{V}_{2\pi k/\mathcal{N}}(n) = e^{in\frac{2\pi k}{\mathcal{N}}} \hat{V}(n). \quad (\text{B.5})$$

Substituting this expression in Eq. (B.3), and inserting the latter in Eq. (B.1), it follows that the total potential can be written as

$$V_{\mathcal{N}}(\theta_a) = \sum_{k=0}^{\mathcal{N}-1} \sum_{n=-\infty}^{\infty} \hat{V}_{2\pi k/\mathcal{N}}(n) e^{in\theta_a} = \sum_{n=-\infty}^{\infty} \hat{V}(n) e^{in\theta_a} \sum_{k=0}^{\mathcal{N}-1} e^{in\frac{2\pi k}{\mathcal{N}}}. \quad (\text{B.6})$$

Comparing this expression with Eq. (B.2), it follows that the coefficients of the Fourier series for the total potential are given by

$$\hat{V}_{\mathcal{N}}(n) = \hat{V}(n) \sum_{k=0}^{\mathcal{N}-1} e^{in\frac{2\pi k}{\mathcal{N}}}. \quad (\text{B.7})$$

Interestingly, these coefficients vanish unless n is a multiple of \mathcal{N}

$$\text{If } n \pmod{\mathcal{N}} \neq 0 \implies \sum_{k=0}^{\mathcal{N}-1} e^{in\frac{2\pi k}{\mathcal{N}}} = 0 \implies \hat{V}_{\mathcal{N}}(n) = 0, \quad (\text{B.8})$$

$$\text{If } n \pmod{\mathcal{N}} = 0 \implies \sum_{k=0}^{\mathcal{N}-1} e^{in\frac{2\pi k}{\mathcal{N}}} = \mathcal{N} \implies \hat{V}_{\mathcal{N}}(n) = \mathcal{N} \hat{V}(n). \quad (\text{B.9})$$

To sum up, the Fourier series of the total potential $V_{\mathcal{N}}(\theta)$ can be easily obtained in terms of the Fourier series of a single term $V(\theta)$ and it only receives contributions from the modes that are multiples of \mathcal{N} . In our case of interest the potential is real and even, this translates into

$$V_{\mathcal{N}}(\theta_a) = 2\mathcal{N} \sum_{t=1}^{\infty} \hat{V}(t\mathcal{N}) \cos(t\mathcal{N}\theta_a), \quad (\text{B.10})$$

where the factor of two comes from the negative modes and the constant term (i.e. θ_a -independent) has been obviated.

C Analytical axion mass dependence from hypergeometric functions

We show here that the Fourier series coefficients of the single world axion potential in Eq. (2.27),

$$\hat{V}(n) = -\frac{m_{\pi}^2 f_{\pi}^2}{1+z} \int_0^{2\pi} \cos(nt) \sqrt{1+z^2+2z\cos(t)} dt, \quad (\text{C.1})$$

can be written for large Fourier modes, $n \gg 1$, as a simple analytical formula that exponentially decays with n . Moreover, by applying the result in Appendix B, it will be shown that the potential for the $Z_{\mathcal{N}}$ axion approaches a single cosine and a simple formula for the $Z_{\mathcal{N}}$ axion mass follows.

Let us start by relating the Fourier series decomposition of the single world potential in Eq. (C.1) with the Gauss hypergeometric functions (see for example Eq. (9.112) in Ref. [55]),

$${}_2F_1 \left(\begin{matrix} p, n+p \\ n+1 \end{matrix} \middle| w^2 \right) = \frac{w^{-n}}{2\pi} \frac{\Gamma(p)n!}{\Gamma(p+n)} \int_0^{2\pi} \frac{\cos(nt) dt}{(1-2w\cos t+w^2)^p}. \quad (\text{C.2})$$

Via the identification $w = -z$ and $p = -1/2$, $\hat{V}(n)$ can be written as

$$\hat{V}(n) = (-1)^{n+1} \frac{m_\pi^2 f_\pi^2}{1+z} z^n \frac{\Gamma(n-1/2)}{\Gamma(-1/2) n!} {}_2F_1 \left(\begin{matrix} -1/2, n-1/2 \\ n+1 \end{matrix} \middle| z^2 \right). \quad (\text{C.3})$$

For convenience, the hypergeometric function can be also expressed as (see Eq. (9.131) from Ref. [55])

$${}_2F_1 \left(\begin{matrix} \alpha, \beta \\ \gamma \end{matrix} \middle| w \right) = (1-w)^{-\alpha} {}_2F_1 \left(\begin{matrix} \alpha, \gamma-\beta \\ \gamma \end{matrix} \middle| \frac{w}{w-1} \right), \quad (\text{C.4})$$

so that

$${}_2F_1 \left(\begin{matrix} -1/2, n-1/2 \\ n+1 \end{matrix} \middle| z^2 \right) = (1-z^2)^{1/2} {}_2F_1 \left(\begin{matrix} -1/2, 3/2 \\ n+1 \end{matrix} \middle| \frac{z^2}{z^2-1} \right). \quad (\text{C.5})$$

The relation in Eq. (C.3) is exact. However, only the modes n which are multiples of \mathcal{N} contribute to the potential (see Appendix B), and therefore it is pertinent to focus on the large n limit. While the limit of the Gauss hypergeometric function when one or more of its parameters become large is difficult to compute in general, some asymptotic expansions of the hypergeometric function are known in the literature. In particular, following Ref. [121, 122],

$$\lim_{\gamma \rightarrow \infty} {}_2F_1 \left(\begin{matrix} \alpha, \beta \\ \gamma \end{matrix} \middle| w \right) = 1 + \frac{\alpha\beta}{\gamma} w + \mathcal{O}\left(\frac{w}{\gamma}\right)^2. \quad (\text{C.6})$$

In turn, the prefactor in Eq. (C.3) simplifies in the large n limit to

$$\lim_{n \rightarrow \infty} \frac{\Gamma(n-1/2)}{\Gamma(-1/2) n!} = -\frac{1}{2\sqrt{\pi}} n^{-3/2}. \quad (\text{C.7})$$

Putting all this together, it follows that, in the large n limit, the coefficient of the Fourier series for a single world is given by

$$\hat{V}(n) = (-1)^n \frac{m_\pi^2 f_\pi^2}{2\sqrt{\pi}} \sqrt{\frac{1-z}{1+z}} n^{-3/2} z^n, \quad (\text{C.8})$$

which in turn implies in this limit that the total $Z_{\mathcal{N}}$ potential in Eq. (B.10) can be written as

$$V_{\mathcal{N}}(\theta_a) = \mathcal{N} \sum_{t=1}^{\infty} (-1)^{t\mathcal{N}} \frac{m_\pi^2 f_\pi^2}{2\sqrt{\pi}} \sqrt{\frac{1-z}{1+z}} (t\mathcal{N})^{-3/2} z^{t\mathcal{N}} \cos(t\mathcal{N}\theta_a). \quad (\text{C.9})$$

This expression allows us to understand several properties of the total potential. Firstly, it can be shown now that the total potential approaches a single cosine in the large \mathcal{N} , since all the other modes are then exponentially suppressed with respect to the first one,

$$\lim_{\mathcal{N} \rightarrow \infty} \left| \frac{\hat{V}_{\mathcal{N}}(t\mathcal{N})}{\hat{V}_{\mathcal{N}}(\mathcal{N})} \right| = \lim_{\mathcal{N} \rightarrow \infty} \left| \frac{\hat{V}(t\mathcal{N})}{\hat{V}(\mathcal{N})} \right| = t^{-3/2} z^{(t-1)\mathcal{N}} \rightarrow 0, \quad (\text{C.10})$$

and thus the potential reads

$$V_{\mathcal{N}}(\theta_a) \xrightarrow{\mathcal{N} \gg 1} \frac{m_\pi^2 f_\pi^2}{\sqrt{\pi}} \sqrt{\frac{1-z}{1+z}} \mathcal{N}^{-1/2} \cdot (-1)^{\mathcal{N}} z^{\mathcal{N}} \cos(\mathcal{N}\theta_a), \quad (\text{C.11})$$

Secondly, we can also obtain an analytical expression for the axion mass that confirms the dependence obtained from the holomorphicity arguments in Section 2.3.1, and completes the expresion with the correct prefactor,

$$m_a^2 f_a^2 \simeq \frac{m_\pi^2 f_\pi^2}{\sqrt{\pi}} \sqrt{\frac{1-z}{1+z}} \mathcal{N}^{3/2} z^{\mathcal{N}}. \quad (\text{C.12})$$

Finally, it is now trivial to show that the potential in the large \mathcal{N} limit has \mathcal{N} minima (maxima) located at $\theta_a = \{\pm 2\pi\ell/\mathcal{N}\}$ for $\ell = 0, 1, \dots, \frac{\mathcal{N}-1}{2}$, for odd (even) \mathcal{N} .

The results above assumed the large \mathcal{N} limit. However, note that the conclusion about the location of the extrema is true for any \mathcal{N} . This can be easily seen after obtaining the *exact* Fourier expansion of the $Z_{\mathcal{N}}$ axion potential in Eq. (2.8), which reads,

$$V_{\mathcal{N}}(\theta_a) = -m_{\pi}^2 f_{\pi}^2 \mathcal{N} \sum_{t=1}^{\infty} (-1)^{t\mathcal{N}+1} \sum_{\ell=t\mathcal{N}}^{\infty} \frac{(2\ell)!(2\ell)!}{2^{4\ell-1} (2\ell-1)(\ell!)^2 (\ell-t\mathcal{N})!(\ell+t\mathcal{N})!} \beta^{\ell} \cos(t\mathcal{N}\theta_a). \quad (\text{C.13})$$

For even \mathcal{N} , it trivially follows that $\theta_a = 0$ is a maximum, as all factors in this expression are positive except for the factor $(1-2\ell) < 0$. For odd \mathcal{N} instead the $(-1)^{t\mathcal{N}}$ factor fluctuates in sign, but the first term ($t=1$) is positive and dominates the expansion (e.g. it is exponentially larger in magnitude than the $t=2$ term which is negative). The periodicity of the potential extends these conclusions to the location of all extrema.

References

- [1] R. D. Peccei and H. R. Quinn, “CP Conservation in the Presence of Instantons,” *Phys. Rev. Lett.* **38** (1977) 1440–1443. [[328\(1977\)](#)]. [3](#)
- [2] R. D. Peccei and H. R. Quinn, “Constraints Imposed by CP Conservation in the Presence of Instantons,” *Phys. Rev.* **D16** (1977) 1791–1797. [3](#)
- [3] S. Weinberg, “A New Light Boson?,” *Phys. Rev. Lett.* **40** (1978) 223–226. [3](#)
- [4] F. Wilczek, “Problem of Strong P and T Invariance in the Presence of Instantons,” *Phys. Rev. Lett.* **40** (1978) 279–282. [3](#)
- [5] A. R. Zhitnitsky, “On Possible Suppression of the Axion Hadron Interactions. (In Russian),” *Sov. J. Nucl. Phys.* **31** (1980) 260. [[Yad. Fiz.31,497\(1980\)](#)]. [3](#)
- [6] M. Dine, W. Fischler, and M. Srednicki, “A Simple Solution to the Strong CP Problem with a Harmless Axion,” *Phys. Lett.* **B104** (1981) 199–202. [3](#)
- [7] J. E. Kim, “Weak Interaction Singlet and Strong CP Invariance,” *Phys. Rev. Lett.* **43** (1979) 103. [3](#), [5](#)
- [8] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, “Can Confinement Ensure Natural CP Invariance of Strong Interactions?,” *Nucl. Phys.* **B166** (1980) 493–506. [3](#)
- [9] L. Di Luzio, F. Mescia, and E. Nardi, “Redefining the Axion Window,” *Phys. Rev. Lett.* **118** no. 3, (2017) 031801, [arXiv:1610.07593 \[hep-ph\]](#). [3](#), [14](#)
- [10] M. Farina, D. Pappadopulo, F. Rompineve, and A. Tesi, “The photo-philic QCD axion,” *JHEP* **01** (2017) 095, [arXiv:1611.09855 \[hep-ph\]](#).
- [11] L. Di Luzio, F. Mescia, and E. Nardi, “Window for preferred axion models,” *Phys. Rev.* **D96** no. 7, (2017) 075003, [1705.05370](#). [14](#)
- [12] P. Agrawal, J. Fan, M. Reece, and L.-T. Wang, “Experimental Targets for Photon Couplings of the QCD Axion,” *JHEP* **02** (2018) 006, [arXiv:1709.06085 \[hep-ph\]](#).
- [13] G. Marques-Tavares and M. Teo, “Light axions with large hadronic couplings,” *JHEP* **05** (2018) 180, [arXiv:1803.07575 \[hep-ph\]](#).

- [14] L. Di Luzio, M. Giannotti, E. Nardi, and L. Visinelli, “The landscape of QCD axion models,” *Phys. Rept.* **870** (2020) 1–117, [arXiv:2003.01100 \[hep-ph\]](#).
- [15] L. Darmé, L. Di Luzio, M. Giannotti, and E. Nardi, “Selective enhancement of the QCD axion couplings,” [arXiv:2010.15846 \[hep-ph\]](#). 3
- [16] D. Budker, P. W. Graham, M. Ledbetter, S. Rajendran, and A. Sushkov, “Proposal for a Cosmic Axion Spin Precession Experiment (CASPER),” *Phys. Rev.* **X4** no. 2, (2014) 021030, [arXiv:1306.6089 \[hep-ph\]](#). 4
- [17] D. F. Jackson Kimball *et al.*, “Overview of the Cosmic Axion Spin Precession Experiment (CASPER),” [arXiv:1711.08999 \[physics.ins-det\]](#). 4
- [18] V. A. Rubakov, “Grand unification and heavy axion,” *JETP Lett.* **65** (1997) 621–624, [arXiv:hep-ph/9703409 \[hep-ph\]](#). 4
- [19] Z. Berezhiani, L. Gianfagna, and M. Giannotti, “Strong CP problem and mirror world: The Weinberg-Wilczek axion revisited,” *Phys. Lett.* **B500** (2001) 286–296, [arXiv:hep-ph/0009290 \[hep-ph\]](#).
- [20] L. Gianfagna, M. Giannotti, and F. Nesti, “Mirror world, supersymmetric axion and gamma ray bursts,” *JHEP* **10** (2004) 044, [arXiv:hep-ph/0409185](#).
- [21] S. D. H. Hsu and F. Sannino, “New solutions to the strong CP problem,” *Phys. Lett.* **B605** (2005) 369–375, [arXiv:hep-ph/0408319 \[hep-ph\]](#).
- [22] A. Hook, “Anomalous solutions to the strong CP problem,” *Phys. Rev. Lett.* **114** no. 14, (2015) 141801, [arXiv:1411.3325 \[hep-ph\]](#).
- [23] H. Fukuda, K. Harigaya, M. Ibe, and T. T. Yanagida, “Model of visible QCD axion,” *Phys. Rev.* **D92** no. 1, (2015) 015021, [arXiv:1504.06084 \[hep-ph\]](#).
- [24] C.-W. Chiang, H. Fukuda, M. Ibe, and T. T. Yanagida, “750 GeV diphoton resonance in a visible heavy QCD axion model,” *Phys. Rev.* **D93** no. 9, (2016) 095016, [arXiv:1602.07909 \[hep-ph\]](#).
- [25] S. Dimopoulos, A. Hook, J. Huang, and G. Marques-Tavares, “A Collider Observable QCD Axion,” *JHEP* **11** (2016) 052, [arXiv:1606.03097 \[hep-ph\]](#).
- [26] T. Gherghetta, N. Nagata, and M. Shifman, “A Visible QCD Axion from an Enlarged Color Group,” *Phys. Rev.* **D93** no. 11, (2016) 115010, [arXiv:1604.01127 \[hep-ph\]](#).
- [27] A. Kobakhidze, “Heavy axion in asymptotically safe QCD,” [arXiv:1607.06552 \[hep-ph\]](#).
- [28] P. Agrawal and K. Howe, “Factoring the Strong CP Problem,” *JHEP* **12** (2018) 029, [arXiv:1710.04213 \[hep-ph\]](#).
- [29] P. Agrawal and K. Howe, “A Flavorful Factoring of the Strong CP Problem,” *JHEP* **12** (2018) 035, [arXiv:1712.05803 \[hep-ph\]](#).
- [30] M. K. Gaillard, M. B. Gavela, R. Houtz, P. Quilez, and R. Del Rey, “Color unified dynamical axion,” *Eur. Phys. J.* **C78** no. 11, (2018) 972, [arXiv:1805.06465 \[hep-ph\]](#).
- [31] M. A. Buen-Abad and J. Fan, “Dynamical axion misalignment with small instantons,” *JHEP* **12** (2019) 161, [arXiv:1911.05737 \[hep-ph\]](#).

- [32] A. Hook, S. Kumar, Z. Liu, and R. Sundrum, “High Quality QCD Axion and the LHC,” *Phys. Rev. Lett.* **124** no. 22, (2020) 221801, [arXiv:1911.12364 \[hep-ph\]](#).
- [33] C. Csáki, M. Ruhdorfer, and Y. Shirman, “UV Sensitivity of the Axion Mass from Instantons in Partially Broken Gauge Groups,” *JHEP* **04** (2020) 031, [arXiv:1912.02197 \[hep-ph\]](#).
- [34] T. Gherghetta and M. D. Nguyen, “A Composite Higgs with a Heavy Composite Axion,” [arXiv:2007.10875 \[hep-ph\]](#). 4
- [35] A. Hook, “Solving the Hierarchy Problem Discretely,” *Phys. Rev. Lett.* **120** no. 26, (2018) 261802, [arXiv:1802.10093 \[hep-ph\]](#). 4, 5, 7, 8, 10, 11, 12
- [36] M. Giannotti, “Mirror world and axion: Relaxing cosmological bounds,” *Int. J. Mod. Phys. A* **20** (2005) 2454–2458, [arXiv:astro-ph/0504636](#). 4, 18
- [37] A. Hook and J. Huang, “Probing axions with neutron star inspirals and other stellar processes,” *JHEP* **06** (2018) 036, [arXiv:1708.08464 \[hep-ph\]](#). 5, 19, 20, 21, 22, 23
- [38] J. Huang, M. C. Johnson, L. Sagunski, M. Sakellariadou, and J. Zhang, “Prospects for axion searches with Advanced LIGO through binary mergers,” *Phys. Rev. D* **99** no. 6, (2019) 063013, [arXiv:1807.02133 \[hep-ph\]](#). 5, 19, 22, 23
- [39] M. A. Shifman, *Instantons in Gauge Theories*. 1994. 5
- [40] J. E. Kim, “A Composite Invisible Axion,” *Phys. Rev.* **D31** (1985) 1733. 5, 16
- [41] K. Choi and J. E. Kim, “Dynamical Axion,” *Phys. Rev.* **D32** (1985) 1828. 5, 16
- [42] L. Di Luzio, B. Gavela, P. Quilez, and A. Ringwald, “Dark Matter from an even lighter QCD axion: trapped misalignment (To appear),” [arXiv:2102.01082 \[hep-ph\]](#). 5, 8, 9, 15, 18, 19, 22, 24
- [43] P. Di Vecchia and G. Veneziano, “Chiral Dynamics in the Large n Limit,” *Nucl. Phys.* **B171** (1980) 253–272. 6
- [44] H. Leutwyler and A. V. Smilga, “Spectrum of Dirac operator and role of winding number in QCD,” *Phys. Rev. D* **46** (1992) 5607–5632. 7
- [45] G. Grilli di Cortona, E. Hardy, J. Pardo Vega, and G. Villadoro, “The QCD axion, precisely,” *JHEP* **01** (2016) 034, [arXiv:1511.02867 \[hep-ph\]](#). 6, 7, 8, 18
- [46] S. Borsanyi *et al.*, “Calculation of the axion mass based on high-temperature lattice quantum chromodynamics,” *Nature* **539** no. 7627, (2016) 69–71, [arXiv:1606.07494 \[hep-lat\]](#). 7
- [47] L. Ubaldi, “Effects of theta on the deuteron binding energy and the triple-alpha process,” *Phys. Rev.* **D81** (2010) 025011, [arXiv:0811.1599 \[hep-ph\]](#). 9, 21
- [48] D. Lee, U.-G. Meißner, K. A. Olive, M. Shifman, and T. Vonk, “ θ -dependence of light nuclei and nucleosynthesis,” *Phys. Rev. Res.* **2** no. 3, (2020) 033392, [arXiv:2006.12321 \[hep-ph\]](#). 9, 21
- [49] Z. Berezhiani, D. Comelli, and F. L. Villante, “The Early mirror universe: Inflation, baryogenesis, nucleosynthesis and dark matter,” *Phys. Lett.* **B503** (2001) 362–375, [arXiv:hep-ph/0008105 \[hep-ph\]](#). 9

- [50] Z. Berezhiani, “Mirror world and its cosmological consequences,” *Int. J. Mod. Phys. A* **19** (2004) 3775–3806, [arXiv:hep-ph/0312335](#).
- [51] R. Foot, “Mirror dark matter: Cosmology, galaxy structure and direct detection,” *Int. J. Mod. Phys. A* **29** (2014) 1430013, [arXiv:1401.3965 \[astro-ph.CO\]](#). 9
- [52] R. Foot, A. Kobakhidze, K. L. McDonald, and R. R. Volkas, “Poincare protection for a natural electroweak scale,” *Phys. Rev. D* **89** no. 11, (2014) 115018, [arXiv:1310.0223 \[hep-ph\]](#). 10
- [53] R. R. Volkas, A. J. Davies, and G. C. Joshi, “Naturalness of the invisible axion model,” *Phys. Lett.* **B215** (1988) 133–138. 10
- [54] L. N. Trefethen and J. A. C. Weideman, “The exponentially convergent trapezoidal rule,” *SIAM Review* **56** no. 3, (2014) 385–458, <https://doi.org/10.1137/130932132>. <https://doi.org/10.1137/130932132>. 10
- [55] I. GRADSHTEYN and I. RYZHIK, “8-9 - special functions,” in *Table of Integrals, Series, and Products*, I. GRADSHTEYN and I. RYZHIK, eds., pp. 904 – 1080. Academic Press, 1980. <http://www.sciencedirect.com/science/article/pii/B9780122947605500209>. 12, 26, 27
- [56] R. Holman, S. D. H. Hsu, T. W. Kephart, E. W. Kolb, R. Watkins, and L. M. Widrow, “Solutions to the Strong CP Problem in a World with Gravity,” *Phys. Lett.* **B282** (1992) 132–136, [arXiv:hep-ph/9203206 \[hep-ph\]](#). 14
- [57] M. Kamionkowski and J. March-Russell, “Planck scale physics and the Peccei-Quinn mechanism,” *Phys. Lett. B* **282** (1992) 137–141, [arXiv:hep-th/9202003](#).
- [58] S. M. Barr and D. Seckel, “Planck scale corrections to axion models,” *Phys. Rev. D* **46** (1992) 539–549.
- [59] S. Ghigna, M. Lusignoli, and M. Roncadelli, “Instability of the Invisible Axion,” *Phys. Lett.* **B283** (1992) 278–281. 14
- [60] H. M. Georgi, L. J. Hall, and M. B. Wise, “Grand Unified Models with an Automatic Peccei-Quinn Symmetry,” *Nucl. Phys.* **B192** (1981) 409–416.
- [61] S. B. Giddings and A. Strominger, “Loss of Incoherence and Determination of Coupling Constants in Quantum Gravity,” *Nucl. Phys.* **B307** (1988) 854–866.
- [62] S. R. Coleman, “Why There is Nothing Rather Than Something: a Theory of the Cosmological Constant,” *Nucl. Phys.* **B310** (1988) 643–668.
- [63] G. Gilbert, “Wormhole Induced Proton Decay,” *Nucl. Phys.* **B328** (1989) 159–170.
- [64] S.-J. Rey, “The Axion Dynamics in Wormhole Background,” *Phys. Rev.* **D39** (1989) 3185.
- [65] J. Alvey and M. Escudero, “The axion quality problem: global symmetry breaking and wormholes,” *JHEP* **01** (2021) 032, [arXiv:2009.03917 \[hep-ph\]](#). 14
- [66] L. Randall, “Composite axion models and Planck scale physics,” *Phys. Lett.* **B284** (1992) 77–80. 14, 17

- [67] B. A. Dobrescu, “The Strong CP Problem Versus Planck Scale Physics,” *Phys. Rev.* **D55** (1997) 5826–5833, [arXiv:hep-ph/9609221 \[hep-ph\]](#).
- [68] D. Butter and M. K. Gaillard, “The Axion mass in modular invariant supergravity,” *Phys. Lett.* **B612** (2005) 304–310, [arXiv:hep-th/0502100 \[hep-th\]](#).
- [69] M. Redi and R. Sato, “Composite Accidental Axions,” *JHEP* **05** (2016) 104, [arXiv:1602.05427 \[hep-ph\]](#). 17
- [70] H. Fukuda, M. Ibe, M. Suzuki, and T. T. Yanagida, “A ”gauged” $U(1)$ Peccei–Quinn symmetry,” *Phys. Lett.* **B771** (2017) 327–331, [arXiv:1703.01112 \[hep-ph\]](#).
- [71] L. Di Luzio, E. Nardi, and L. Ubaldi, “Accidental Peccei-Quinn symmetry protected to arbitrary order,” *Phys. Rev. Lett.* **119** no. 1, (2017) 011801, [arXiv:1704.01122 \[hep-ph\]](#).
- [72] H. Fukuda, M. Ibe, M. Suzuki, and T. T. Yanagida, “Gauged Peccei-Quinn symmetry – A case of simultaneous breaking of SUSY and PQ symmetry,” *JHEP* **07** (2018) 128, [arXiv:1803.00759 \[hep-ph\]](#).
- [73] M. Ibe, M. Suzuki, and T. T. Yanagida, “ $B - L$ as a Gauged Peccei-Quinn Symmetry,” *JHEP* **08** (2018) 049, [arXiv:1805.10029 \[hep-ph\]](#). 17
- [74] B. Lillard and T. M. P. Tait, “A High Quality Composite Axion,” [arXiv:1811.03089 \[hep-ph\]](#).
- [75] M. Ardu, L. Di Luzio, G. Landini, A. Strumia, D. Teresi, and J.-W. Wang, “Axion quality from the (anti)symmetric of $SU(\mathcal{N})$,” *JHEP* **11** (2020) 090, [arXiv:2007.12663 \[hep-ph\]](#).
- [76] L. Di Luzio, “Accidental $SO(10)$ axion from gauged flavour,” *JHEP* **11** (2020) 074, [arXiv:2008.09119 \[hep-ph\]](#).
- [77] Q. Bonnefoy, P. Cox, E. Dudas, T. Gherghetta, and M. D. Nguyen, “Flavoured Warped Axion,” [arXiv:2012.09728 \[hep-ph\]](#).
- [78] Q. Bonnefoy, E. Dudas, and S. Pokorski, “Axions in a highly protected gauge symmetry model,” *Eur. Phys. J. C* **79** no. 1, (2019) 31, [arXiv:1804.01112 \[hep-ph\]](#). 14
- [79] R. Alonso and A. Urbano, “Wormholes and masses for Goldstone bosons,” [arXiv:1706.07415 \[hep-ph\]](#). 14
- [80] M. B. Gavela, M. Ibe, P. Quilez, and T. T. Yanagida, “Automatic Peccei–Quinn symmetry,” *Eur. Phys. J. C* **79** no. 6, (2019) 542, [arXiv:1812.08174 \[hep-ph\]](#). 17
- [81] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper, and J. March-Russell, “String Axiverse,” *Phys. Rev. D* **81** (2010) 123530, [arXiv:0905.4720 \[hep-th\]](#). 17
- [82] D. Brzeminski, Z. Chacko, A. Dev, and A. Hook, “A Time-Varying Fine Structure Constant from Naturally Ultralight Dark Matter,” [arXiv:2012.02787 \[hep-ph\]](#). 17
- [83] K. Ehret *et al.*, “New ALPS Results on Hidden-Sector Lightweights,” *Phys. Lett. B* **689** (2010) 149–155, [arXiv:1004.1313 \[hep-ex\]](#). 19
- [84] **CAST** Collaboration, S. Andriamonje *et al.*, “An Improved limit on the axion-photon coupling from the CAST experiment,” *JCAP* **04** (2007) 010, [arXiv:hep-ex/0702006](#).

- [85] **CAST** Collaboration, V. Anastassopoulos *et al.*, “New CAST Limit on the Axion-Photon Interaction,” *Nature Phys.* **13** (2017) 584–590, [arXiv:1705.02290 \[hep-ex\]](#).
- [86] M. Betz, F. Caspers, M. Gasiot, M. Thumm, and S. Rieger, “First results of the CERN Resonant Weakly Interacting sub-eV Particle Search (CROWS),” *Phys. Rev. D* **88** no. 7, (2013) 075014, [arXiv:1310.8098 \[physics.ins-det\]](#).
- [87] I. Shilon, A. Dudarev, H. Silva, and H. ten Kate, “Conceptual Design of a New Large Superconducting Toroid for IAXO, the New International AXion Observatory,” *IEEE Trans. Appl. Supercond.* **23** no. 3, (2013) 4500604, [arXiv:1212.4633 \[physics.ins-det\]](#).
- [88] **OSQAR** Collaboration, R. Ballou *et al.*, “New exclusion limits on scalar and pseudoscalar axionlike particles from light shining through a wall,” *Phys. Rev. D* **92** no. 9, (2015) 092002, [arXiv:1506.08082 \[hep-ex\]](#).
- [89] F. Della Valle, A. Ejlli, U. Gastaldi, G. Messineo, E. Milotti, R. Pengo, G. Ruoso, and G. Zavattini, “The PVLAS experiment: measuring vacuum magnetic birefringence and dichroism with a birefringent Fabry–Perot cavity,” *Eur. Phys. J. C* **76** no. 1, (2016) 24, [arXiv:1510.08052 \[physics.optics\]](#).
- [90] R. Bähre *et al.*, “Any light particle search II —Technical Design Report,” *JINST* **8** (2013) T09001, [arXiv:1302.5647 \[physics.ins-det\]](#). 19
- [91] D. Wouters and P. Brun, “Constraints on Axion-like Particles from X-Ray Observations of the Hydra Galaxy Cluster,” *Astrophys. J.* **772** (2013) 44, [arXiv:1304.0989 \[astro-ph.HE\]](#). 19
- [92] M. D. Marsh, H. R. Russell, A. C. Fabian, B. P. McNamara, P. Nulsen, and C. S. Reynolds, “A New Bound on Axion-Like Particles,” *JCAP* **12** (2017) 036, [arXiv:1703.07354 \[hep-ph\]](#).
- [93] C. S. Reynolds, M. D. Marsh, H. R. Russell, A. C. Fabian, R. Smith, F. Tombesi, and S. Veilleux, “Astrophysical limits on very light axion-like particles from Chandra grating spectroscopy of NGC 1275,” [arXiv:1907.05475 \[hep-ph\]](#).
- [94] M. Meyer and T. Petrushevska, “Search for Axionlike-Particle-Induced Prompt γ -Ray Emission from Extragalactic Core-Collapse Supernovae with the *Fermi* Large Area Telescope,” *Phys. Rev. Lett.* **124** no. 23, (2020) 231101, [arXiv:2006.06722 \[astro-ph.HE\]](#). [Erratum: *Phys.Rev.Lett.* 125, 119901 (2020)].
- [95] M. Meyer, M. Giannotti, A. Mirizzi, J. Conrad, and M. Sánchez-Conde, “Fermi Large Area Telescope as a Galactic Supernovae Axionscope,” *Phys. Rev. Lett.* **118** no. 1, (2017) 011103, [arXiv:1609.02350 \[astro-ph.HE\]](#).
- [96] **Fermi-LAT** Collaboration, M. Ajello *et al.*, “Search for Spectral Irregularities due to Photon–Axionlike-Particle Oscillations with the Fermi Large Area Telescope,” *Phys. Rev. Lett.* **116** no. 16, (2016) 161101, [arXiv:1603.06978 \[astro-ph.HE\]](#).
- [97] **H.E.S.S.** Collaboration, A. Abramowski *et al.*, “Constraints on axionlike particles with H.E.S.S. from the irregularity of the PKS 2155-304 energy spectrum,” *Phys. Rev. D* **88** no. 10, (2013) 102003, [arXiv:1311.3148 \[astro-ph.HE\]](#).

- [98] A. Ayala, I. Domínguez, M. Giannotti, A. Mirizzi, and O. Straniero, “Revisiting the bound on axion-photon coupling from Globular Clusters,” *Phys. Rev. Lett.* **113** no. 19, (2014) 191302, [arXiv:1406.6053](#) [[astro-ph.SR](#)].
- [99] A. Payez, C. Evoli, T. Fischer, M. Giannotti, A. Mirizzi, and A. Ringwald, “Revisiting the SN1987A gamma-ray limit on ultralight axion-like particles,” *JCAP* **02** (2015) 006, [arXiv:1410.3747](#) [[astro-ph.HE](#)].
- [100] A. Abeln *et al.*, “Axion search with BabyIAXO in view of IAXO,” in *40th International Conference on High Energy Physics*. 12, 2020. [arXiv:2012.06634](#) [[physics.ins-det](#)].
- [101] **BabyIAXO** Collaboration, A. Abeln *et al.*, “Conceptual Design of BabyIAXO, the intermediate stage towards the International Axion Observatory,” [arXiv:2010.12076](#) [[physics.ins-det](#)]. 19
- [102] C. O’Hare, “cajohare/axionlimits: Axionlimits,” July, 2020. <https://doi.org/10.5281/zenodo.3932430>. 19
- [103] H. Georgi, D. B. Kaplan, and L. Randall, “Manifesting the Invisible Axion at Low-energies,” *Phys. Lett.* **169B** (1986) 73–78. 18
- [104] G. G. Raffelt, “Astrophysical axion bounds,” *Lect. Notes Phys.* **741** (2008) 51–71, [arXiv:hep-ph/0611350](#) [[hep-ph](#)]. [51(2006)]. 19
- [105] A. Arvanitaki, M. Baryakhtar, and X. Huang, “Discovering the QCD Axion with Black Holes and Gravitational Waves,” *Phys. Rev. D* **91** no. 8, (2015) 084011, [arXiv:1411.2263](#) [[hep-ph](#)]. 19
- [106] A. Arvanitaki and S. Dubovsky, “Exploring the String Axiverse with Precision Black Hole Physics,” *Phys. Rev. D* **83** (2011) 044026, [arXiv:1004.3558](#) [[hep-th](#)].
- [107] M. J. Stott, “Ultralight Bosonic Field Mass Bounds from Astrophysical Black Hole Spin,” [arXiv:2009.07206](#) [[hep-ph](#)].
- [108] V. M. Mehta, M. Demirtas, C. Long, D. J. E. Marsh, L. Mcallister, and M. J. Stott, “Superradiance Exclusions in the Landscape of Type IIB String Theory,” [arXiv:2011.08693](#) [[hep-th](#)]. 19
- [109] R. Balkin, J. Serra, K. Springmann, and A. Weiler, “The QCD axion at finite density,” *JHEP* **07** (2020) 221, [arXiv:2003.04903](#) [[hep-ph](#)]. 20
- [110] T. D. Cohen, R. J. Furnstahl, and D. K. Griegel, “Quark and gluon condensates in nuclear matter,” *Phys. Rev.* **C45** (1992) 1881–1893. 20
- [111] J. Gasser, H. Leutwyler, and M. Sainio, “Form-factor of the sigma term,” *Phys. Lett. B* **253** (1991) 260–264. 20
- [112] **Borexino** Collaboration, G. Bellini *et al.*, “Final results of Borexino Phase-I on low energy solar neutrino spectroscopy,” *Phys. Rev.* **D89** no. 11, (2014) 112007, [arXiv:1308.0443](#) [[hep-ex](#)]. 21
- [113] K. Mukai, “X-ray Emissions from Accreting White Dwarfs: a Review,” *Publ. Astron. Soc. Pac.* **129** no. 976, (2017) 062001, [arXiv:1703.06171](#) [[astro-ph.HE](#)]. 21

- [114] M. Hoferichter, J. Ruiz de Elvira, B. Kubis, and U.-G. Meißner, “High-Precision Determination of the Pion-Nucleon σ Term from Roy-Steiner Equations,” *Phys. Rev. Lett.* **115** (2015) 092301, [arXiv:1506.04142 \[hep-ph\]](#). 21
- [115] J. Alarcon, J. Martin Camalich, and J. Oller, “The chiral representation of the πN scattering amplitude and the pion-nucleon sigma term,” *Phys. Rev. D* **85** (2012) 051503, [arXiv:1110.3797 \[hep-ph\]](#). 21
- [116] J. H. Taylor and J. M. Weisberg, “A new test of general relativity: Gravitational radiation and the binary pulsar PS R 1913+16,” *Astrophys. J.* **253** (1982) 908–920. 23
- [117] M. Burgay *et al.*, “An Increased estimate of the merger rate of double neutron stars from observations of a highly relativistic system,” *Nature* **426** (2003) 531–533, [arXiv:astro-ph/0312071 \[astro-ph\]](#).
- [118] C. M. Will, “The Confrontation between general relativity and experiment,” *Living Rev. Rel.* **4** (2001) 4, [arXiv:gr-qc/0103036 \[gr-qc\]](#). 23
- [119] **LIGO Scientific Collaboration and Virgo Collaboration** Collaboration, B. P. Abbott *et al.*, “Gw170817: Observation of gravitational waves from a binary neutron star inspiral,” *Phys. Rev. Lett.* **119** (Oct, 2017) 161101. <https://link.aps.org/doi/10.1103/PhysRevLett.119.161101>. 23
- [120] L. Hui, J. P. Ostriker, S. Tremaine, and E. Witten, “Ultralight scalars as cosmological dark matter,” *Phys. Rev. D* **95** no. 4, (2017) 043541, [arXiv:1610.08297 \[astro-ph.CO\]](#). 23
- [121] M. Cvitković, A.-S. Smith, and J. Pande, “Asymptotic expansions of the hypergeometric function with two large parameters—application to the partition function of a lattice gas in a field of traps,” *Journal of Physics A: Mathematical and Theoretical* **50** no. 26, (Jun, 2017) 265206. <https://doi.org/10.1088%2F1751-8121%2Faa7213>. 27
- [122] F. Olver, D. Lozier, R. Boisvert, and C. Clark, “Nist handbook of mathematical functions,”. 27