

**DEUTSCHES ELEKTRONEN-SYNCHROTRON**  
Ein Forschungszentrum der Helmholtz-Gemeinschaft



DESY 21-013  
DO-TH 21/04  
TTP21-004  
SAGEX-21-03  
arXiv:2102.12237  
February 2021

**The QED Initial State Corrections to  
the Forward-Backward Asymmetry of  
 $e^+e^- \rightarrow \gamma^*/Z^0^*$  to Higher Orders**

J. Blümlein, A. De Freitas

*Deutsches Elektronen-Synchrotron DESY, Zeuthen*

K. Schönwald

*Institut für Theoretische Teilchenphysik, Karlsruher Institut für Technologie (KIT), Karlsruhe*

ISSN 0418–9833

**PLATANENALLEE 6 — 15738 ZEUTHEN**

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your reports and preprints are promptly included in the  
HEP literature database  
send them to (if possible by air mail):

<p>DESY Zentralbibliothek Notkestraße 85 22607 Hamburg Germany</p>	<p>DESY Bibliothek Platanenallee 6 15738 Zeuthen Germany</p>
--	--

# The QED Initial State Corrections to the Forward-Backward Asymmetry of $e^+e^- \rightarrow \gamma^*/Z^0^*$ to Higher Orders

J. Blümlein<sup>a</sup>, A. De Freitas<sup>a</sup>, and K. Schönwald<sup>b</sup>

<sup>a</sup> *Deutsches Elektronen–Synchrotron, DESY,  
Platanenallee 6, D–15738 Zeuthen, Germany*

<sup>b</sup> *Institut für Theoretische Teilchenphysik,  
Karlsruher Institut für Technologie (KIT) D–76128 Karlsruhe, Germany*

## Abstract

The QED initial state corrections are calculated to the forward–backward asymmetry for  $e^+e^- \rightarrow \gamma^*/Z^0^*$  in the leading logarithmic approximation to  $O(\alpha^6 L^6)$  extending the known corrections up to  $O(\alpha^2 L^2)$  in analytic form. We use the method of massive on-shell operator matrix elements and present the radiators both in Mellin- $N$  and momentum fraction  $z$ -space. Numerical results are presented for various energies around the  $Z$ -peak by also including energy cuts. These corrections are of relevance for the precision measurements at the FCC-ee.

Dedicated to the Memory of Tini Veltman, who made it possible to probe the Standard Model at high precision.

# 1 Introduction

The measurement of the forward–backward asymmetry for the process  $e^+e^- \rightarrow \gamma^*/Z^*$  provides an excellent possibility to determine the running fine structure constant  $\alpha_{\text{QED}}(s) = 4\pi a(s)$  near  $s = M_Z^2$  at high precision. At the planned future  $e^+e^-$  facilities which operate at high energy and at large luminosity, like the ILC, CLIC [1–4], the FCC\\_ee [5], and also muon colliders [6], if operating in the vicinity of the  $Z$ -peak, one will obtain highly precise data. Any measurement based on these data needs a theoretical description of even higher precision [7]. Recently, higher order inclusive corrections were calculated for  $e^+e^- \rightarrow \gamma^*/Z^*$  to  $O(\alpha_s^6 L^5)$  up to the first three orders in  $L$  at the respective order in  $\alpha$  in Refs. [8–11] confirming the results to  $O(\alpha^2)$  in Ref. [12] and correcting Refs. [13, 14], where  $L = \ln(s/m_e^2)$ . This will lead to a change of the analysis codes TOPAZ [15, 16] and ZFITTER [17] and may require a re-analysis of the data taken at LEP [18].

The first order QED initial state radiative (ISR) corrections to the forward–backward asymmetry have been mutually calculated, see [16, 19–24]. Furthermore, the initial-final state interference and final state corrections are known at this order, cf. [16] for a survey. Furthermore, electroweak [25, 26] and QCD corrections [27] have also been calculated. Starting at  $O(\alpha^2)$  also the contributions to the leading order series of  $O(\alpha^k L^k)$ ,  $k \geq 1$ , with  $L = \ln(s/m_e^2)$ <sup>1</sup> receive besides the inclusive contribution another one, related to the angular structure, being present in all sub-leading terms as well. Yet the  $O(\alpha^k L^k)$  terms are universal, since they do not depend on the process-dependent Wilson coefficients, cf. [11]. The inclusive terms were computed in [11] and include besides the anomalous dimensions and massive OMEs also the inclusive massless Wilson coefficients up to  $O(\alpha^2)$  [29, 30].

In the present paper we will calculate the angular dependent leading logarithmic contributions to the radiators to  $O(\alpha^6 L^6)$  as a first specific contribution which emerges for the forward–backward asymmetry. To 2nd order, these corrections were obtained in [23]. The corresponding radiators can be represented by iterated integrals over the alphabet of the harmonic polylogarithms [31] and cyclotomic harmonic polylogarithms [32] for cyclotomy  $c = 4$ . We also determine efficient representations for these quantities allowing a fast numerical analysis and present the corresponding corrections for the forward–backward asymmetry in the vicinity of the  $Z$ -resonance. These corrections, unlike the inclusive ones, do not lead to distribution–valued radiators.

The paper is organized as follows. In Section 2 we calculate the higher order QED initial state corrections to the forward–backward asymmetry. Here we use the packages **Sigma** [33, 34] and **HarmonicSums** [31, 32, 35–38]. The leading logarithmic radiators of  $H_{FB}(z)$  to  $O(\alpha^6 L^6)$  of the angular-dependent terms are presented in Section 3. We also derive the expansions of the radiators in the regions  $z \rightarrow 0$  and  $z \rightarrow 1$ . Numerical and phenomenological results are presented in Section 4. In Appendix A we give a brief account of the aforementioned radiators in Mellin  $N$ -space. The radiators are given in computer-readable form in an attachment to this paper.

## 2 The Forward-Backward Asymmetry around the $Z$ -Peak

The forward-backward asymmetry is formed out of the partial cross sections integrating over the angle  $\theta$  for the forward and backward hemispheres separately,

$$\sigma_F = 2\pi \int_0^1 d \cos(\theta) \frac{d\sigma}{d\Omega}, \quad \sigma_B = 2\pi \int_{-1}^0 d \cos(\theta) \frac{d\sigma}{d\Omega}. \quad (1)$$

---

<sup>1</sup>The leading order contributions to the direct terms were obtain in [28].

The angle  $\theta$  is defined between the incoming electron  $e^-$  and the outgoing muon  $\mu^-$  from  $\gamma^*/Z^*$  decay. The forward-backward asymmetry is defined by

$$A_{\text{FB}}(s) = \frac{\sigma_F(s) - \sigma_B(s)}{\sigma_T(s)}, \quad (2)$$

with  $\sigma_T(s) = \sigma_F(s) + \sigma_B(s)$ . At Born level this reduces to [39]

$$\sigma_{FB}^{(0)}(s) = \sigma_F^{(0)}(s) - \sigma_B^{(0)}(s) = \frac{\pi\alpha^2}{s} N_{C,f} \left(1 - \frac{4m_f^2}{s}\right) G_3(s), \quad (3)$$

$$\sigma_T^{(0)}(s) = \sigma_F^{(0)}(s) + \sigma_B^{(0)}(s) = \frac{4\pi\alpha^2}{3s} N_{C,f} \sqrt{1 - \frac{4m_f^2}{s}} \left[ \left(1 + \frac{2m_f^2}{s}\right) G_1(s) - 6\frac{m_f^2}{s} G_2(s) \right], \quad (4)$$

with  $m_f$  the final state fermion mass,  $m_f \equiv m_\mu$ .  $N_{C,f}$  is the number of colors of the final state fermion, with  $N_{C,f} = 1$  in the present case.  $s$  is the cms energy, and the effective couplings  $G_i(s)|_{i=1\dots 3}$  read

$$G_1(s) = G_{1,1} + G_{1,2} + G_{1,3} = Q_e^2 Q_f^2 + 2Q_e Q_f v_e v_f \text{Re}[\chi_Z(s)] + (v_e^2 + a_e^2)(v_f^2 + a_f^2) |\chi_Z(s)|^2, \quad (5)$$

$$G_2(s) = (v_e^2 + a_e^2) a_f^2 |\chi_Z(s)|^2, \quad (6)$$

$$G_3(s) = G_{3,1} + G_{3,2} = 2Q_e Q_f a_e a_f \text{Re}[\chi_Z(s)] + 4v_e v_f a_e a_f |\chi_Z(s)|^2. \quad (7)$$

For later use we define

$$\sigma_T^{\gamma\gamma} = F_1 G_{1,1}, \quad \sigma_T^{\gamma Z} = \sigma_T^{Z\gamma} = \frac{1}{2} F_1 G_{1,2}, \quad \sigma_T^{ZZ} = F_1 G_{1,3} - 6F_3 G_2 \quad (8)$$

$$\sigma_{FB}^{\gamma Z} = \sigma_{FB}^{Z\gamma} = \frac{1}{2} F_2 G_{3,1}, \quad \sigma_{FB}^{ZZ} = F_2 G_{3,2}. \quad (9)$$

with

$$\begin{aligned} F_1 &= \frac{4\pi\alpha^2}{3s} N_{C,f} \sqrt{1 - \frac{4m_f^2}{s}} \left(1 + \frac{2m_f^2}{s}\right), & F_2 &= \frac{\pi\alpha^2}{s} N_{C,f} \left(1 - \frac{4m_f^2}{s}\right), \\ F_3 &= \frac{4\pi\alpha^2}{3s} N_{C,f} \left(1 - \frac{4m_f^2}{s}\right) \frac{m_f^2}{s}. \end{aligned} \quad (10)$$

The reduced  $Z$ -propagator is given by

$$\chi_Z(s) = \frac{s}{s - M_Z^2 + iM_Z\Gamma_Z}, \quad (11)$$

where  $M_Z$  and  $\Gamma_Z$  are the mass and the width of the  $Z$  boson.  $Q_{e,f}$  are the electromagnetic charges of the electron ( $Q_e = -1$ ) and the final state fermion, respectively, and the electro-weak couplings  $v_i$  and  $a_i$  read

$$v_e = \frac{1}{\sin\theta_w \cos\theta_w} [I_{w,e}^3 - 2Q_e \sin^2\theta_w], \quad a_e = \frac{1}{\sin\theta_w \cos\theta_w} I_{w,e}^3, \quad (12)$$

$$v_f = \frac{1}{\sin\theta_w \cos\theta_w} [I_{w,f}^3 - 2Q_f \sin^2\theta_w], \quad a_f = \frac{1}{\sin\theta_w \cos\theta_w} I_{w,f}^3. \quad (13)$$

$\theta_w$  denotes the weak mixing angle, and  $I_{w,i}^3 = \pm 1/2$  the third component of the weak isospin for up and down particles, respectively.

When accounting for initial-state-radiation, one obtains the following representation of the  $A_{FB}$  introducing the radiators  $H_e^{LL}$  and  $H_{FB}^{LL}$  [23], using the notation in [19],

$$A_{FB}(s) = \frac{1}{\sigma_T(s)} \int_{z_0}^1 dz \frac{4z}{(1+z)^2} \tilde{H}_e^{LL}(z) \sigma_{FB}^{(0)}(zs), \quad \sigma_T(s) = \int_{z_0}^1 dz H_e(z) \sigma_T^{(0)}(zs), \quad (14)$$

with

$$\tilde{H}_e^{LL}(z) = [H_e^{LL}(z) + H_{FB}^{LL}(z)]. \quad (15)$$

The normalization factor  $\sigma_T(s)$  will be calculated considering all corrections derived in Ref. [11]. We will consider different values for the threshold  $z_0$ . Like in Refs. [8–10]  $z_0$  is chosen as  $z_0 = 4m_\tau^2/s$ , with  $m_\tau$  the mass of the  $\tau$  lepton. Another choice will be  $z_0 = 0.99$  or  $0.999$  in accordance with [40]. The variable  $z$  is given by  $z = s'/s$ , where  $s'$  denotes the virtuality of the gauge boson  $\gamma^*$  or  $Z^*$ . Furthermore,  $H_e(z)$  is the radiator in the inclusive case, given in Ref. [11] and  $H_{FB}^{LL}(z)$  denotes the leading-log radiator in the angular dependent case up to  $O(\alpha^6 L^6)$ ,

$$H_{FB}^{LL}(z) = \int_0^1 dx_1 \int_0^1 dx_2 \left( \frac{(1+z)^2}{(x_1+x_2)^2} - 1 \right) \Gamma_{ee}^{LL}(x_2) \Gamma_{ee}^{LL}(x_1) \delta(x_1 x_2 - z). \quad (16)$$

Here  $\Gamma_{ee}^{LL}(x)$  is the leading-log operator matrix element. The product  $\Gamma_{ee}^{LL}(x_1) \Gamma_{ee}^{LL}(x_2)$  is consistently expanded up to  $O(\alpha^6 L^6)$ . The latter radiator does only contain universal contributions. The radiators obey the expansion

$$H_i(z) = \delta_{ie} \delta(1-z) + \sum_{k=1}^{\infty} a^k \sum_{l=0}^k L^l H_i^{(k,l)}(z), \quad i = e, FB, \quad (17)$$

$$H_i^{LL}(z) = \delta_{ie} \delta(1-z) + \sum_{k=1}^{\infty} (aL)^k H_i^{(k),LL}(z), \quad (18)$$

where  $a = \alpha/(4\pi)$  and  $\alpha$  denotes the fine structure constant.

We proceed in the following way in order to evaluate Eq. (16) analytically. First, we calculate the Mellin transform

$$\mathcal{M}[H_{FB}^{LL}(z)](n) = \int_0^1 dz z^n H_{FB}^{LL}(z) = \int_0^1 dx_1 \int_0^1 dx_2 x_1^n x_2^n \left( \frac{(1+x_1 x_2)^2}{(x_1+x_2)^2} - 1 \right) \Gamma_{ee}^{LL}(x_2) \Gamma_{ee}^{LL}(x_1). \quad (19)$$

Since this integral is not suited to be integrated with the package **HarmonicSums** directly, we compute the generating function

$$\begin{aligned} \mathcal{G}[H_{FB}^{LL}(z)](t) &= \sum_{n=0}^{\infty} t^n \mathcal{M}[H_{FB}^{LL}(z)](n) \\ &= \int_0^1 dx_1 \int_0^1 dx_2 \frac{1}{1-t x_1 x_2} \left( \frac{(1+x_1 x_2)^2}{(x_1+x_2)^2} - 1 \right) \Gamma_{ee}^{LL}(x_2) \Gamma_{ee}^{LL}(x_1), \end{aligned} \quad (20)$$

which resums the Mellin-kernel into a denominator which can be easily integrated over. After the integration over  $x_1$  and  $x_2$  we are left with generalized iterated integrals evaluated at argument  $z = 1$  which contain the parameter  $t$  in their letters. We use differential equations to pull this parameter into the argument. This is straightforward since the limit  $t \rightarrow 0$  always exists and is easily expressed in terms of known constants. Afterwards we can use the `HarmonicSums` command `GetMoment` to get the  $n$ -space expression and `GeneralInvMellin` to arrive at the final result. The radiators are consistently expanded in  $(aL)$  to  $O((aL)^6)$  and are expressed using the variable  $\sqrt{z}$  rather than  $z$  to obtain a unique representation concerning the contributing iterated integrals.

At  $O(\alpha)$  the complete QED initial state corrections [16, 21, 22] are known and are given by

$$A_{FB}^{(1)}(s) = \frac{1}{\sigma_T(s)} \int_{z_0}^1 dz \frac{4z}{(1+z)^2} a(s) [\tilde{H}_e^{(1),LL}(z)L + \tilde{H}_e^{(1,0)}(z)] \sigma_T^{(0)}(zs), \quad (21)$$

and

$$H_e^{(1),LL}(z) = 4 \left[ \frac{1+z^2}{1-z} \right]_+ \quad (22)$$

$$\tilde{H}_e^{(1,0)}(z) = 4 \left[ - \left[ \frac{1+z^2}{1-z} \right]_+ + \left( 2\zeta_2 - \frac{1}{2} \right) \delta(1-z) + \frac{1+z^2}{1-z} [2\ln(1+z) - 2\ln(2) - \ln(z)] \right]. \quad (23)$$

### 3 The leading-log radiators $H_{FB}^{LL}$

We obtain the following leading-log radiators  $H_{FB}^{(k),LL}$  :

$$H_{FB}^{(1),LL}(z) = 0 \quad (24)$$

$$H_{FB}^{(2),LL}(z) = \frac{2(1-z)(1+z)^2}{z} + 2\pi \frac{(1-z)^2}{\sqrt{z}} - 8(1+z)H_0 - 8(1-z)^2 \frac{H_{\{4,0\}}}{\sqrt{z}} \quad (25)$$

$$\begin{aligned} H_{FB}^{(3),LL}(z) = & -\frac{16(1-z)(4+11z+4z^2)}{3z} - \pi \left[ \frac{4(2-3z-2z^2-3z^3+2z^4)}{3z^{3/2}} \right. \\ & + \frac{4(1-z)(1+5z)}{\sqrt{z}} H_0 + \frac{16(1-z)^2}{\sqrt{z}} H_{\{4,1\}} \Big] + \left[ \frac{4(1+z)(5-18z-19z^2)}{3z} \right. \\ & - \frac{16(1-z)(1-7z)}{\sqrt{z}} H_{\{4,0\}} - 96(1+z)H_{\{4,1\}} \Big] H_0 - 8(1+z)H_0^2 \\ & + \left[ \frac{16(1-z)(1+z)^2}{z} - \frac{64(1-z)^2}{\sqrt{z}} H_{\{4,0\}} \right] H_1 + \left[ \frac{16(2-3z-2z^2-3z^3+2z^4)}{3z^{3/2}} \right. \\ & + \frac{64(1-z)^2}{\sqrt{z}} H_{\{4,1\}} + \left[ -\frac{16(1-z)(1+z)^2}{z} + 64(1+z)H_0 \right. \\ & + \frac{64(1-z)^2}{\sqrt{z}} H_{\{4,0\}} \Big] H_{-1} - 64(1+z)H_{0,1} + \frac{32(1-z)^2}{\sqrt{z}} H_{0,\{4,0\}} + 96(1+z)H_{0,\{4,1\}} \\ & + \frac{64(1-z)^2}{\sqrt{z}} H_{1,\{4,0\}} - \frac{64(1-z)^2}{\sqrt{z}} H_{\{4,0\},\{4,1\}} - 64(1+z)H_{-1,0} \\ & \left. \left. - \frac{64(1-z)^2}{\sqrt{z}} H_{-1,\{4,0\}} + 20(1+z)\zeta_2 \right] \right] \end{aligned} \quad (26)$$

with  $H_{\vec{w}} \equiv H_{\vec{w}}(\sqrt{z})$  and  $\zeta_k, k \in \mathbb{N}, k \geq 2$  are the values of Riemann's  $\zeta$  function at integer argument. The iterated integrals are given by

$$H_{b,\vec{a}}(z) = \int_0^z dy f_b(y) H_{\vec{a}}(y), \quad H_\emptyset = 1. \quad (27)$$

The indices  $b, a_i$  refer to the letters in the following alphabet

$$\mathfrak{A} = \left\{ f_0 = \frac{1}{z}, f_1 = \frac{1}{1-z}, f_{-1} = \frac{1}{1+z}, f_{\{4,0\}} = \frac{1}{1+z^2}, f_{\{4,1\}} = \frac{z}{1+z^2} \right\}, \quad (28)$$

where the last two letters in (28) are cyclotomic. The function  $H_{FB}^{(2),LL}(z)$  has been calculated in [23] in an equivalent representation. The radiators  $H_{FB}^{(4-6),LL}(z)$  are too voluminous to be presented here and are given in computer-readable form in the attachment.

We finally illustrate the structure of the radiators expanding them for their leading small and a few large  $z$  terms. One obtains

$$zH_{FB}^{(2),LL}(z) \simeq 2 + O(\sqrt{z}) \quad (29)$$

$$zH_{FB}^{(3),LL}(z) \simeq -\frac{8\pi}{3\sqrt{z}} + \frac{10}{3} \ln(z) - \frac{32}{3} + O(\sqrt{z} \ln(z)) \quad (30)$$

$$\begin{aligned} zH_{FB}^{(4),LL}(z) \simeq & \frac{32}{9z} - \frac{11}{6} \ln^2(z) + \frac{10}{27} \ln(z) - 38 - 43\zeta_2 \\ & + \frac{\pi}{\sqrt{z}} \left( \frac{4}{3} \ln(z) + \frac{328}{27} \right) + O(\sqrt{z} \ln^2(z)) \end{aligned} \quad (31)$$

$$\begin{aligned} zH_{FB}^{(5),LL}(z) \simeq & -\frac{512}{81z} + \frac{5}{36} \ln^3(z) + \frac{97}{18} \ln^2(z) + \left[ -\frac{299}{27} + \frac{215\zeta_2}{18} \right] \ln(z) + \frac{72283}{243} \\ & + \frac{2551}{27} \zeta_2 + \frac{320}{9} \zeta_3 + \frac{\pi}{\sqrt{z}} \left( \frac{2}{3} \ln^2(z) + \frac{92}{27} \ln(z) + \frac{460}{81} \right. \\ & \left. + \frac{4}{9} \zeta_2 \right) + O(\sqrt{z} \ln^3(z)) \end{aligned} \quad (32)$$

$$\begin{aligned} zH_{FB}^{(6),LL}(z) \simeq & -\frac{10240}{729z} - \frac{256}{81z} \ln(z) + \frac{97}{720} \ln^4(z) + \frac{79}{162} \ln^3(z) + \left( \frac{17059}{2430} - \frac{19}{180} \zeta_2 \right) \ln^2(z) \\ & + \left( -\frac{159109}{7290} - \frac{2825}{81} \zeta_2 + \frac{136}{9} \zeta_3 \right) \ln(z) - \frac{2459744}{10935} + \frac{87173}{1215} \zeta_2 - \frac{53216}{405} \zeta_3 \\ & + \frac{5659}{300} \zeta_2^2 + \frac{\pi}{\sqrt{z}} \left( -\frac{1}{18} \ln^3(z) - \frac{32}{9} \ln^2(z) + \left[ -\frac{39212}{1215} - \frac{73}{9} \zeta_2 \right] \ln(z) \right. \\ & \left. - \frac{401806}{3645} - \frac{15904}{405} \zeta_2 - \frac{128}{9} \zeta_3 \right) + O(\sqrt{z} \ln^4(z)). \end{aligned} \quad (33)$$

For the large  $z$  representation we set  $u = 1 - \sqrt{z}$  and obtain

$$H_{FB}^{(2),LL}(z) \simeq 32u - 16u^2 + \frac{112}{3}u^3 + \frac{56}{3}u^4 + O(u^5) \quad (34)$$

$$\begin{aligned} H_{FB}^{(3),LL}(z) \simeq & \frac{64}{3}u - \frac{416}{3}u^2 + \frac{1408}{9}u^3 - \frac{736}{9}u^4 + \left( -256u + 128u^2 - \frac{896}{3}u^3 - \frac{448}{3}u^4 \right) \\ & \times \ln(2u) + O(u^5) \end{aligned} \quad (35)$$

$$H_{FB}^{(4),LL}(z) \simeq \frac{27328}{27}u - \frac{13664}{27}u^2 + \frac{162080}{81}u^3 + \frac{36112}{81}u^4 + \left( 1024u - 512u^2 \right.$$

$$+ \frac{3584}{3}u^3 + \frac{1792}{3}u^4 \Big) (\ln^2(2u) - \zeta_2) + \left( 1024 - \frac{3328}{3}u + \frac{2176}{3}u^2 \right) \\ \times u^2 \ln(2u) + O(u^5) \quad (36)$$

$$H_{FB}^{(5),LL}(z) \simeq \frac{387712}{81}u - \frac{523328}{81}u^2 + \frac{2954144}{243}u^3 - \frac{328688}{243}u^4 + \left( \frac{2048}{3} + \frac{11264}{3}u \right. \\ \left. - \frac{34304}{9}u^2 + \frac{28928}{9}u^3 \right) u (\zeta_2 - \ln^2(2u)) + \left( -\frac{8192}{3} + \frac{4096}{3}u - \frac{28672}{9}u^2 \right. \\ \left. - \frac{14336}{9}u^3 \right) u (2\zeta_3 - 3\zeta_2 \ln(2u) + \ln^3(2u)) + \left( -\frac{73216}{9} + \frac{30464}{9}u \right. \\ \left. - \frac{419840}{27}u^2 - \frac{108544}{27}u^3 \right) u \ln(2u) + O(u^5) \quad (37)$$

$$H_{FB}^{(6),LL}(z) \simeq \frac{153977536}{3645}u - \frac{136846688}{3645}u^2 + \frac{1140995264}{10935}u^3 + \frac{60397216}{10935}u^4 + \left( -\frac{894976}{27} \right. \\ \left. + \frac{300032}{27}u - \frac{24477184}{405}u^2 - \frac{7326464}{405}u^3 \right) u (\zeta_2 - \ln^2(2u)) + \left( \frac{16384}{5} - \frac{8192}{5}u \right. \\ \left. + \frac{57344}{15}u^2 + \frac{28672}{15}u^3 \right) u \left( \zeta_2^2 + \frac{5}{3} \ln^4(2u) \right) + \left( \frac{32768}{9} + \frac{81920}{9}u - \frac{1132544}{135}u^2 \right. \\ \left. + \frac{1276928}{135}u^3 \right) u (2\zeta_3 - 3\zeta_2 \ln(2u) + \ln^3(2u)) + \left( -\frac{886784}{27} + \frac{1338368}{27}u \right. \\ \left. - \frac{35939456}{405}u^2 + \frac{5414336}{405}u^3 - \left[ -\frac{131072}{3} + \frac{65536}{3}u - \frac{458752}{9}u^2 \right. \right. \\ \left. \left. - \frac{229376}{9}u^3 \right] \zeta_3 \right) u \ln(2u) + \left( -32768 + 16384u - \frac{114688}{3}u^2 - \frac{57344}{3}u^3 \right) \\ \times u \zeta_2 \ln^2(2u) + O(u^5). \quad (38)$$

These terms are the beginning of the respective series used in the numerical representation to which we turn now.

## 4 Numerical results

For the numerical evaluation of the new radiators we calculated 60 terms of the expansions around  $\sqrt{z} = 0$  and  $\sqrt{z} = 1$  using `HarmonicSums`. The constants associated with cyclotomic harmonic polylogarithms for cyclotomy  $c=4$ , which are needed for the expansion around  $\sqrt{z} = 1$  are known at least up to weight  $w = 5$  [32]. However, all constants except of  $\pi$  cancel in the final results. We switch between these two expansions at  $z = 1/4$ , where the absolute difference of the respective approximation is always smaller than  $10^{-12}$ . The usual harmonic polylogarithms are calculated using the implementations in [42, 43]. In the numerical illustration we refer to the electroweak parameters given in [41].

In Figure 1 we illustrate the effect of the initial-state QED corrections on the forward–backward asymmetry for a wider range of cms energies around the  $Z$ -peak with  $\sqrt{s} \in [85, 96]$  GeV to provide an overall impression on the size of the radiative corrections, which will be detailed below.

Let us define

$$\Delta A_{FB} = 1 - \frac{A_{FB}^{(l)}}{A_{FB}^{(0)}}, \quad (39)$$




Figure 1:  $A_{FB}$  and its initial state QED corrections as a function of  $\sqrt{s}$ . Black (dashed) the Born approximation, blue (dotted) the  $O(\alpha)$  improved approximation, red (full) also including the leading-log improvement up to  $O(\alpha^6)$  for  $s'/s \geq 4m_\tau^2/s$ .




Figure 2:  $\Delta A_{FB}$  in % as defined in Eq. (39) as a function of  $\sqrt{s}$ . Black (dashed) the  $O(\alpha)$  improved approximation, blue (dotted) the  $O(\alpha^2 L^2)$  improved approximation, red (full) also including the leading-log improvement up to  $O(\alpha^6)$  for  $s'/s \geq 4m_\tau^2/s$ .

where  $l$  denotes the order to which the initial state radiation is taken into account.  $l = 0$

corresponds to the Born approximation. In Figure 2 we illustrate the relative size of the ISR corrections to the Born cross section.

In Table 1 we illustrate the effect of the different orders of the QED initial state corrections of the forward–backward asymmetry at the  $Z$ -peak and two more values of  $s = s_{\pm}$  [40]. The ISR corrections are not monotonic order by order, even leading to a sign change in the corrected values of  $A_{FB}$  at  $s = M_Z^2$ . Here the ISR corrections are largest.

If we compare the known one-loop corrected values of  $A_{FB}$  with the highest radiative correction calculated in this paper we obtain further corrections of  $-3\%$  for  $s_-$  and  $-1\%$  for  $s_+$ . These values supersede previous estimates, based on assumptions, in [40].

	$A_{FB}(s_-)$	$A_{FB}(M_Z^2)$	$A_{FB}(s_+)$
$\mathcal{O}(\alpha^0)$	-0.3564803	0.0225199	0.2052045
$\mathcal{O}(\alpha L^1)$	-0.2945381	-0.0094232	0.1579347
$\mathcal{O}(\alpha L^0)$	-0.2994478	-0.0079610	0.1611962
$\mathcal{O}(\alpha^2 L^2)$	-0.3088363	0.0014514	0.1616887
$\mathcal{O}(\alpha^3 L^3)$	-0.3080578	0.0000198	0.1627252
$\mathcal{O}(\alpha^4 L^4)$	-0.3080976	0.0001587	0.1625835
$\mathcal{O}(\alpha^5 L^5)$	-0.3080960	0.0001495	0.1625911
$\mathcal{O}(\alpha^6 L^6)$	-0.3080960	0.0001499	0.1625911

Table 1:  $A_{FB}$  evaluated at  $s_- = (87.9 \text{ GeV})^2$ ,  $M_Z^2$  and  $s_+ = (94.3 \text{ GeV})^2$  for the cut  $z > 4m_\tau^2/s$ .

If cuts of  $s'$  are applied the ISR corrections turn out to be smaller, as illustrated in Tables 2 and 3. As noted in [19] also an approximative treatment of the forward–backward asymmetry

$$A_{FB}(s) \approx \frac{1}{\sigma_T(s)} \int_{z_0}^1 dz \frac{4z}{(1+z)^2} H_e(z) \sigma_{FB}^{(0)}(zs), \quad (40)$$

has been used in the literature [24] close to the  $Z$ -peak, where  $H_e(z)$  denotes the inclusive

	$A_{FB}(s_-)$	$A_{FB}(M_Z^2)$	$A_{FB}(s_+)$
$\mathcal{O}(\alpha^0)$	-0.2714847	0.01545540	0.2539839
$\mathcal{O}(\alpha L^1)$	-0.2913715	0.01244012	0.2787800
$\mathcal{O}(\alpha L^0)$	-0.2923065	0.01266498	0.2794039
$\mathcal{O}(\alpha^2 L^2)$	-0.2922610	0.01489549	0.2760952
$\mathcal{O}(\alpha^3 L^3)$	-0.2917702	0.01430419	0.2764488
$\mathcal{O}(\alpha^4 L^4)$	-0.2918380	0.01439578	0.2763865
$\mathcal{O}(\alpha^5 L^5)$	-0.2918313	0.01438608	0.2763934
$\mathcal{O}(\alpha^6 L^6)$	-0.2918318	0.01438681	0.2763929

Table 2:  $A_{FB}$  evaluated at  $s_- = (87.9 \text{ GeV})^2$ ,  $M_Z^2$  and  $s_+ = (94.3 \text{ GeV})^2$  for the cut  $z > 0.99$ .

radiator appearing in case of  $\sigma_T(s)$  [11]. In the sub-leading corrections to the radiators [11], Wilson coefficients appear, which are different for  $\sigma_T$  and  $\sigma_{FB}$ . Therefore the radiator for  $\sigma_T$

cannot be used in its sub-leading terms for this reason. The numerical results in Table 4–6 show, that contributions due to the angular radiators are indeed suppressed if measuring at the  $Z$ -peak w.r.t. to the inclusive ones also in higher orders. This holds quite irrespectively of the cuts in  $s'$ .

	$A_{FB}(s_-)$	$A_{FB}(M_Z^2)$	$A_{FB}(s_+)$
$\mathcal{O}(\alpha^0)$	−0.2683019	0.01538044	0.2576633
$\mathcal{O}(\alpha L^1)$	−0.2905494	0.01623701	0.2797643
$\mathcal{O}(\alpha L^0)$	−0.2913658	0.01630129	0.2805232
$\mathcal{O}(\alpha^2 L^2)$	−0.2899912	0.01655798	0.2787005
$\mathcal{O}(\alpha^3 L^3)$	−0.2898133	0.01641903	0.2787137
$\mathcal{O}(\alpha^4 L^4)$	−0.2898401	0.01645212	0.2786941
$\mathcal{O}(\alpha^5 L^5)$	−0.2898362	0.01644636	0.2786984
$\mathcal{O}(\alpha^6 L^6)$	−0.2898368	0.01644713	0.2786978

Table 3:  $A_{FB}$  evaluated at  $s_- = (87.9 \text{ GeV})^2$ ,  $M_Z^2$  and  $s_+ = (94.3 \text{ GeV})^2$  for the cut  $z > 0.999$ .

	all ( $s_-$ )	ang. ( $s_-$ )	all ( $M_Z^2$ )	ang. ( $M_Z^2$ )	all ( $s_+$ )	ang. ( $s_+$ )
$\mathcal{O}(\alpha^2 L^2)$	$-9 \cdot 10^{-3}$	$-9 \cdot 10^{-5}$	$9 \cdot 10^{-3}$	$-1 \cdot 10^{-5}$	$5 \cdot 10^{-4}$	$-6 \cdot 10^{-5}$
$\mathcal{O}(\alpha^3 L^3)$	$8 \cdot 10^{-4}$	$-4 \cdot 10^{-6}$	$-1 \cdot 10^{-3}$	$-8 \cdot 10^{-7}$	$1 \cdot 10^{-3}$	$-3 \cdot 10^{-6}$
$\mathcal{O}(\alpha^4 L^4)$	$-4 \cdot 10^{-5}$	$-3 \cdot 10^{-7}$	$1 \cdot 10^{-4}$	$-7 \cdot 10^{-8}$	$-1 \cdot 10^{-4}$	$-7 \cdot 10^{-8}$
$\mathcal{O}(\alpha^5 L^5)$	$2 \cdot 10^{-6}$	$-1 \cdot 10^{-8}$	$-9 \cdot 10^{-6}$	$-4 \cdot 10^{-9}$	$8 \cdot 10^{-6}$	$1 \cdot 10^{-8}$
$\mathcal{O}(\alpha^6 L^6)$	$-6 \cdot 10^{-8}$	$-6 \cdot 10^{-10}$	$4 \cdot 10^{-7}$	$-1 \cdot 10^{-10}$	$-3 \cdot 10^{-8}$	$2 \cdot 10^{-9}$

Table 4: Contribution to  $A_{FB}$  at different orders of  $\alpha L$  evaluated at  $s_- = (87.9 \text{ GeV})^2$ ,  $M_Z^2$  and  $s_+ = (94.3 \text{ GeV})^2$  for the cut  $z > 4m_\tau^2/s$ . all: full contributions, ang.: only the angular dependent contributions.

	all ( $s_-$ )	ang. ( $s_-$ )	all ( $M_Z^2$ )	ang. ( $M_Z^2$ )	all ( $s_+$ )	ang. ( $s_+$ )
$\mathcal{O}(\alpha^2 L^2)$	$5 \cdot 10^{-5}$	$-5 \cdot 10^{-8}$	$2 \cdot 10^{-3}$	$-2 \cdot 10^{-9}$	$-3 \cdot 10^{-3}$	$5 \cdot 10^{-8}$
$\mathcal{O}(\alpha^3 L^3)$	$5 \cdot 10^{-4}$	$-3 \cdot 10^{-8}$	$-6 \cdot 10^{-4}$	$-1 \cdot 10^{-9}$	$4 \cdot 10^{-4}$	$3 \cdot 10^{-8}$
$\mathcal{O}(\alpha^4 L^4)$	$-7 \cdot 10^{-5}$	$-9 \cdot 10^{-9}$	$9 \cdot 10^{-5}$	$-3 \cdot 10^{-10}$	$-6 \cdot 10^{-5}$	$9 \cdot 10^{-9}$
$\mathcal{O}(\alpha^5 L^5)$	$7 \cdot 10^{-6}$	$-2 \cdot 10^{-9}$	$-1 \cdot 10^{-5}$	$-4 \cdot 10^{-11}$	$7 \cdot 10^{-6}$	$2 \cdot 10^{-9}$
$\mathcal{O}(\alpha^6 L^6)$	$-5 \cdot 10^{-7}$	$-2 \cdot 10^{-10}$	$7 \cdot 10^{-7}$	$-4 \cdot 10^{-12}$	$-5 \cdot 10^{-7}$	$2 \cdot 10^{-10}$

Table 5: Contribution to  $A_{FB}$  at different orders of  $\alpha L$  evaluated at  $s_- = (87.9 \text{ GeV})^2$ ,  $M_Z^2$  and  $s_+ = (94.3 \text{ GeV})^2$  for the cut  $z > 0.99$ . all: full contributions, ang.: only the angular dependent contributions.

For completeness we also quantify the final state and initial–final interference contributions at  $O(\alpha)$ , cf. e.g. [16,19,21,22] for the respective numerator,  $\sigma_{FB}$ , and denominator,  $\sigma_T$  contributions

to  $A_{FB} = \sigma_{FB}/\sigma_T$ . The inclusive final state correction to  $\sigma_{FB}$  and  $\sigma_T$  imply the factors

$$1 + \delta_{FB}^F, \quad \text{with } \delta_{FB}^F = 0 \quad \text{and} \quad 1 + \delta_T^F, \quad \text{with } \delta_T^F = 3a \quad (41)$$

for the  $\mu^+\mu^-$  final state. The multiplicative numerical correction to  $A_{FB}$  is 0.998149.

The initial-final interference term [16, 22] has a more involved representation compared to the final state corrections, since there are different correction terms to the different pieces of the angular averaged Born cross section. In particular these terms are important because of logarithmically large terms of  $\ln(1 - s'_{\min}/s)$  in the region  $s'_{\min}$  close to  $s$  as chosen in [40]. The correction term  $1 + \delta_{FB}^{FI}$  modifying  $A_{FB}$  to  $O(\alpha)$  is given by

$$\begin{aligned} \delta_{FB}^{FI} &= -\frac{1}{\sigma_T^{\text{Born}}} \left\{ \sigma_{FB}^{\gamma Z} \text{Re}[R_T^{\gamma\gamma} + R_T^{ZZ}] + \sigma_{FB}^{ZZ} \text{Re}[R_T^{ZZ}] \right\} \\ &\quad + \frac{1}{A_{FB}} \left\{ \text{Re}[R_{FB}^{\gamma\gamma}] \frac{\sigma_T^{\gamma\gamma}}{\sigma_T^{\text{Born}}} + \text{Re}[R_{FB}^{\gamma\gamma} + R_{FB}^{ZZ}] \frac{\sigma_T^{\gamma Z}}{\sigma_T^{\text{Born}}} + \text{Re}[R_{FB}^{ZZ}] \frac{\sigma_T^{ZZ}}{\sigma_T^{\text{Born}}} \right\}, \end{aligned} \quad (42)$$

with  $R_A^{ij}$ ,  $A = T, FB$ ,  $i, j = \gamma, Z$ , the corresponding radiators [16, 22, 26] and

$$\delta_{FB}^{FI} = -0.0821 \quad \text{for } \sqrt{s} = 87.9 \text{ GeV} \quad s'/s > 0.99 \quad (43)$$

$$\delta_{FB}^{FI} = +0.0928 \quad \text{for } \sqrt{s} = 94.3 \text{ GeV} \quad s'/s > 0.99 \quad (44)$$

$$\delta_{FB}^{FI} = -0.1954 \quad \text{for } \sqrt{s} = 87.9 \text{ GeV} \quad s'/s > 0.999 \quad (45)$$

$$\delta_{FB}^{FI} = +0.2223 \quad \text{for } \sqrt{s} = 94.3 \text{ GeV} \quad s'/s > 0.999 \quad (46)$$

The tight cuts in (43–46) imply a correction term of up to  $\pm 9\%$  for  $z > 0.99$  and up to  $\pm 22\%$  for  $z > 0.999$ .

	all ( $s_-$ )	ang. ( $s_-$ )	all ( $M_Z^2$ )	ang. ( $M_Z^2$ )	all ( $s_+$ )	ang. ( $s_+$ )
$\mathcal{O}(\alpha^2 L^2)$	$1 \cdot 10^{-3}$	$-5 \cdot 10^{-10}$	$3 \cdot 10^{-4}$	$2 \cdot 10^{-11}$	$-2 \cdot 10^{-3}$	$5 \cdot 10^{-10}$
$\mathcal{O}(\alpha^3 L^3)$	$2 \cdot 10^{-4}$	$-4 \cdot 10^{-10}$	$-1 \cdot 10^{-4}$	$2 \cdot 10^{-11}$	$1 \cdot 10^{-5}$	$4 \cdot 10^{-10}$
$\mathcal{O}(\alpha^4 L^4)$	$-3 \cdot 10^{-5}$	$-2 \cdot 10^{-10}$	$3 \cdot 10^{-5}$	$9 \cdot 10^{-12}$	$-2 \cdot 10^{-5}$	$2 \cdot 10^{-10}$
$\mathcal{O}(\alpha^5 L^5)$	$4 \cdot 10^{-6}$	$-5 \cdot 10^{-11}$	$-6 \cdot 10^{-6}$	$3 \cdot 10^{-12}$	$4 \cdot 10^{-6}$	$5 \cdot 10^{-11}$
$\mathcal{O}(\alpha^6 L^6)$	$-5 \cdot 10^{-7}$	$-1 \cdot 10^{-11}$	$8 \cdot 10^{-7}$	$5 \cdot 10^{-13}$	$-6 \cdot 10^{-7}$	$1 \cdot 10^{-11}$

Table 6: Contribution to  $A_{FB}$  at different orders of  $\alpha L$  evaluated at  $s_- = (87.9 \text{ GeV})^2$ ,  $M_Z^2$  and  $s_+ = (94.3 \text{ GeV})^2$  for the cut  $z > 0.999$ . all: full contributions, ang.: only the angular dependent contributions.

## A $N$ space results

In the following we are presenting the first radiators in Mellin  $N$ -space.

$$H_{FB}^{(1),LL}(N) = 0 \quad (47)$$

$$\begin{aligned} H_{FB}^{(2),LL}(N) &= \frac{8(3N^2 + 3N - 1)P_1}{(N-1)N^2(N+1)^2(N+2)(2N-1)(2N+3)} \\ &\quad - \frac{32(4N^2 + 4N - 1)(-1)^N}{(2N-1)(2N+1)(2N+3)} [S_{-1} + \ln(2)] \end{aligned} \quad (48)$$

$$\begin{aligned}
H_{FB}^{(3),LL}(N) = & \frac{64(3N^2 + 3N - 1)P_2}{(N-1)N^2(N+1)^2(N+2)(2N-1)(2N+3)} S_1 \\
& - \frac{4P_7}{3(N-1)^2N^3(N+1)^3(N+2)^2(2N-3)(2N-1)^2(2N+3)^2(2N+5)} \\
& + (-1)^N \left\{ \frac{8P_4}{N(N+1)(2N-1)(2N+1)(2N+3)} S_{-2} \right. \\
& - \frac{256(4N^2 + 4N - 1)}{(2N-1)(2N+1)(2N+3)} \left[ S_{-1,1} - \frac{1}{2} \ln^2(2) + \sum_{i=1}^N \frac{\ln(2) + S_{-1}(i)}{1+2i} \right] \\
& - \frac{128P_5}{3(2N-3)(2N-1)^2(2N+1)^2(2N+3)^2(5+2N)} S_{-1} \\
& + \frac{4P_3}{N(N+1)(2N-1)(2N+1)(2N+3)} \zeta_2 \\
& \left. - \frac{128P_6}{3(2N-3)(2N-1)^2(2N+1)^2(2N+3)^2(2N+5)} \ln(2) \right\}, \tag{49}
\end{aligned}$$

with the polynomials

$$P_1 = 4N^4 + 8N^3 - N^2 - 5N - 3 \tag{50}$$

$$P_2 = 4N^4 + 8N^3 - N^2 - 5N - 3, \tag{51}$$

$$P_3 = 32N^4 + 16N^3 - 48N^2 - 14N + 9, \tag{52}$$

$$P_4 = 64N^4 + 80N^3 - 24N^2 - 22N + 9, \tag{53}$$

$$P_5 = 128N^7 + 256N^6 - 320N^5 - 144N^4 + 1128N^3 + 104N^2 - 936N + 27, \tag{54}$$

$$P_6 = 896N^7 + 2944N^6 - 512N^5 - 7344N^4 - 2760N^3 + 2816N^2 + 36N - 243, \tag{55}$$

$$\begin{aligned}
P_7 = & 5120N^{14} + 35840N^{13} + 73856N^{12} - 22784N^{11} - 252848N^{10} - 201200N^9 \\
& + 193136N^8 + 264800N^7 - 43255N^6 - 113445N^5 + 7512N^4 + 13587N^3 \\
& + 1539N^2 + 8262N - 1620. \tag{56}
\end{aligned}$$

Since the expressions for the higher order radiators are voluminous, they are given in a computer-readable file attached to this paper. Here the functions  $S_{\vec{a}} \equiv S_{\vec{a}}(N)$  denote the harmonic sums [36, 37]

$$S_{b,\vec{a}}(N) = \sum_{k=1}^N \frac{(\text{sign}(b))^k}{k^{|b|}} S_{\vec{a}}(k), \quad S_{\emptyset} = 1, \quad a, b_i \in \mathbb{Z} \setminus \{0\}. \tag{57}$$

Despite the fact that already at 2nd order in  $z$ -space cyclotomic harmonic polylogarithms occur, cyclotomic harmonic sums are only contributing from 3rd order onward. As usual, the  $N$ -space expressions turn out to be structurally simpler than the  $z$ -space expressions. However, there are evanescent poles at half integer arguments  $N = 1/2, 3/2, \dots$ . As has been seen in the small  $z$  expansion of the radiators, they are all tractable for  $N > 1$ . The pure singlet kernels are introducing a rightmost pole at  $N = 1$  and poles left to this are allowed.

One may numerically represent the higher order corrections also starting from Mellin  $N$  space, as described in Refs. [38, 45]. For this asymptotic representations and the recurrences of the respective expressions need to be known, which can be easily obtained using the package `HarmonicSums`. The recurrences are implied by the quantities itself in terms of the hierachic

sum-structures.

### Acknowledgments.

We would like to thank W. Hollik, P. Janot, P. Marquard, G. Passarino, and C. Schneider for discussions. This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 764850, SAGEX, COST action CA16201: Unraveling new physics at the LHC through the precision frontier.

## References

- [1] E. Accomando *et al.* Phys. Rept. **299** (1998) 1–78 [hep-ph/9705442];  
 J.A. Aguilar-Saavedra *et al.* hep-ph/0106315;  
 International Linear Collider Reference Design Report, ILC-REPORT-2007-001, Eds. J. Brau, Y. Okada, and N. Walker; Vol. 1–4.  
 G. Aarons *et al.* [ILC Collaboration], International Linear Collider Reference Design Report Volume 2: Physics At The ILC, [arXiv:0709.1893 [hep-ph]];  
<http://www.linearcollider.org/ILC>
- [2] H. Aihara *et al.* [Linear Collider Collaboration], *The International Linear Collider. A Global Project*, arXiv:1901.09829 [hep-ex].
- [3] J. Mnich, *The International Linear Collider: Prospects and Possible Timelines*, arXiv: 1901.10206 [hep-ex].
- [4] S. van der Meer, The CLIC Project and the Design for an  $e^+e^-$  Collider, CLIC-NOTE-68, (1988);  
 R.W. Assmann et al., CLIC Study Team, A 3 TeV  $e^+e^-$  Linear Collider Based on CLIC Technology, CERN 2000-008;  
 E. Accomando *et al.* [CLIC Physics Working Group], *Physics at the CLIC multi-TeV linear collider*, arXiv:hep-ph/0412251;  
 P. Roloff *et al.* [CLIC and CLICdp Collaborations], *The Compact Linear  $e^+e^-$  Collider (CLIC): Physics Potential*, arXiv:1812.07986 [hep-ex].
- [5] A. Abada *et al.* [FCC Collaboration], *Future Circular Collider : 2 The Lepton Collider (FCC-ee)*, CERN-ACC-2018-0057, Eur. Phys. J. Special Topics **228** (2019) no.2, 261–623;  
<http://tlep.web.cern.ch/>
- [6] J.P. Delahaye *et al.*, *Muon Colliders*, arXiv:1901.06150 [physics.acc-ph].
- [7] M.J.G. Veltman, Rev. Mod. Phys. **72** (2000) 341–349.
- [8] J. Blümlein, A. De Freitas, C.G. Raab and K. Schönwald, Phys. Lett. B **791** (2019) 206–209 [arXiv:1901.08018 [hep-ph]].
- [9] J. Blümlein, A. De Freitas, C.G. Raab and K. Schönwald, Phys. Lett. B **801** (2020) 135196 [arXiv:1910.05759 [hep-ph]].
- [10] J. Blümlein, A. De Freitas, C. Raab and K. Schönwald, Nucl. Phys. B **956** (2020) 115055 [arXiv:2003.14289 [hep-ph]].
- [11] J. Ablinger, J. Blümlein, A. De Freitas and K. Schönwald, Nucl. Phys. B **955** (2020) 115045 [arXiv:2004.04287 [hep-ph]].
- [12] J. Blümlein, A. De Freitas and W.L. van Neerven, Nucl. Phys. B **855** (2012) 508–569 [arXiv:1107.4638 [hep-ph]].
- [13] F.A. Berends, W.L. van Neerven and G.J.H. Burgers, Nucl. Phys. B **297** (1988) 429–478. Erratum: [Nucl. Phys. B **304** (1988) 921–922].

- [14] B.A. Kniehl, M. Krawczyk, J.H. Kühn and R.G. Stuart, Phys. Lett. B **209** (1988) 337–342.
- [15] G. Montagna, O. Nicrosini, F. Piccinini and G. Passarino, Comput. Phys. Commun. **117** (1999) 278–289 [arXiv:hep-ph/9804211 [hep-ph]].
- [16] D.Y. Bardin, M. Grünewald and G. Passarino, *Precision calculation project report*, arXiv:hep-ph/9902452 [hep-ph];  
D.Y. Bardin and G. Passarino, *The standard model in the making: Precision study of the electroweak interactions*, International series of monographs on physics **104** (Calendron Press, Oxford, 1999).
- [17] D.Y. Bardin, P. Christova, M. Jack, L. Kalinovskaya, A. Olchevski, S. Riemann and T. Riemann, Comput. Phys. Commun. **133** (2001) 229–395 [arXiv:hep-ph/9908433 [hep-ph]];  
A.B. Arbuzov, M. Awramik, M. Czakon, A. Freitas, M.W. Grünewald, K. Möning, S. Riemann and T. Riemann, Comput. Phys. Commun. **174** (2006) 728–758 [arXiv:hep-ph/0507146 [hep-ph]].
- [18] S. Schael *et al.* Phys. Rept. **427** (2006) 257–454 [hep-ex/0509008].
- [19] M. Böhm, W. Hollik, D.Y. Bardin, W. Beenakker, F.A. Berends, M.S. Bilenky, G. Burgers, J.E. Campagne, A. Djouadi, O. Fedorenko, *et al.*, *Forward-backward asymmetries*, CERN-TH-5536-89.
- [20] M. Greco and A. F. Grillo, Lett. Nuovo Cim. **15** (1976) 174–178;  
M. Greco, G. Panchari-Srivastava and Y. Srivastava, Nucl. Phys. B **171** (1980) 118–140 [Erratum: Nucl. Phys. B **197** (1982) 543–545];  
O. Nicrosini and L. Trentadue, Z. Phys. C **39** (1988) 479–486;  
J. E. Campagne and R. Zitoun, Phys. Lett. B **222** (1989) 497–500;  
O. Nicrosini and L. trentadue, *Structure functions techniques in  $e^+e^-$  collisions* in : Radiative corrections for  $e^+e^-$  Collisions, (Springer, Berlin, 1989), Ed. J.H. Kühn, 25–54;  
W. Beenakker, F.A. Berends and S.C. van der Marck, Phys. Lett. B **251** (1990) 299–304;  
S. Jadach, B.F.L. Ward and Z. Was, Phys. Lett. B **257** (1991) 213–218;  
S. Jadach and S. Yost, Phys. Rev. D **100** (2019) no.1, 013002 [arXiv:1801.08611 [hep-ph]].
- [21] D.Y. Bardin, M.S. Bilenky, A. Chizhov, A. Sazonov, Y. Sedykh, T. Riemann and M. Sachwitz, Phys. Lett. B **229** (1989) 405–408.
- [22] D.Y. Bardin, M.S. Bilenky, A. Sazonov, Y. Sedykh, T. Riemann and M. Sachwitz, Phys. Lett. B **255** (1991) 290–296 [arXiv:hep-ph/9801209 [hep-ph]].
- [23] W. Beenakker, F.A. Berends and W.L. van Neerven, *Applications of renormalization group methods to radiative corrections*, in : Radiative corrections for  $e^+e^-$  Collisions, (Springer, Berlin, 1989), Ed. J.H. Kühn, 3–24. Print-89-0445 (Leiden).
- [24] T. Riemann and Z. Was, Mod. Phys. Lett. A **4** (1989) 2487–2491;  
Z. Was and S. Jadach, Phys. Rev. D **41** (1990) 1425–1437;  
G. Montagna, O. Nicrosini and L. Trentadue, Phys. Lett. B **231** (1989) 492–496.
- [25] B.W. Lynn and R.G. Stuart, Nucl. Phys. B **253** (1985) 216–230;  
D.Y. Bardin, M. S. Bilenky, G. Mitselmakher, T. Riemann and M. Sachwitz, Z. Phys. C **44** (1989) 493–502.;  
W.J.P. Beenakker, *Electroweak corrections: techniques and applications*, PhD Thesis, (Leiden University, 1989);  
D.Y. Bardin, M.S. Bilenky, A. Chizhov, A. Sazonov, O. Fedorenko, T. Riemann and M. Sachwitz, Nucl. Phys. B **351** (1991) 1–48 [arXiv:hep-ph/9801208 [hep-ph]];  
A. Leike and T. Riemann, Z. Phys. C **51** (1991) 113–118;  
W. Beenakker, S.C. van der Marck and W. Hollik, Nucl. Phys. B **365** (1991) 24–78;  
W. Beenakker and W. Hollik, Phys. Lett. B **269** (1991) 425–431;  
W. Beenakker, A. Denner and A. Kraft, Nucl. Phys. B **410** (1993) 219–244;  
W. Beenakker and G. Passarino, Phys. Lett. B **425** (1998) 199–207 [arXiv:hep-ph/9710376 [hep-ph]];  
G. Duckeck and W. Hollik, *Electroweak precision tests at LEP*, Springer Tracts Mod. Phys. **162** (2000) 1–161;  
J.H. Kühn, A.A. Penin and V.A. Smirnov, Eur. Phys. J. C **17** (2000) 97–105 [arXiv:hep-ph/9912503 [hep-ph]];

- B. Feucht, J.H. Kühn, A.A. Penin and V.A. Smirnov, Phys. Rev. Lett. **93** (2004) 101802 [arXiv:hep-ph/0404082 [hep-ph]];  
 G. Bell, J.H. Kühn and J. Rittinger, Eur. Phys. J. C **70** (2010) 659–671 [arXiv:1004.4117 [hep-ph]];  
 A.V. Manohar and M. Trott, Phys. Lett. B **711** (2012) 313–316 [arXiv:1201.3926 [hep-ph]]; I. Dubovsky, A. Freitas, J. Gluza, T. Riemann and J. Usovitsch, JHEP **08** (2019) 113 [arXiv:1906.08815 [hep-ph]].
- [26] D.Y. Bardin, M.S. Bilenky, A. Chizhov, A. Sazonov, O. Fedorenko, T. Riemann and M. Sachwitz, Nucl. Phys. B **351** (1991) 1–48 [arXiv:hep-ph/9801208 [hep-ph]].
- [27] J. Jersak, E. Laermann and P.M. Zerwas, Phys. Lett. B **98** (1981) 363–366; Phys. Rev. D **25** (1982) 1218–1228 [Erratum: Phys. Rev. D **36** (1987) 310];  
 A. Djouadi, Z. Phys. C **39** (1988) 561–568;  
 A.B. Arbuzov, D.Y. Bardin and A. Leike, Mod. Phys. Lett. A **7** (1992) 2029–2038 [Erratum: Mod. Phys. Lett. A **9** (1994) 1515];  
 J.B. Stav and H.A. Olsen, Phys. Rev. D **52** (1995) 1359–1368;  
 G. Altarelli and B. Lampe, Nucl. Phys. B **391** (1993) 3–22;  
 P. Nason and B.R. Webber, Nucl. Phys. B **421** (1994) 473–517 [Erratum: Nucl. Phys. B **480** (1996) 755];  
 P.J. Rijken and W.L. van Neerven, Phys. Lett. B **386** (1996) 422–428 [arXiv:hep-ph/9604436 [hep-ph]];  
 Nucl. Phys. B **487** (1997) 233–282 [arXiv:hep-ph/9609377 [hep-ph]]; Phys. Lett. B **392** (1997) 207–215 [arXiv:hep-ph/9609379 [hep-ph]];  
 V. Ravindran and W.L. van Neerven, Phys. Lett. B **445** (1998) 214–222 [arXiv:hep-ph/9809411 [hep-ph]];  
 W. Bernreuther, R. Bonciani, T. Gehrmann, R. Heinesch, T. Leineweber, P. Mastrolia and E. Remiddi, Nucl. Phys. B **750** (2006) 83–107; [arXiv:hep-ph/0604031 [hep-ph]];  
 J.A. Aguilar-Saavedra, W. Bernreuther and Z.G. Si, Phys. Rev. D **86** (2012) 115020 [arXiv:1209.6352 [hep-ph]];  
 J. Gao and H.X. Zhu, Phys. Rev. Lett. **113** (2014) no.26, 262001 [arXiv:1410.3165 [hep-ph]];  
 W. Bernreuther, L. Chen, O. Dekkers, T. Gehrmann and D. Heisler, JHEP **01** (2017) 053 [arXiv:1611.07942 [hep-ph]];  
 D. d’Enterria and C. Yan, *Revised QCD effects on the  $Z \rightarrow b\bar{b}$  forward-backward asymmetry*, [arXiv:2011.00530 [hep-ph]];  
 A.B. Arbuzov, T.V. Kopylova and G.A. Seikhanova, Mod. Phys. Lett. A **35** (2020) no.25, 2050210 [arXiv:2003.14054 [hep-ph]].
- [28] M. Skrzypek, Acta Phys. Polon. **B23** (1992) 135–172;  
 M. Jezabek, Z. Phys. **C56** (1992) 285–288;  
 M. Przybycien, Acta Phys. Polon. B **24** (1993) 1105–1114 [hep-th/9511029];  
 J. Blümlein, S. Riemersma and A. Vogt, Eur. Phys. J. C **1** (1998) 255–259 [hep-ph/9611214];  
 A.B. Arbuzov, Phys. Lett. B **470** (1999) 252–258 [hep-ph/9908361];  
 A.B. Arbuzov, JHEP **0107** (2001) 043 [hep-ph/9907500];  
 J. Blümlein and H. Kawamura, Nucl. Phys. B **708** (2005) 467–510 [hep-ph/0409289]; Eur. Phys. J. C **51** (2007) 317–333 [arXiv:hep-ph/0701019].
- [29] R. Hamberg, W.L. van Neerven and T. Matsuura, Nucl. Phys. B **359** (1991) 343–405 Erratum: [Nucl. Phys. B **644** (2002) 403–404].
- [30] R.V. Harlander and W.B. Kilgore, Phys. Rev. Lett. **88** (2002) 201801 [hep-ph/0201206].
- [31] E. Remiddi and J.A.M. Vermaasen, Int. J. Mod. Phys. **A15** (2000) 725–754 [hep-ph/9905237].
- [32] J. Ablinger, J. Blümlein and C. Schneider, J. Math. Phys. **52** (2011) 102301 [arXiv:1105.6063 [math-ph]].
- [33] C. Schneider, Sémin. Lothar. Combin. **56** (2007) 1–36 article B56b.
- [34] C. Schneider, Simplifying Multiple Sums in Difference Fields, in: *Computer Algebra in Quantum Field Theory: Integration, Summation and Special Functions* Texts and Monographs in Symbolic Computation eds. C. Schneider and J. Blümlein (Springer, Wien, 2013) 325–360 [arXiv:1304.4134 [cs.SC]].
- [35] J. Ablinger, J. Blümlein and C. Schneider, J. Math. Phys. **54** (2013) 082301 [arXiv:1302.0378 [math-ph]];  
 J. Ablinger, J. Blümlein, C.G. Raab and C. Schneider, J. Math. Phys. **55** (2014) 112301 [arXiv:1407.1822 [hep-th]];

J. Blümlein, Comput. Phys. Commun. **159** (2004) 19–54 [arXiv:hep-ph/0311046 [hep-ph]];  
J. Ablinger, J. Blümlein and C. Schneider, J. Phys. Conf. Ser. **523** (2014) 012060 [arXiv:1310.5645 [math-ph]];  
J. Ablinger, PoS (LL2014) 019 [arXiv:1407.6180[cs.SC]]; *A Computer Algebra Toolbox for Harmonic Sums Related to Particle Physics*, Diploma Thesis, JKU Linz, 2009, arXiv:1011.1176[math-ph]; *Computer Algebra Algorithms for Special Functions in Particle Physics*, Ph.D. Thesis, Linz U. (2012) arXiv:1305.0687[math-ph]; PoS (LL2016) 067; Experimental Mathematics 26 (2017) [arXiv:1507.01703 [math.CO]]; PoS (RAD-COR2017) 001 [arXiv:1801.01039 [cs.SC]]; arXiv:1902.11001 [math.CO]; PoS (LL2018) 063.

- [36] J. Vermaseren, Int. J. Mod. Phys. **A14** (1999) 2037–2076 [hep-ph/9806280].
- [37] J. Blümlein and S. Kurth, Phys. Rev. **D60** (1999) 014018 [hep-ph/9810241].
- [38] J. Blümlein, Comput. Phys. Commun. **180** (2009) 2218–2249 [arXiv:0901.3106 [hep-ph]].
- [39] M. Böhm, A. Denner, and H. Joos, *Gauge Theories of the Strong and Electroweak Interaction*, (B.G. Teubner, Stuttgart, 2001);  
A. Denner and S. Dittmaier, *Electroweak Radiative Corrections for Collider Physics*, arXiv:1912.06823 [hep-ph].
- [40] P. Janot, JHEP **02** (2016) 053 [Erratum: JHEP **11** (2017) 164] [arXiv:1512.05544 [hep-ph]].
- [41] P.A. Zyla et al. (Particle Data Group), Prog. Theor. Exp. Phys. (2020) 083C01.
- [42] T. Gehrmann and E. Remiddi, Comput. Phys. Commun. **141** (2001) 296–312 [hep-ph/0107173].
- [43] J. Ablinger, J. Blümlein, M. Round and C. Schneider, Comput. Phys. Commun. **240** (2019) 189–201 [arXiv:1809.07084 [hep-ph]].
- [44] J. Blümlein and H. Kawamura, Phys. Lett. B **553** (2003) 242–250 [hep-ph/0211191];  
A.B. Arbuzov, Phys. Part. Nucl. **50** (2019) no.6, 721–825.
- [45] J. Blümlein, Comput. Phys. Commun. **133** (2000) 76–104 [hep-ph/0003100];  
J. Blümlein and S.O. Moch, Phys. Lett. B **614** (2005) 53–61 [hep-ph/0503188].