


DEUTSCHES ELEKTRONEN-SYNCHROTRON
Ein Forschungszentrum der Helmholtz-Gemeinschaft



DESY 21-030
TTK-21-07
arXiv:2103.06773
March 2021

Expectation Management

J. Braathen

Deutsches Elektronen-Synchrotron DESY, Hamburg

M. D. Goodsell

*Laboratoire de Physique Théorique et Hautes Energies (LPTHE),
Sorbonne Université et CNRS, Paris, France*

S. Paßehr

Institut für Theoretische Teilchenphysik und Kosmologie, RWTH Aachen

E. Pinsard

*Laboratoire de Physique de Clermont, CNRS/IN2P3,
Université Clermont Auvergne, Aubière, France*

ISSN 0418-9833

NOTKESTRASSE 85 - 22607 HAMBURG

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your reports and preprints are promptly included in the
HEP literature database
send them to (if possible by air mail):

| | |
|---|---|
| DESY Zentralbibliothek Notkestraße 85 22607 Hamburg Germany | DESY Bibliothek Platanenallee 6 15738 Zeuthen Germany |
|---|---|

Expectation management

Johannes Braathen^{a *}, Mark D. Goodsell^{b †}, Sebastian Paßehr^{c ‡} and
Emanuelle Pinsard^{d §}

^a *Deutsches Elektronen-Synchrotron DESY, Notkestraße 85, D-22607 Hamburg, Germany.*

^b *Laboratoire de Physique Théorique et Hautes Energies (LPTHE),
UMR 7589, Sorbonne Université et CNRS, 4 place Jussieu, 75252 Paris Cedex 05, France.*

^c *Institute for Theoretical Particle Physics and Cosmology,
RWTH Aachen University, Sommerfeldstraße 16, 52074 Aachen, Germany.*

^d *Laboratoire de Physique de Clermont (UMR 6533), CNRS/IN2P3,
Univ. Clermont Auvergne, 4 Av. Blaise Pascal, F-63178 Aubière Cedex, France*

We consider the application of a Fleischer–Jegerlehner-like treatment of tadpoles to the calculation of neutral scalar masses (including the Higgs) in general theories beyond the Standard Model. This is especially useful when the theory contains new scalars associated with a small expectation value, but comes with its own disadvantages. We show that these can be overcome by combining with effective field theory matching. We provide the formalism in this modified approach for matching the quartic coupling of the Higgs via pole masses at one loop, and apply it to both a toy model and to the μ NMSSM as prototypes where the standard treatment can break down.

arXiv:2103.06773v2 [hep-ph] 5 Aug 2021

* e-mail: johannes.braathen@desy.de
† e-mail: goodsell@lpthe.jussieu.fr
‡ e-mail: passehr@physik.rwth-aachen.de
§ e-mail: emanuelle.pinsard@clermont.in2p3.fr

1 Introduction

The mass of the SM-like Higgs boson, discovered by ATLAS and CMS [1–3], is now an electroweak precision observable, thanks to its outstandingly accurate determination at the LHC [4–6], and it plays an important role in constraining the allowed parameter space of Beyond-the-Standard-Model (BSM) theories. On the one hand, the Higgs mass is a prediction in supersymmetric theories (see Ref. [7] and references therein for a recent review) and interestingly it depends most heavily on the electroweak couplings and scale – quantities that are already known from other observations – while it is only at loop level that a dependence on the scale of supersymmetric particles appears. This property has spurred significant developments in precision scalar-mass calculations, advanced in recent years by the KUTS initiative [8–54] as described in the report [7]. On the other hand, in non-supersymmetric theories, the Higgs mass is not a prediction by itself, but it can be used to extract the Higgs quartic coupling and, in turn, investigate the stability of the electroweak vacuum. In this context, a precise calculation is essential to produce reliable results on vacuum stability (see Refs. [55–59] for works in the SM) and to correctly appreciate the potential impact of new particles [60–65].

We refer the interested reader to Ref. [7] and references therein for an in-depth review of Higgs-mass computations, and we only recall here the main steps involved (applicable for any BSM theory). The standard calculational technique begins with the extraction of SM-like parameters – namely the electroweak and strong gauge couplings, the quark and lepton Yukawa couplings, and the Higgs vacuum expectation value (vev) – from observables. Adding then the BSM parameters to these, the Higgs (and other particle) masses can be calculated, along with any other desired predictions. The relevant observables for the electroweak sector are typically, as in calculations in the SM, either $M_Z, M_W, \alpha(0)$ or $M_Z, G_F, \alpha(0)$ where $M_{Z,W}$ are the Z and W boson masses, $\alpha(0)$ is the fine-structure constant extracted in the Thompson limit, and G_F is the Fermi constant. This latter quantity is extracted from muon three-body decays, whereas the others are related essentially to self-energies. In general, this extraction of the SM-like couplings and the Higgs vev can be performed at one-loop for any theory, but the two-loop relationships are only known for the SM and a small subset of other models in certain limits.

At the tree level, the expectation value v of the Higgs boson is related to the other parameters in the theory by the requirement that the theory be at the minimum of the potential. To be concrete, consider the Higgs potential of the SM, $V = \mu^2 |H|^2 + \lambda |H|^4$; then the minimisation condition gives

$$0 = \mu^2 + \lambda v^2. \quad (1.1)$$

Since we do not have an observable for μ^2 we typically use this equation to eliminate it, giving the Higgs mass to be

$$m_h^2 = \mu^2 + 3\lambda v^2 = 2\lambda v^2. \quad (1.2)$$

However, once we go beyond tree level, there are several possible choices. The approach typically taken in BSM theories, and in the SM in Ref. [66], is to insist that the expectation value v is a fixed “observable”, and instead keep solving for μ^2 order-by-order in perturbation theory. In this way,

$$\mu^2 = -\lambda v^2 - \frac{1}{v} \left. \frac{\partial \Delta V}{\partial h} \right|_{h=0} \equiv -\lambda v^2 - \frac{1}{v} t_h, \quad (1.3)$$

where ΔV are the loop corrections to the effective potential, and then the Higgs pole mass M_h reads

$$M_h^2 = 2\lambda v^2 - \frac{1}{v} t_h + \Pi_{hh}(M_h^2) \equiv 2\lambda v^2 + \Delta M_h^2, \quad (1.4)$$

where $\Pi_{hh}(M_h^2)$ is the Higgs self-energy evaluated on-shell. One of the chief advantages of this approach is that tadpole diagrams do not appear in any processes, since they vanish by construction.

On the other hand, while this is in principle a straightforward procedure to follow, it is complicated by the fact that the self-energies and effective potential implicitly depend on μ^2 . In Landau gauge, or the gaugeless limit, this leads to the “Goldstone Boson Catastrophe” at two loops [67–70] – its solution appears by consistently solving the above equation order by order [27, 33]. Indeed, one way to formalise this is as a finite (or possibly IR-divergent) counterterm for μ^2 :

$$\mathcal{L} \supset -(\mu^2 + \delta\mu^2 + \lambda v^2) v h - \frac{1}{2} (\mu^2 + \delta\mu^2 + 3\lambda v^2) h^2 + \dots, \quad (1.5)$$

where $\delta\mu^2 = -\frac{1}{v} t_h$. Another drawback is that it manifestly breaks gauge invariance, since the loop corrections above depend on the gauge; and it also means that the expectation value v is not an $\overline{\text{MS}}$ parameter, so the renormalisation-group equations for the expectation value are no longer just given by those of μ^2 and λ , but have extra contributions [71, 72].

However, there is a further drawback to the above procedure which we wish to highlight in this paper. When considering a BSM theory with additional scalars that may have an expectation value, it is typical to take the same approach as for the scalar field in the SM and fix their expectation values, solving the additional tadpole equations for other dimensionful parameters – for example, their mass-squared parameters, or sometimes a cubic scalar coupling. To take the example of a real singlet S with mass-squared *Lagrangian parameter* m_S^2 – not to be confused with the pole mass, which we denote M_S – and expectation value v_S , this means that analogously to eq. (1.3),

$$m_S^2 = (m_S^2)^{\text{tree}} - \frac{1}{v_S} \frac{\partial \Delta V}{\partial S}. \quad (1.6)$$

If the loop corrections are not large, and v_S is not small, this is completely acceptable – so for models such as the NMSSM there is generally no problem. However, if we consider a different theory or regions of the parameter space where v_S is small, for example if $m_S \gg v$ and $v_S \propto v^2$ (as may be found in examples of EFT matching [41]) then we can easily find the case that $\delta m_S^2 > (m_S^2)^{\text{tree}}$. This makes the calculation unreliable.

The archetypal example of this problem is the case where the neutral scalar obtaining an expectation value actually comes from an $SU(2)$ triplet \mathbf{T} with expectation value v_T and mass-squared m_T^2 – for example in Dirac-gaugino models [73–76]. In that case, $v_T \propto v^2/m_T^2$ multiplied by other dimensionful parameters of the theory. Moreover, we require that $v_T \lesssim 4 \text{ GeV}$ from electroweak-precision constraints, generally requiring $m_T \gtrsim 1 \text{ TeV}$. So then

$$\delta m_T^2 \sim \frac{1}{4 \text{ GeV}} \times \frac{1}{16\pi^2} \times \mathcal{O}(\text{TeV}^3) \sim 2.5 \times \mathcal{O}(\text{TeV}^2), \quad (1.7)$$

i. e. we see that there is a severe problem whenever v_T/m_T is of the order of a loop factor.

Moreover, for such cases where v_S is small, this procedure works in the opposite way to that which we would desire. In BSM theories the scalar expectation values beyond v are not top-down inputs or tied closely to some observables, whereas we may typically want to define the masses and couplings as fixed by some high-energy boundary conditions (for example constrained or minimal SUGRA conditions where soft masses have a common origin). In this case we would like to solve the tadpole equations for v_S ; even if this would typically lead to coupled cubic equations, nowadays it is almost trivial to solve them numerically, or start from an approximation.

In this paper we will instead examine an alternative procedure, proposed by Fleischer and Jegerlehner in examining Higgs decays in the SM [77], which has the potential to solve both of these issues. Instead of taking the expectation values as fixed, we take them to be the tree-level solutions of the tadpole equations. This means that we do not work at the “true” minimum of the potential and must include tadpole diagrams in all processes. While this implies the addition of some new Feynman diagrams in the Higgs mass calculation, it is not technically more complicated than including finite counterterm insertions for μ^2 . This approach has the additional advantages that, since the Lagrangian is specified in

terms of $\overline{\text{MS}}$ parameters only, the result is manifestly gauge independent, and the expectation values are just the solutions to the tree-level tadpole equations. For these reasons, it has been used and advocated in the SM, in particular at two loops in Ref. [57, 78–82]; and applied to certain extensions of the Two Higgs Doublet Model (THDM) when considering decays [83–86]. We also note that this approach is closely related to the various on-shell renormalisations used in *e.g.* Refs. [87–90] in the THDM and the Minimal Supersymmetric Standard Model (MSSM).

In the example of the SM at the one-loop order, this would mean

$$M_h^2 = 2\lambda v^2 - \frac{6\lambda v}{m_h^2} t_h^{(1)} + \Pi_{hh}^{(1)}(m_h^2), \quad (1.8)$$

where the superscripts in brackets indicate the loop order, and we put the momentum in the self-energy at the tree-level Higgs mass in order to respect the order of perturbation theory. In other words, the tadpole contribution is suppressed by the mass-squared of the Higgs, although – since $m_h^2 = 2\lambda v^2$ – here we find that they have a very similar form to the previous approach. On the other hand, in the case of a heavy singlet or triplet the contributions to the singlet self-energy would be similarly suppressed by m_S^2 , and we can have m_S much greater than the triplet coupling – so the corrections to the singlet mass would be well under control.

On the other hand, in the BSM context this approach was proposed by Ref. [91] for the following very different reason: by no-longer forcing the electroweak expectation value to have its observed value, we allow new physics to disturb the electroweak hierarchy. In the above approach, the contribution $-\frac{6\lambda v}{m_h^2} t_h^{(1)} = -\frac{3}{v} t_h^{(1)}$ is effectively the contribution *from a shift in v* . We can view the calculation as equivalent to counterterms for the expectation value $\delta^{(1)}v$, where

$$\mathcal{L} \supset -(\mu^2 + \lambda v^2) v h - (\mu^2 + 3\lambda v^2) \delta^{(1)}v h - \dots \quad (1.9)$$

so that now

$$\delta^{(1)}v = -\frac{1}{m_h^2} t_h^{(1)}. \quad (1.10)$$

In this case, if there is heavy new physics at a scale $\Lambda \gg m_h$, then we shift the Higgs expectation value up to that new scale suppressed only by a loop factor. Indeed in Ref. [91] the proposal was to use

$$\frac{\delta m_h^2}{m_h^2} \equiv \frac{1}{m_h^2} \left[-\frac{3}{v} t_h^{(1)} + \Pi_{hh}^{(1)}(m_h^2) \right] \quad (1.11)$$

as *a measure of fine-tuning of the theory*.

Another perspective on the difference between the two approaches is given by viewing the SM as an EFT. In this case, in the EFT the SM receives corrections to both μ^2 and λ at the matching scale from integrating out heavy states which can be done with $v = 0$. As discussed in Ref. [27], when expanding in v , in order to respect gauge invariance we must have:

$$\begin{aligned} \Delta V &= \Delta V_0 + \frac{1}{2} \Delta V_{hh}|_{v=0} v^2 + \mathcal{O}(v^4) + \dots, \\ \Pi_{hh}(m_h^2) &= \Delta V_{hh}|_{v=0} + \mathcal{O}(v^2) \end{aligned} \quad (1.12)$$

and therefore $t_h = v \Delta V_{hh}|_{v=0} + \dots$. This shows that the EFT-matching correction to μ^2 , which is $\Delta V_{hh}|_{v=0}$, and the origin of the hierarchy problem, correspond to t_h/v to lowest order in v . Hence in the “standard” approach of eq. (1.4) this cancels out and leaves only corrections proportional to v^2 – whereas in the modified approach it remains and gives a large shift to the Higgs mass.

However, the reappearance of the hierarchy is a problem for the *light* Higgs mass, whereas the problem we wished to solve actually appeared in new, *heavy* states! If we wish to explore theories which may remain natural while having heavy states, such as those in Ref. [91], then the modified tadpole approach should work best. There must consequently be some trade-off between losing control of the light Higgs and losing control of the heavier states (and losing gauge invariance too). In section 2 we will set up the necessary general formalism and explore this in detail for a toy model.

However, there are *two* potential solutions to allow us to have the best of both worlds:

1. Retain counterterms for μ^2 as in eq. (1.5) for the SM Higgs, but *only* for them. This is somewhat tricky to automate, since we must make a special case of the electroweak sector, and we also lose gauge invariance.
2. For cases where the tuning of the hierarchy becomes large, use EFT pole matching [26] with the modified treatment of tadpoles. This way, the heavy states remain entirely under control, we keep the heavy masses and couplings as top-down inputs (that remain genuinely $\overline{\text{MS}}$ or $\overline{\text{DR}}'$), and we have gauge invariance built-in.

In section 4 we will adopt the second approach for the example of the general NMSSM (and apply it specifically to the variant known as the μ NMSSM [63]). We establish the necessary formalism for the matching and give a detailed examination, via implementing the computation in a modified SPheno [92, 93] code generated from SARAH [13, 16, 33, 94–98].

2 Treatment of tadpoles for theories with heavy scalars

For a general renormalisable field theory, once we have solved the vacuum minimisation conditions and diagonalised the mass matrices, we can write the potential in terms of real scalar fields $\{\phi_i\}$ as

$$V = \text{const} + \frac{1}{2} m_i^2 \phi_i^2 + \frac{1}{6} a_{ijk} \phi_i \phi_j \phi_k + \frac{1}{24} \lambda_{ijkl} \phi_i \phi_j \phi_k \phi_l. \quad (2.1)$$

If we take the standard approach and fix the expectation values, adjusting the mass parameters order by order in perturbation theory, then as described in Ref. [27] we can write the pole masses as

$$(M_i^2)^{(1)} = m_i^2 + \Delta_{ii} + \Pi_{ii}^{(1)}(m_i^2) \equiv m_i^2 + \Delta M_i^2. \quad (2.2)$$

To define the shifts Δ_{ii} in a general way, we must start from some basis of fields $\{\phi_i^0\}$ split into expectation values and fluctuations so that $\phi_i^0 \equiv v_i + \hat{\phi}_i^0$ and then diagonalise the fields via $\hat{\phi}_i^0 = R_{ij} \phi_j$. In the simplest case where we solve the tadpole equations for some mass-squared parameters in the original basis and where we ignore pseudoscalars, we can then write

$$\Delta_{ii} = - \sum_k R_{ki}^2 \frac{1}{v_k} \left. \frac{\partial \Delta V}{\partial \hat{\phi}_k^0} \right|_{\hat{\phi}_k^0=0} = - \sum_{k,l} R_{ki}^2 R_{lk} \frac{1}{v_k} t_l^{(1)}. \quad (2.3)$$

The generalisation to solving for other variables (such as cubic scalar couplings) and to include pseudoscalar mass shifts is given in Ref. [33].

On the other hand, taking the modified approach and including the tadpole diagrams, the pole masses up to one loop are simply

$$(M_i^2)^{(1)} = \hat{m}_i^2 - \frac{1}{\hat{m}_j^2} a_{ijj} t_j^{(1)} + \Pi_{ii}^{(1)}(\hat{m}_i^2) \equiv \hat{m}_i^2 + \hat{\Pi}_{ii}^{(1)}(\hat{m}_i^2), \quad (2.4)$$

where we have defined \hat{m}_i^2 to be the tree-level mass when we are using the modified scheme (we will later drop the distinction between m_i and \hat{m}_i , see below) and $\hat{\Pi}_{ij}(p^2)$ for later use to be the self-energies including the tadpoles. The expressions for the tadpoles and self-energies at one loop can be

found *e. g.* in Refs. [27, 99]; this calculation is therefore more straightforward to automate, being purely diagrammatic in nature. An *explicitly* gauge-invariant expression for this (*i. e.* one where there are no gauge-fixing parameters present) will be given in future work.

At this point the reader may object that, no matter what technique we use to calculate masses, the result for a given theory should be the same up to higher-loop corrections. Unfortunately this is made obscure by the difficulties in general in defining the parameters of our theory. To compare the two calculations *for the same parameter point*, in the standard approach we are invited to treat the expectation values as fundamental, so if we start from a theory defined in this way, we must:

1. Calculate loop-level masses in the standard approach for a given choice of expectation values (with the associated problems when those expectation values are small).
2. Extract the Lagrangian parameters from the loop-corrected tadpole equations.
3. Solve the tree-level vacuum stability equations with these new parameters, obtaining the expectation values for use in the alternative approach.
4. Compute the new tree-level spectrum using these expectation values
5. Compute the loop-corrected masses in the alternative approach.

Let us denote the tree-level masses and expectation values in the alternative approach as \hat{m}_i and \hat{v}_i , and for simplicity assume that we solve the tadpole equations for some mass-squared parameters (rather than cubic couplings, say). Then, by passing back to the basis in which the fields are not diagonalised, where the Lagrangian mass parameters are $m_{0,ij}^2 = \hat{m}_{0,ij}^2 + \delta m_{0,ij}^2$ and the Lagrangian couplings are $a_{0,ijk}$, $\lambda_{0,ijkl}$, we can carry out the above steps and solve perturbatively for the expectation values \hat{v}_i in the modified scheme:

$$\begin{aligned} 0 &= (m_{0,ij}^2 + \delta m_{0,ij}^2) v_j + \frac{1}{2} a_{0,ijk} v_j v_k + \frac{1}{6} \lambda_{0,ijkl} v_j v_k v_l + t_{0,i} \\ &= (m_{0,ij}^2 + \delta m_{0,ij}^2) \hat{v}_j + \frac{1}{2} a_{0,ijk} \hat{v}_j \hat{v}_k + \frac{1}{6} \lambda_{0,ijkl} \hat{v}_j \hat{v}_k \hat{v}_l. \end{aligned} \quad (2.5)$$

We have written $t_{0,i}$ for the one-loop tadpole to emphasise that it is in the undiagonalised basis; to go to the mass-diagonal basis we need to rotate by the matrix R_{ij} as above. Writing $\hat{v}_i = v_i + \delta v_i$ we obtain

$$0 = -t_{0,i} + \mathcal{M}_{0,ij}^2 \delta v_j, \quad (2.6)$$

where $\mathcal{M}_{0,ij}^2$ is the tree-level mass matrix of scalars in the standard scheme. This can be trivially solved by rotating to the mass-diagonal basis. We then write the tree-level mass matrix in the alternative scheme as

$$\hat{\mathcal{M}}_{0,ij}^2 = \mathcal{M}_{0,ij}^2 + \delta m_{0,ij}^2 + a_{0,ijk} \delta v_k + \lambda_{0,ijkl} v_k \delta v_l. \quad (2.7)$$

Using the *same* matrix R_{ij} we can rotate this to obtain*

$$\hat{m}_i^2 = (R^T \hat{\mathcal{M}}_0^2 R)_{ii} = m_i^2 + \Delta_{ii} + a_{iik} \frac{t_k^{(1)}}{m_k^2} + \mathcal{O}(2\text{-loop}). \quad (2.8)$$

Inserting this into (2.4) gives (2.2).

Of course, this comes with the associated problems of defining the theory in the standard approach: if we have a small expectation value, then (as we shall illustrate below) the loop corrections in Δ_{ii} can be very large, so the mass of the heavy scalar may differ greatly from the tree-level one. Making a conversion in this way just ensures that we see the same problem in the alternative treatment. Instead, for such points we should start with a theory defined in the *alternative* manner.

* Recall that $a_{ijk} = (a_{0,i'j'k'} + \lambda_{0,i'j'k'l'} v_{l'}) R_{i'i} R_{j'j} R_{k'k}$

Then, to compare the same point for the standard calculation one should:

1. Calculate loop-level masses in the alternative approach for a given choice of masses and couplings.
2. Iteratively solve the loop-level vacuum stability equations to obtain the loop-corrected expectation values v_i for use in the standard scheme.
3. Use these expectation values to compute the tree-level spectrum for use in the standard scheme (if we are using the approach with “consistent tadpoles”)[†]
4. Compute the loop-corrected masses in the standard approach.

In this way, we should obtain the same result (up to higher-order differences) for our desired point as in the alternative scheme. However, the key complicating factor is step 2: it assumes that we can efficiently and accurately find the true minimum of the potential. This can only be done by iteration of the tadpole equations; this involves *recomputing the masses and couplings of the theory at each step* and is therefore often numerically expensive (especially at higher loop orders). On the other hand, if we do this perturbatively, then we are effectively using the alternative scheme!

Disclaimer

While the above discussion is reassuring for the consistency of our calculations, in the following we will *not* (for the most part) compare masses at the same parameter point, for the obvious reason that the results would be almost the same. Instead, what we want to illustrate is the difficulty in even defining our theory: in the standard approach, since we are required to choose a vacuum-expectation value for the heavy singlet fields (which are not physical parameters), the phenomenologist will often use a guess or a tree-level-approximate solution for this, rather than iteratively solve the tadpole equations (which, in any case, would lead to a different input value depending on the chosen loop order). We shall take this naive approach below, and compare (in most cases) theories *with the same tree-level spectrum* by taking the expectation values to be the same in both the standard and modified schemes. Of course, according to the discussion above, these are not the same parameter points: we are instead illustrating the differences in methods of defining the theory, and will show how the alternative scheme gives a much more stable and efficient definition (at least in cases where the hierarchy problem for the light Higgs does not become severe).

2.1 A toy model

Let us now apply the above general expressions to the simplest toy model that can illustrate the differences of prescriptions for dealing with radiative corrections to tadpoles. This consists of the abelian Goldstone model coupled to a real singlet S , and has scalar potential

$$V = \mu^2 |H|^2 + \frac{1}{4} \lambda |H|^4 + \frac{1}{2} m_S^2 S^2 + a_{SH} S |H|^2 + \lambda_{SH} S^2 |H|^2 + a_S S^3 + \lambda_S S^4 \quad (2.9)$$

with the fields

$$H \equiv \frac{1}{\sqrt{2}} (v + h + iG), \quad S \equiv v_S + \hat{S}, \quad (2.10)$$

v and v_S denoting the Higgs and singlet vacuum expectation values (vevs), respectively. The minimisation conditions at the tree level yield the equations

$$-\mu^2 = \frac{1}{4} \lambda v^2 + a_{SH} v_S + \lambda_{SH} v_S^2, \quad (2.11a)$$

[†] In principle it is possible, and simpler, to just use the “true” input masses in the standard approach. This would vitiate the problem to a large extent, but would then lead to the well-known infra-red issues at two loops, or uncanceled logarithms in EFT matching, etc.




Figure 1: *left:* one-loop tadpole diagrams; *middle:* one-loop self-energy diagrams appearing in standard and modified calculation; *right:* additional self-energy diagrams in the modified approach.

$$(m_S^2 + \lambda_{SH} v^2) v_S = -\frac{1}{2} a_{SH} v^2 - 3 a_S v_S^2 - 4 \lambda_S v_S^3 \quad (2.11b)$$

that lead to the tree-level (squared) mass matrix for the scalars (which do not mix with the massless pseudoscalar):

$$\mathcal{M}_{\text{tree}}^2 = \begin{pmatrix} \frac{1}{2} \lambda v^2 & a_{SH} v + 2 \lambda_{SH} v v_S \\ a_{SH} v + 2 \lambda_{SH} v v_S & m_S^2 + \lambda_{SH} v^2 + 6 a_S v_S + 12 \lambda_S v_S^2 \end{pmatrix}. \quad (2.12)$$

The one-particle irreducible one-loop contributions to the one- and two-point functions (see figure 1) of this toy model are given by

$$t_i^{(1)} = -\frac{\kappa}{2} a_{ijj} A(m_j^2), \quad (2.13a)$$

$$\Pi_{ij}^{(1)}(p^2) = \kappa \left[\frac{1}{2} \lambda_{ijkk} A(m_k^2) - \frac{1}{2} a_{ikl} a_{jkl} B(p^2, m_k^2, m_l^2) \right] \quad (2.13b)$$

with A and B denoting the scalar one-point and two-point one-loop integrals in the conventions of *e. g.* Refs. [27, 99], $\kappa \equiv (16\pi^2)^{-1}$ and p^2 denoting the external momentum. In the approach of keeping the vevs fixed, we find for the one-loop pole masses:

$$(M_i^2)^{(1)} = m_i^2 - R_{i1}^2 \frac{1}{v} t_h^{(1)} - R_{i2}^2 \frac{1}{v_S} t_S^{(1)} + \Pi_{ii}(m_i^2), \quad (2.14)$$

where $t_h^{(1)} = \partial\Delta V/\partial h|_{h,\hat{S}=0}$, $t_S^{(1)} = \partial\Delta V/\partial S|_{h,\hat{S}=0}$. Thus the tadpole corrections suffer from the division by the vev; in particular, the mass predictions can become numerically unstable in scenarios with a small singlet vev. Let us see this in practice for our example when m_S^2 is large; in this case

$$v_S \sim -\frac{a_{SH} v^2}{2 m_S^2}, \quad R \sim \begin{pmatrix} 1 & -\frac{a_{SH} v}{m_S^2} \\ \frac{a_{SH} v}{m_S^2} & 1 \end{pmatrix}. \quad (2.15)$$

If we take v small and just look at the singlet mass in the limit $p^2 \rightarrow 0$ for simplicity,[‡] we have

$$\Delta M_S^2 \approx \Pi_{SS}(0) - \frac{1}{v_S} t_S \supset -\frac{3 a_S m_S^2 \kappa}{v_S} (\overline{\log} m_S^2 - 1) + \dots \quad (2.16)$$

where $\overline{\log} m_S^2 \equiv \log m_S^2/Q^2$ for renormalisation scale Q . When the system is really decoupled and $v = 0$, then $v_S \sim m_S^2/(6a_S)$ and this expression remains well-controlled, but when $0 < v \ll m_S$ – which is the case we are interested in – we instead have

$$\Delta M_S^2 \propto \frac{6 a_S m_S^4}{16 \pi^2 a_{SH} v^2} \overline{\log} m_S^2 \quad (2.17)$$

which can be very large compared to m_S^2 .

[‡] This limit is not implemented in our code and serves only the more lucid presentation. In fact, an off-shell evaluation of the self-energies implies unphysical behaviour of Higgs-mass predictions [100].

If we take the modified approach to tadpoles, then the relevant generic expression for the self-energy is

$$\hat{\Pi}_{ij}^{(1)}(p^2) = \frac{1}{16\pi^2} \left[\frac{1}{2} \lambda_{ijkk} A(m_k^2) - \frac{1}{2} a_{ikl} a_{jkl} B(p^2, m_k^2, m_l^2) - \frac{1}{2m_k^2} a_{ijk} a_{kll} A(m_l^2) \right]; \quad (2.18)$$

and for our example

$$\hat{\Pi}_{SS}^{(1)}(m_S^2) \approx \Pi_{SS}(0) - \frac{a_{SH}^2 \kappa}{2m_h^2} A(m_S^2) - \frac{3a_S^2 \kappa}{m_S^2} A(m_S^2) + \dots \sim -\frac{\kappa}{2} \left(\frac{a_{SH}^2}{m_h^2} - 24\lambda_S \right) m_S^2 \overline{\log} m_S^2. \quad (2.19)$$

Provided that $a_{SH} \lesssim m_h$ this is well under control, in contrast to the previous “standard” approach.

2.2 Numerical examples

In this section we shall illustrate the different behaviours of the two approaches to tadpoles in the toy model defined in eq. (2.9) through numerical examples. For this purpose, we present results for the one-loop pole masses M_h and M_S computed diagrammatically both in the standard approach – following eq. (2.2) – and in the modified approach of equation (2.4). We shall consider points defined to have *the same tree-level spectrum*, but whose loop-corrected masses differ according to the scheme used. As described in the disclaimer above, these are not therefore the same points in parameter space: this illustrates *the difficulty in defining the model*.

For all the following figures, we set $\lambda = 0.52$, to reproduce a light “Higgs” (noting that there are no gauge fields) near 125 GeV, and we also fix $\lambda_{SH} = 0$ and $\lambda_S = 1/24$. In each case, we shall fix the $\overline{\text{MS}}$ parameter m_S and solve the tree-level tadpole equations numerically to obtain v_S and fix $v = 246\text{GeV}$. Then the calculation in the modified scheme gives the correct value for the scalar masses. For comparison, in each of the figures 2, 3, 4 and 5 we use these same values as inputs for the conventional scheme, where we treat the derived value for v_S as the “all orders” expectation value; this means that, in the standard scheme, $(m_S^2)^{\text{mod.}} = (m_S^2)^{\text{tree}}$, the tree-level value, and is not actually the $\overline{\text{MS}}$ mass-squared parameter any more. Hence, as mentioned above, these represent different parameter points now; only the tree-level spectra are the same. To avoid ambiguity, we shall therefore use $(m_S^2)^{\text{tree}}$ since it is the input value for both schemes. In this way we see that two ways of defining the theory at tree-level can give, at times, drastically different results. In section 2.3 we provide as a consistency check a comparison of the approaches with a conversion of the parameters.

In figure 2, we show first M_h (left side) and M_S (right side) as a function of the trilinear coupling a_{SH} , at tree level (green curves) and at one loop in the standard (red curves) and modified (blue curves) schemes for the tadpoles. We choose here a scenario with a large Lagrangian mass term $m_S^{\text{tree}} = 2000\text{ GeV}$ and a non-zero trilinear self-coupling $a_S = 100\text{ GeV}$ for the singlet (and we also fix the renormalisation scale to be $Q = 2000\text{ GeV}$). Consequently, we find ourselves exactly in the dangerous region $0 < v \ll m_S$, *c.f.* eq. (2.17), and as expected from our theoretical discussion, we find that the standard treatment of the tadpoles breaks down. On the one hand, for M_h one can observe that the radiative corrections are larger in the standard approach and lead to larger variations of the loop-corrected mass than in the modified tadpole scheme. On the other hand, more strikingly, the results for M_S in the standard approach are manifestly spurious. Indeed, while the loop corrections in the modified scheme remain very small (the green tree-level and blue one-loop curves are almost superimposed), in the standard scheme the corrections are huge: for large $a_{SH} \gtrsim v$ – meaning not too small values of the singlet vev v_S – they already amount to several hundred GeV, and if one decreases a_{SH} (thereby increasing ΔM_S^2 , *c.f.* eq. (2.17)) the singlet pole mass becomes tachyonic below $a_{SH} = v$.

Next, in figure 3, we fix the trilinear coupling $a_{SH} = 150\text{ GeV}$ and now consider M_h (left) and M_S (right) as a function of the Lagrangian mass term m_S^{tree} . We also set $Q = m_S^{\text{tree}}$ and $a_S = 100\text{ GeV}$. Once again, with our choice of a non-zero singlet trilinear self-coupling a_S and relatively small a_{SH} – hence also a small singlet vev – we expect the standard approach to exhibit instabilities. For M_h (left




Figure 2: M_h (left) and M_S (right) as a function of a_{SH} . $m_S^{\text{tree}} = Q = 2000$ GeV, $a_S = 100$ GeV, $\lambda = 0.52$, $\lambda_{SH} = 0$, $\lambda_S = 1/24$. The tree-level values are shown with the green curves, while the red and blue curves correspond to the one-loop results using respectively the standard (eq. (2.2)) and modified (eq. (2.4)) treatments of tadpoles.




Figure 3: M_h (left) and M_S (right) as a function of m_S^{tree} . $Q = m_S^{\text{tree}}$, $a_{SH} = 150$ GeV, $a_S = 100$ GeV, $\lambda = 0.52$, $\lambda_{SH} = 0$, $\lambda_S = 1/24$. The colours for the different curves are the same as in figure 2.




Figure 4: M_h (left) and M_S (right) as a function of a_{SH} . $m_S^{\text{tree}} = 1000$ GeV, $Q = 5000$ GeV, $a_S = 0$ GeV, $\lambda = 0.52$, $\lambda_{SH} = 0$, $\lambda_S = 1/24$. The colours for the different curves are the same as in figure 2.




Figure 5: M_h (left) and M_S (right) as a function of a_{SH} . $m_S^{\text{tree}} = Q = 500$ GeV, $a_S = 0$ GeV, $\lambda_{SH} = 0$, $\lambda = 0.52$, $\lambda_S = 1/24$. The colours for the different curves are the same as in figure 2.

side of figure 3) both approaches behave relatively well and no instability seems to occur, although the radiative corrections are significantly larger in the standard scheme. However, for M_S the calculation in the standard approach (red curve) once again breaks down when m_S^{tree} is increased – equivalently for small v_S – while the loop corrections to M_S in the modified approach (blue curve) remain minute.

In figure 4, we illustrate the behaviour of eq. (2.19). We plot once more M_h (left) and M_S (right) as a function of the trilinear coupling a_{SH} , but now for a scenario where $a_S = 0$ (in order to avoid large corrections ΔM_S^2 in the standard scheme), and with $m_S^{\text{tree}} = 1000$ GeV and $Q = 5000$ GeV so as to increase the size of the logarithms $\overline{\log}(m_S^2)^{\text{tree}}$. For small values of a_{SH} , both schemes (red and blue curves) produce very similar results, however, as a_{SH} becomes larger the radiative corrections to M_h as well as M_S increase significantly in the modified tadpole scheme, leading to less reliable predictions (especially for $a_{SH} \gtrsim 300$ –400 GeV).

Finally, we present in figure 5 an example of scenario in which both ways to treat the tadpole contributions give reliable results. We take a small singlet mass parameter $m_S = 500$ GeV, set $a_S = 0$ and maintain $a_{SH} < 200$ GeV. We observe here that the radiative corrections to M_h and M_S remain well behaved in both approaches.

2.3 Comparisons at the same point

Here, for clarity (and as a consistency check) we shall follow the (first) prescription in section 2 and compare the two schemes for computing the one-loop masses in our toy model at the same parameter point. We consider the same input parameters as in figure 3, except that now we scan over the true $\overline{\text{MS}}$ mass m_S in both schemes. The calculation in the modified scheme is therefore identical to those in figure 3, but we then solve the tadpole equations for μ^2 and m_S^2 at the one-loop order to find the values of v, v_S ; while the value for v changes little, the equation for v_S becomes

$$0 = (m_S^2 + \lambda_{SH} v^2) v_S + \frac{1}{2} a_{SH} v^2 + 3 a_S v_S^2 + 4 \lambda_S v_S^3 + t_S(m_S^2). \quad (2.20)$$

We then use this new value for v_S to compute the tree and loop-level spectra in the standard scheme. In figure 6 we employ consistent tadpoles, so that we obtain a value for $(m_S^2)^{\text{tree}}$ which satisfies eq. (2.11b) and use this to compute the tree-level spectrum, and as input for the loop computation with the appropriate perturbative shifts to the loop mass; neglecting mixing between the light and heavy scalars we have

$$(M_S^2)^{\text{tree}} \simeq -\frac{a_{SH} v^2}{2 v_S} + v_S (3 a_S + 8 v_S \lambda_S), \quad (2.21a)$$

$$M_S^2 \simeq (M_S^2)^{\text{tree}} - \frac{1}{v_S} t_S((m_S^2)^{\text{tree}}) + \Pi_{SS}((M_S^2)^{\text{tree}}; (m_S^2)^{\text{tree}}). \quad (2.21b)$$




Figure 6: M_h (left) and M_S (right) as a function of the true $\overline{\text{MS}}$ parameter m_S in both the standard and modified schemes, where the standard scheme is performed according to the “consistent tadpole” prescription. Other parameters as in figure 3.




Figure 7: M_h (left) and M_S (right) as a function of the true $\overline{\text{MS}}$ parameter m_S in both the standard and modified schemes, where the standard scheme does not involve “consistent tadpoles” but the true $\overline{\text{MS}}$ mass m_S is used everywhere. Other parameters as in figure 3.

We have written $(m_S^2)^{\text{tree}}$ in the arguments of the tadpoles and self-energies to show the explicit dependence in the loop functions. In the left and right-hand plots of figure 6 we therefore see that the shift between m_S^2 and $(m_S^2)^{\text{tree}}$ becomes very large, and this leads to a breakdown of the (primitive) iterative algorithm that we use to solve for v_S , hence the standard scheme curves end near $m_S = 1250$ GeV, while the modified scheme has no such issue and the difference between loop-corrected and tree-level masses is negligible. This gives a different perspective on the general problem of calculating masses in such models. On the other hand, we see that, while the tree-level masses can differ significantly (even for the light “Higgs”) the loop masses agree to a high precision, as they should.

For a final comparison, we give in figure 7 the same computation but where, instead of “consistent tadpoles” we use the true $\overline{\text{MS}}$ mass m_S^2 obtained from eq. (2.20) in all of the loop functions so that

$$(M_S^2)^{\text{tree}} \simeq -\frac{a_{SH}v^2}{2v_S} + v_S(3a_S + 8v_S\lambda_S) - \frac{1}{v_S}t_S(m_S^2), \quad (2.22a)$$

$$M_S^2 \simeq (M_S^2)^{\text{tree}} + \Pi_{SS}((M_S^2)^{\text{tree}}; m_S^2). \quad (2.22b)$$

Aside from a shuffling of the tadpole term between the “tree-level” mass in the standard scheme, which now ensures that all of the curves on the right-hand side of figure 7 lie on top of each other (modulo the same proviso that the algorithm for finding v_S breaks down) the differences between these two versions of the standard scheme then only exist at two loops. From figure 7 it would seem that avoiding the consistent tadpoles would be preferable in these cases, but of course then the above equations mix tree-level and loop-level quantities, so we have problems with EFT matching at one loop (because subleading logarithms do not cancel) and infra-red issues at two loops.

3 Pole mass matching with tadpole insertions

When matching two theories via pole masses, care must be taken that subleading logarithms are correctly subtracted. The best way to do this is to expand the expressions on both sides of the matching relation in terms of the same parameters; the most efficient way to do this is to use those of the high-energy theory (HET) even though this adds a layer of complication because it is the SM parameters that we know from the bottom-up observations. To this end we require the shifts in the vacuum expectation value as well as gauge, Yukawa and of course quartic couplings.

The most straightforward way to match the vacuum-expectation value of the Higgs is via matching the Z mass, which gives (see *e. g.* Refs. [26, 29, 41]):

$$v_{\text{SM}}^2 = v_{\text{HET}}^2 + \frac{4}{g_Y^2 + g_2^2} \left[\hat{\Pi}_{ZZ}^{\text{HET}}(0) - \hat{\Pi}_{ZZ}^{\text{SM}}(0) \right] + \mathcal{O}(v^4). \quad (3.1)$$

If we match the one-loop Higgs mass in the SM to the HET, where the light Higgs mass at tree level is m_0 , then we have

$$2 \lambda_{\text{SM}} v_{\text{SM}}^2 + \hat{\Pi}_{hh}^{\text{SM}}(2 \lambda_{\text{SM}} v_{\text{SM}}^2) = m_0^2 + \hat{\Pi}_{hh}^{\text{HET}}(m_0^2) \quad (3.2a)$$

$$\lambda_{\text{SM}} = \frac{1}{2 v_{\text{HET}}^2} \left\{ m_0^2 + \hat{\Pi}_{hh}^{\text{HET}}(m_0^2) - \hat{\Pi}_{hh}^{\text{SM}}(m_0^2) - \frac{4 m_0^2}{v_{\text{HET}}^2 (g_Y^2 + g_2^2)} \left[\hat{\Pi}_{ZZ}^{\text{HET}}(0) - \hat{\Pi}_{ZZ}^{\text{SM}}(0) \right] \right\}. \quad (3.2b)$$

It should be noted that – in order to preserve gauge invariance, and cancel large logarithms exactly without introducing spurious subleading ones – the matching of the quartic coupling should be performed according to this equation, as opposed to performing some iteration, matching eigenvalues of the mass matrices, or separately matching the expectation values and Higgs mass (as performed in some codes) [32, 51]. With the prescription of including tadpole diagrams, this leads to

$$\hat{\Pi}_{hh} \equiv \Pi_{hh} - a^{hhk} \frac{1}{m_k^2} t_k, \quad \hat{\Pi}_{ZZ} \equiv \Pi_{ZZ} - g^{ZZk} \frac{1}{m_k^2} t_k. \quad (3.3)$$

In the SM with $\mathcal{L} \supset -\lambda_{\text{SM}} |H|^4$ we have

$$\hat{\Pi}_{hh}^{\text{SM}} \equiv \Pi_{hh}^{\text{SM}} - \frac{6 \lambda v}{m_h^2} t_h^{\text{SM}} = \Pi_{hh}^{\text{SM}} - \frac{3}{v} t_h^{\text{SM}}, \quad \hat{\Pi}_{ZZ}^{\text{SM}} \equiv \Pi_{ZZ}^{\text{SM}} - \frac{2 M_Z^2}{v m_h^2} t_k^{\text{SM}}, \quad (3.4)$$

and so

$$\hat{\Pi}_{hh}^{\text{SM}} - \frac{m_h^2}{M_Z^2} \hat{\Pi}_{ZZ}^{\text{SM}} = \Pi_{hh}^{\text{SM}} - \frac{m_h^2}{M_Z^2} \Pi_{ZZ}^{\text{SM}} - \frac{3}{v} t_h^{\text{SM}} + \frac{2}{v} t_h^{\text{SM}} = \Delta M_{\text{SM}}^2 - \frac{m_h^2}{M_Z^2} \Pi_{ZZ}^{\text{SM}}, \quad (3.5)$$

where the ΔM_{SM}^2 is now just the standard set of vacuum conditions as in eqs. (1.4) or (2.2). So what we have shown is that the modified treatment of tadpoles cancels out exactly in the matching of the light Higgs, *for the SM part*. Of course, the shift in the matching condition should only depend on the Lagrangian parameters, which are not affected by the treatment of tadpoles, so the same is true for the matching in the HET part *up to terms of higher order in v* .

We have already implicitly shown how the change in scheme affects the matching of the gauge bosons; now for fermions we have

$$\Gamma_{F_i F_j}(p) = i(\not{p} - m_F) \delta_{ij} + i \left[\not{p} \left(P_L \hat{\Sigma}_{ij}^L(p^2) + P_R \hat{\Sigma}_{ij}^R(p^2) \right) + P_L \hat{\Sigma}_{ij}^{SL}(p^2) + P_R \hat{\Sigma}_{ij}^{SR}(p^2) \right]. \quad (3.6)$$

For fermions at one loop we can write the mass-matrix corrections as

$$\delta m_F = -\Sigma^{SL} - \frac{1}{2} (\Sigma^R m + m \Sigma^L). \quad (3.7)$$

This means that our tadpole shift just affects

$$\delta \Sigma^{SL} = \delta \Sigma^{SR} = \frac{1}{m_k^2} y^{ijk} \frac{\partial V}{\partial \phi_k}, \quad (3.8)$$

where y^{ijk} are the Yukawa couplings, that can be written in terms of Weyl spinors $\{\psi_i\}$ as

$$\mathcal{L} \supset -\frac{1}{2} y^{ijk} \psi_i \psi_j \phi_k. \quad (3.9)$$

To match the Yukawa couplings via the pole masses of the quarks, the matching of the electroweak expectation value must also be included; working in the basis with diagonalised Yukawa couplings, we can match the diagonal elements as (using $Y^F \equiv y^{Fh}$ for h the SM Higgs and a general fermion F)

$$M_F = v Y^F - \Sigma^{SL} - \frac{1}{2} (\Sigma^R m + m \Sigma^L), \quad (3.10a)$$

$$\begin{aligned} Y_{\text{SM}}^F &= Y_{\text{HET}}^F + \frac{1}{v_{\text{HET}}} \left[(\delta m_F)^{\text{HET}} - (\delta m_F)^{\text{SM}} - \frac{1}{m_k^2} y_{\text{HET}}^{FFk} t_k + \frac{1}{m_h^2} Y_{\text{SM}}^F t_h^{\text{SM}} \right] \\ &\quad - \frac{Y_{\text{HET}}^F}{2 M_Z^2} \left[\hat{\Pi}_{ZZ}^{\text{HET}}(0) - \hat{\Pi}_{ZZ}^{\text{SM}}(0) \right] \\ &= Y_{\text{HET}}^F + \frac{1}{v_{\text{HET}}} \left[(\delta m_F)^{\text{HET}} - (\delta m_F)^{\text{SM}} - \frac{1}{m_k^2} y_{\text{HET}}^{FFk} t_k \right] - \frac{Y_{\text{HET}}^F}{2 M_Z^2} \left[\hat{\Pi}_{ZZ}^{\text{HET}}(0) - \Pi_{ZZ}^{\text{SM}}(0) \right], \end{aligned} \quad (3.10b)$$

where we once again see that the shift in the tadpole scheme cancels out exactly in the SM part. This procedure is particularly important since the shift to the expectation value arising in eq. (3.1) is very large, as discussed in the introduction. In this case, since the corrections to μ^2 – and therefore also to v^2 – are very large, it becomes impractical in an implementation to actually use the “correct” value of v^2 in the high-energy theory. Indeed, this can even become impossible, if $\delta\mu^2$ is such that μ^2 would become positive in the SM! Instead, provided we take v much less than the matching scale, we can just treat it as perturbation parameter to extract the SM values. In our numerical calculation in the next section we do exactly this: we just use the SM value of v in both high- and low-energy theories, but use the correct shifts of the expectation values in the matching of the parameters. This is very similar to a standard EFT calculation, which assumes *e.g.* in split supersymmetry that the heavy Higgs masses are tuned according to the mixing angle given as an input, and takes $v = 0$ explicitly, since we are not interested in corrections to Lagrangian parameters of order v^2/M^2 where M is the matching scale.

4 Application in the μ NMSSM

In the introduction, we explained that the modified treatment of tadpoles can be useful for stability under perturbation theory of heavy scalar masses when they are associated with a small expectation value. In section 2 we showed how it worked in practice in a toy model. In section 3 we described how, for theories where the new scalars are substantially above the electroweak scale, it can be practically applied via EFT matching of the pole masses. Here, we shall apply this technique to a real test case, the μ NMSSM.

4.1 NMSSM, μ NMSSM and GNMSSM

The superpotential of the most general form of the NMSSM – the GNMSSM – is [101, 102]

$$W_{\text{GNMSSM}} = Y_u Q \cdot H_u U - Y_d Q \cdot H_d D - Y_e L \cdot H_d E + \frac{1}{3} \kappa S^3 + (\mu + \lambda S) H_u \cdot H_d + \xi S + \frac{1}{2} \mu_S S^2$$

and the supersymmetry-breaking terms in the Higgs sector are

$$V_{\text{soft}} \supset m_S^2 |S|^2 + m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2 + \left(B_\mu H_u \cdot H_d + T_\lambda S H_u \cdot H_d + \frac{1}{3} T_\kappa S^3 + \frac{1}{2} B_S S^2 + \xi_S S + \text{h. c.} \right). \quad (4.1)$$

Once the singlet develops an expectation value, we can write effective terms

$$\mu_{\text{eff}} \equiv \mu + \frac{1}{\sqrt{2}} \lambda v_S, \quad B_{\text{eff}} \equiv B_\mu + \frac{1}{\sqrt{2}} T_\lambda v_S + \lambda \left(\xi + \frac{1}{\sqrt{2}} \mu_S v_S + \frac{1}{2} \kappa v_S^2 \right) \quad (4.2)$$

and the tadpole equations become

$$0 = -B_{\text{eff}} \cot \beta + m_{H_u}^2 + \mu_{\text{eff}}^2 - \frac{M_Z^2}{2} c_{2\beta} + \frac{1}{2} \lambda c_\beta^2, \quad (4.3a)$$

$$0 = -B_{\text{eff}} \tan \beta + m_{H_d}^2 + \mu_{\text{eff}}^2 + \frac{M_Z^2}{2} c_{2\beta} + \frac{1}{2} \lambda s_\beta^2, \quad (4.3b)$$

$$0 = v_S (B_S + m_S^2 + \mu_S^2 + 2 \kappa \xi) + \frac{1}{\sqrt{2}} v_S^2 (T_\kappa + 3 \kappa \mu_S) + \kappa^2 v_S^3 + \sqrt{2} \mu_S \xi + \sqrt{2} \xi_S + \frac{1}{2 \sqrt{2}} v^2 \left(2 \lambda \mu_{\text{eff}} - (T_\lambda + 2 \kappa \lambda v_S + \mu_S \lambda) s_{2\beta} \right). \quad (4.3c)$$

The first two lines are essentially modified versions of the MSSM tadpole equations with an extra term from the λ coupling. The third line, however, is the crucial one for our discussion. In a general non-supersymmetric theory, we can redefine singlet fields to remove their tadpole terms. However, in the GNMSSM, which has tadpole parameters ξ in the superpotential and ξ_S in the soft-breaking terms, we can only remove one of these, or the combination $\sqrt{2} \mu_S \xi + \sqrt{2} \xi_S$.

Clearly in the GNMSSM, it is most logical to choose a linear combination of the singlet tadpole terms ξ and ξ_S (or just one) as the variable to be eliminated by the tadpole equations. However, this is not possible in the NMSSM or μ NMSSM, since these terms vanish by the assumption of (at least partial) \mathbb{Z}_3 symmetry. Then aside from $(m_{H_u}^2, m_{H_d}^2)$ or (μ, B_μ) , the dimensionful parameter that we can now choose for elimination via the singlet tadpole equation is one of $\{m_S^2, \mu_{\text{eff}}, T_\lambda, T_\kappa\}$.

We are interested in the case that the singlet is rather heavier than the SM-like Higgs, so that $v^2/m_S^2 \ll 1$. This is clearly at best problematic in the NMSSM, since $\mu_{\text{eff}}, B_{\text{eff}} \propto v_S$ so if we imagine $v_S \sim \text{GeV}$ we will have very light higgsinos, pseudoscalar/charged Higgs and difficulties solving the tadpole equations. Hence we turn to the μ NMSSM, where we neglect all terms that break the \mathbb{Z}_3 symmetry except for μ and B_μ , and find

$$v_S \simeq -v^2 \left(\frac{2 \lambda \mu - T_\lambda s_{2\beta}}{2 \sqrt{2} m_S^2} \right), \quad (4.4)$$

where the true value can be found numerically.

The logical choice for this case is to solve for T_λ . In this case we have

$$\Delta T_\lambda = -\frac{2 \sqrt{2}}{v^2 s_{2\beta}} \frac{\partial \Delta V}{\partial v_S}, \quad (4.5)$$

and the terms in the mass matrix become

$$\mathcal{M}_{h_u^0 h_u^0}^2 \supset -\frac{v_S t_S^{(1)}}{v^2 s_\beta c_\beta} + \dots \propto \frac{t_S^{(1)}}{m_S^2}, \quad \mathcal{M}_{h_u^0 s_R}^2 = -\frac{m_S^2 v_S + t_S^{(1)}}{v s_\beta} + \dots \quad (4.6)$$

Note that this is in the “flavour basis” before we diagonalise the fields at tree level, so the contributions to the light Higgs and heavy singlet masses are $\propto t_S^{(1)}/m_S^2$.

On the other hand, this choice leads to a (potentially very) large quantum correction to T_λ . Suppose we want to investigate gauge-mediation scenarios where trilinears are small (nearly vanishing), or are otherwise specified by the top-down inputs – this would be completely inappropriate. Furthermore, we have to not only take into account shifts in the masses but also the *couplings* – this is moderately cumbersome to implement at one loop, but much more so if we want to compute the two-loop corrections. Indeed, it is not included in the algorithm to generate “consistent vacuum equations” of Refs. [27, 70], which assumes that the parameters that we solve the tadpole equations for only affect scalar masses.

To solve both of these issues the simplest choice is to solve for m_S^2 , and this leads to exactly the same problem as in the toy model, that the corrections to the singlet mass scale as $t_S^{(1)}/v_S$ leading to numerical instabilities for tiny v_S . Hence this model is an excellent prototype for comparing the different approaches to solving the tadpole equations.

4.2 Numerical comparison of tadpole schemes

In the μ NMSSM and GNMSSM, we not only have a Higgs sector, but also squarks, sleptons, a gluino and electroweakinos. In particular the colourful states have a large impact on the mass of the light Higgs, and, when they are heavy enough to be safe from current collider searches, they cause the “little hierarchy problem” to manifest itself. If we try to apply our modified tadpole scheme directly to these models, then we find all of the problems associated with this little hierarchy in our Higgs-mass calculation. Therefore it is only sensible to use EFT matching for the light Higgs mass. In this section we shall endeavour to show that with such an approach we can solve the technical difficulties with computing the masses of both light and heavy Higgs bosons.

We shall present here numerical investigations of several scenarios of the μ NMSSM and GNMSSM illustrating the differences between the two approaches to the treatment of tadpoles, both using EFT matching. For this, we compare results obtained using the original version of **SPheno** code obtained directly from **SARAH** (for the model **SMSSM**), as well as with a version of the **Fortran** output extensively modified according to the prescriptions described in section 3.[§]

In these calculations we must refer the reader again to our disclaimer, that we shall compare parameter points that generate the same *tree-level spectrum* in the two schemes, but that differ from each other at higher order; because this provides the clearest illustration of the problems faced (namely how to even define the parameter point). In contrast to the toy model, we will give no examples with a complete conversion of parameters, *i. e.* a comparison of both calculations at the same point, since the actual procedure of converting between the schemes is too onerous for technical reasons. In the **SARAH/SPheno** code, while a numerical solution of the tadpole equations (required for providing $\overline{\text{MS}}$ input to the standard scheme) is in principle possible, it is labourious and not implemented for loop computations where the variable to solve for is the vacuum expectation value.^{||} Therefore, again we take the tree-level value of v_S as input for the modified scheme, and treat it as the “all-orders” expectation value in the standard scheme (with consistent tadpoles) thus ensuring the same tree-level spectrum, but potentially vastly different results at one loop due to the large corrections to m_S^2 in the standard scheme. Again we stress that this is typical of the ambiguity in defining a parameter point that the phenomenologist is invited to suffer, thanks to the expedient in the standard scheme of hiding loop corrections in the definition of the expectation values.

[§] This private code is not intended for public release, although it is available on request from the authors. The new functionality should eventually be made available in a future release of **SARAH**.

^{||} This development in **SARAH** is envisioned in the future.




Figure 8: M_{h_1} (left) and M_{h_2} and M_{h_3} (right) as a function of v_S in a scenario of the μ NMSSM. The other inputs are taken as follows: $\lambda = \kappa = 0.1$, $T_\lambda = 200$ GeV, $T_\kappa = -10$ GeV, $\mu = 100$ GeV, $B_\mu = 6 \cdot 10^5$ GeV². Tree-level values are shown with green curves, while the red and blue curves correspond to the pole masses computed at one loop, respectively with the standard and modified approaches to the tadpoles. The colour coding of the lines remains the same for all figures in this section.

In section 4.2.1 we give an example of the above reasoning in the μ NMSSM. For illustration in section 4.2.2 we also give examples in the GNMSSM where we solve the tadpoles for the same variables ($m_{H_u}^2$, $m_{H_d}^2$, m_S^2) which allows us to compare several different scenarios.

4.2.1 μ NMSSM

In figure 8, we present the behaviour of the three CP-even mass eigenvalues – *i. e.* the lightest Higgs mass M_{h_1} (left side) and the masses of the additional CP-even states M_{h_2} and M_{h_3} (right side) – as a function of the singlet vev v_S in a μ NMSSM scenario, where the underlying parameters are given in the caption. The tree-level values are shown in green, while the one-loop results using the standard and the modified treatments of tadpoles are in red and blue respectively. We consider here a low range of values for v_S , so that, following our discussion in the previous section, we expect the standard approach to perform poorly for the singlet-like mass eigenstate. This is indeed what we observe if we turn to the right-side plot: for lower v_S ($\lesssim 0.1$ – 1 GeV) the singlet-like scalar is the heaviest eigenstate h_3 , while after level crossing it is h_2 for larger v_S . For the entire range of v_S the mass corrections in the standard approach are huge, and they grow as large as 50 TeV for $v_S = 0.001$ GeV – *i. e.* 250% of the tree-level result! On the other hand, if we look instead at the lightest Higgs boson h_1 , we find that the radiative corrections are somewhat larger with the modified treatment of the tadpole diagrams, and increase significantly with v_S in this scenario – due to the contributions from the tadpole diagram with a relatively large value of $T_\lambda = 200$ GeV and a relatively small tree-level mass of the singlet-like state.

4.2.2 GNMSSM

While the μ NMSSM provides an excellent prototype for the case of a heavy singlet with a small expectation value, where we cannot hide the loop corrections in a tadpole term, since it is a subset of the GNMSSM we can find more varied scenarios exhibiting the same behaviour. Of course, this is with the proviso that (with less justification in general) we restrict ourselves to solving the tadpole equations for m_S^2 .

We have devised three types of scenarios:

- Scenario 1: large singlet vev and intermediate λ ;
- Scenario 2: small singlet vev and small λ ;
- Scenario 3: small singlet vev but large λ .

| Scenario | 1 | 2 | 3 |
|-----------------------------|------------------|-------------------|-------------------|
| m_0 [GeV] | 2000 | 1500 | 1500 |
| λ | 0.1 [†] | 0.01 | 0.15 |
| κ | 0.005 | 0.05 | 0.05 |
| T_λ [GeV] | 1000 | 1000 [†] | 7500 [†] |
| v_S [GeV] | 3000 | 1.0 [†] | 1.0 [†] |
| μ [GeV] | 500 | 200 | 200 |
| μ_S [GeV] | 0 | -200 | -200 |
| ξ [GeV ²] | $1.0 \cdot 10^8$ | $1.7 \cdot 10^6$ | $5.0 \cdot 10^4$ |
| B_μ [GeV ²] | $2.0 \cdot 10^5$ | $1.0 \cdot 10^6$ | $4.0 \cdot 10^5$ |

Table 1: Definitions of the input parameters in the considered μ NMSSM scenarios. Some of the BSM parameters are not modified, and remain the same for the three scenario. Namely, we take: $\tan\beta = 10$, $m_{12} = 2$ TeV, $A_0 = 3$ TeV, $B_0 = 0$, $m_A = 500$ GeV, $T_\kappa = -0.5$ GeV. The renormalisation scale is kept at $Q = 3$ TeV for all computations. Finally, the numbers marked with a “†” are varied for some of the parameter scans.

Table 1 summarises the values taken for the BSM input parameters relevant for **SPheno** – note that we have adjusted the soft terms m_0 (scalar mass) and A_0 (scalar trilinear coupling) in order to obtain a mass for the lightest Higgs boson within the interval [123 GeV, 127 GeV]. We should also emphasise that the numbers in table 1 are given to **SPheno** as high-scale inputs (as this only requires a limited set of values). We then convert these into low-scale input parameters using the standard version of the μ NMSSM **SPheno** code, and the plots presented in the following are obtained by varying one of the low-scale inputs. In light of the analytic expressions in the previous section, we can expect the two approaches to the tadpoles to give relatively similar results in scenario 1, where the singlet vev is large. However, in scenarios 2 and 3, the singlet vev is taken to be small, so that the differences between the two schemes should be more pronounced. Scenario 3 furthermore allows us to investigate the effect of increasing the coupling λ .

We show first in figure 9 the behaviour of the lightest Higgs mass M_{h_1} (left side) and of the additional CP-even Higgs-boson masses M_{h_2} and M_{h_3} (right side) as a function of the superpotential coupling λ . Among the two BSM states h_2 and h_3 , the former is singlet-like while the latter is doublet-like, in this figure. As can be seen in the right-hand side plot of figure 9, the heavy Higgs bosons receive only minute mass corrections in either of the approaches for tadpoles. For the lightest scalar mass M_{h_1} , the results in the two schemes are also in excellent agreement. However, we have cut off the plot before $\lambda = 0.19$ because beyond this value perturbativity is lost: in the *standard approach* the singlet-like pseudoscalar Higgs becomes tachyonic at one loop (from a tree-level mass of 750 GeV!). If we continued the plot into this regime we would see the predictions diverging, with the standard approach predicting ever decreasing masses and the modified approach increasing ones for larger λ (compare 104 GeV and 138 GeV respectively for $\lambda = 0.3$).

Next, we turn to scenario 2, *i. e.* we consider a small $\lambda = 0.01$ and small singlet vev $v_S = 1$ GeV. Figure 10 shows the behaviour of the CP-even masses as a function of the soft trilinear coupling T_λ , at tree level and one loop (the colouring of the curves is the same as previously explained). We should emphasise that we have made sure to fulfill constraints from vacuum stability (and the absence of a charge-breaking minimum) on T_λ – see Ref. [63] – and the tree-level mass of the charged Higgs boson remains positive for the entire range of T_λ investigated here. While for M_{h_1} (left side) and M_{h_2} (lower curves of the right-side plot) it seems essentially impossible to distinguish the two approaches to the tadpole treatment, the radiative corrections to M_{h_3} – the mass of the singlet-like scalar – are clearly much larger with the standard method, and the result of the modified scheme is certainly more reliable. As a concrete comparison, we have for the intermediate value $T_\lambda = 2$ TeV a one-loop correction to M_{h_3} of 2752 GeV (*i. e.* 24% of the tree-level result) in the standard approach, but only of -4.5 GeV in the modified scheme.




Figure 9: M_{h_1} (left) and M_{h_2} and M_{h_3} (right) as a function of λ , in scenario 1. The other inputs are taken as in table 1. Tree-level values are shown with green curves, while the red and blue curves correspond to the pole masses computed at one loop, respectively with the standard and modified approaches to the tadpoles.




Figure 10: M_{h_1} (left) and M_{h_2} and M_{h_3} (right) as a function of the soft trilinear coupling T_λ , in scenario 2. The values of the other BSM parameters are taken as in table 1.




Figure 11: M_{h_1} (left) and M_{h_2} and M_{h_3} (right) as a function of v_S , in scenario 2. Input values for the other BSM parameters are given in table 1.




Figure 12: M_{h_1} (left) and M_{h_2} and M_{h_3} (right) as a function of T_λ , in scenario 3. The other BSM inputs are taken as in table 1.




Figure 13: M_{h_1} (left) and M_{h_2} and M_{h_3} (right) as a function of v_S , in scenario 3. The values of the other relevant inputs are given in table 1.

We can confirm that the large difference between the two treatments of the tadpoles arises from the small value of the singlet vev v_S . Indeed, in figure 11, we present the same three CP-even scalar masses for v_S varying between 0.5 and 100 GeV. One can observe that the results using both approaches for all three masses are in good agreement for large values of the singlet vev. A short comment should be made for M_{h_1} : indeed, as v_S increases the results from the two schemes seem to grow apart, and it is somewhat difficult to determine which one should be trusted more in this case. We note that the radiative corrections to M_{h_1} keep increasing with v_S in the standard approach while their size remains relatively stable in the modified scheme. On the other hand, if we consider the situation for $v_S \gtrsim 0.5$ GeV, the breakdown of the standard approach for small singlet vevs becomes obvious. Indeed, considering the different results for the mass M_{h_3} of the CP-even singlet-like scalar at $v_S = 0.5$ GeV, the one-loop corrections in the standard scheme amount to 6.5 TeV – in other words, 40% of the tree-level result – compared to only -3.3 GeV (-0.02% of the tree-level mass) in the modified scheme.

Lastly, we consider the type of scenario 3, *i. e.* what happens if we keep a small singlet vev $v_S = 1$ GeV but increase the coupling λ to 0.15. In figure 12, we present the CP-even scalar masses as a function of T_λ – having once again made sure to maintain vacuum stability [63]. Considering first the masses of the two doublet-like scalars h_1 and h_2 , we observe an excellent agreement of the results from the two tadpole schemes for low to intermediate values of T_λ – for $0 \leq T_\lambda \lesssim 4$ TeV. However, as T_λ becomes larger, the corrections to M_{h_1} and M_{h_2} in the modified approach start growing out of control. This appears similar to the loss of accuracy of the modified scheme that we encountered in the toy model of section 2 when increasing the trilinear coupling a_{SH} , which plays the same role as T_λ – see eq. (2.19) and figure 4. Turning however to the singlet-like mass M_{h_3} we find (as in figure 10 for scenario 2) that the radiative corrections are huge with the standard treatment of tadpoles, but remain well-behaved

with the modified one. Interestingly, having increased the value of λ has not made the breakdown of the standard calculation for the singlet-like mass more severe than in scenario 2. Nevertheless, the one-loop result M_{h_3} using the modified tadpole scheme is undoubtedly more reliable here.

Finally, we present in figure 13 the behaviour of the CP-even scalar masses as a function of the singlet vev v_S – restricting our attention to the low range $0.5 \text{ GeV} \leq v_S \leq 5 \text{ GeV}$. As can be read from table 1, we have chosen for this figure a large value of the soft trilinear coupling $T_\lambda = 7.5 \text{ TeV}$, which corresponds to the right parts of the plots in figure 12. Therefore, it is not surprising that we observe some discrepancy between the results of the two tadpole schemes for all three masses, as discussed above. More interestingly, we can compare the size of the loop corrections to M_{h_3} in the two approaches, as we vary v_S . On the one hand, in the standard approach, the one-loop corrections increase from 2.3 TeV (19% of the tree-level result) for $v_S = 2.5 \text{ GeV}$ to as much as 9 TeV (40% of the tree-level mass) for $v_S = 0.75 \text{ GeV}$, for instance. On the other hand, in the modified scheme, the effects remain minute and vary from -46 GeV for $v_S = 2.5 \text{ GeV}$ to -3.6 GeV for $v_S = 0.75 \text{ GeV}$ (this amounts to -0.38% and -0.02% of the results at tree level, respectively).

5 Conclusions

We have shown the advantages and limitations of taking a different prescription for the solution of tadpole equations. In contrast to previous applications of this technique, in the SM or as a measure of fine-tuning, we have shown that it can be very useful when new scalars having a small expectation value are present in the theory, and in the case that they are much heavier than the electroweak scale, it is best employed via the matching of pole masses in an EFT approach. While this technique offers the advantages of perturbative stability for the heavy scalar masses, easy generalisability (the corrections are simply computed diagrammatically rather than via taking derivatives of the tadpole equations) and gauge invariance, it can also lead to numerical instabilities in extracting the *light* Higgs mass, and the loss of the ability to match the electroweak expectation value.

In future work, other than a general numerical implementation in SARAH, it would be interesting to explore a hybrid approach (along the lines of option 1 described at the end of the introduction), where only the electroweak expectation value is fixed by appropriate counterterms. On the other hand, we intend to consider the corrections at two loops in this approach, and we shall also provide general expressions for the one-loop self-energies which are explicitly gauge independent.

Acknowledgements

We would like to thank Sven Heinemeyer and Pietro Slavich for interesting discussions. MDG acknowledges support from the grant ‘‘HiggsAutomator’’ of the Agence Nationale de la Recherche (ANR) (ANR-15-CE31-0002). JB is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy – EXC 2121 ‘‘Quantum Universe’’ – 390833306. The work of SP is supported by the BMBF Grant No. 05H18PACC2. This project received support from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860881-HIDDeN.

References

- [1] ATLAS Collaboration, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”, *Phys. Lett.* **B716**, 1 (2012), [arXiv:1207.7214](#). [p 2]
- [2] CMS Collaboration, “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC”, *Phys. Lett.* **B716**, 30 (2012), [arXiv:1207.7235](#). [p 2]
- [3] ATLAS Collaboration, CMS Collaboration, “Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments”, *Phys. Rev. Lett.* **114**, 191803 (2015), [arXiv:1503.07589](#). [p 2]
- [4] ATLAS Collaboration, CMS Collaboration, “Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV”, *JHEP* **08**, 045 (2016), [arXiv:1606.02266](#). [p 2]
- [5] CMS Collaboration, A. M. Sirunyan, *et al.*, “Combined measurements of Higgs boson couplings in proton–proton collisions at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J.* **C79**, 421 (2019), [arXiv:1809.10733](#). [p 2]
- [6] ATLAS, G. Aad, *et al.*, “Combined measurements of Higgs boson production and decay using up to 80 fb^{-1} of proton-proton collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS experiment”, *Phys. Rev. D* **101**, 012002 (2020), [arXiv:1909.02845](#). [p 2]
- [7] P. Slavich, S. Heinemeyer (eds.), E. Bagnaschi, *et al.*, “Higgs-mass predictions in the MSSM and beyond”, (2020), [arXiv:2012.15629](#). [p 2]
- [8] W. Hollik, S. Paßehr, “Two-loop top-Yukawa-coupling corrections to the Higgs boson masses in the complex MSSM”, *Phys. Lett.* **B733**, 144 (2014), [arXiv:1401.8275](#). [p 2]
- [9] S. Borowka, T. Hahn, S. Heinemeyer, G. Heinrich, W. Hollik, “Momentum-dependent two-loop QCD corrections to the neutral Higgs-boson masses in the MSSM”, *Eur. Phys. J.* **C74**, 2994 (2014), [arXiv:1404.7074](#). [p 2]
- [10] E. Bagnaschi, G. F. Giudice, P. Slavich, A. Strumia, “Higgs Mass and Unnatural Supersymmetry”, *JHEP* **09**, 092 (2014), [arXiv:1407.4081](#). [p 2]
- [11] W. Hollik, S. Paßehr, “Higgs boson masses and mixings in the complex MSSM with two-loop top-Yukawa-coupling corrections”, *JHEP* **10**, 171 (2014), [arXiv:1409.1687](#). [p 2]
- [12] G. Degrossi, S. Di Vita, P. Slavich, “Two-loop QCD corrections to the MSSM Higgs masses beyond the effective-potential approximation”, *Eur. Phys. J.* **C75**, 61 (2015), [arXiv:1410.3432](#). [p 2]
- [13] M. Goodsell, K. Nickel, F. Staub, “Two-Loop Higgs mass calculations in supersymmetric models beyond the MSSM with SARAH and SPheno”, *Eur. Phys. J.* **C75**, 32 (2015), [arXiv:1411.0675](#). [pp 2, 5]
- [14] M. D. Goodsell, K. Nickel, F. Staub, “Two-loop corrections to the Higgs masses in the NMSSM”, *Phys. Rev.* **D91**, 035021 (2015), [arXiv:1411.4665](#). [p 2]
- [15] M. Mühlleitner, D. T. Nhung, H. Rzehak, K. Walz, “Two-loop contributions of the order $\mathcal{O}(\alpha_t\alpha_s)$ to the masses of the Higgs bosons in the CP-violating NMSSM”, *JHEP* **05**, 128 (2015), [arXiv:1412.0918](#). [p 2]
- [16] M. Goodsell, K. Nickel, F. Staub, “Generic two-loop Higgs mass calculation from a diagrammatic approach”, *Eur. Phys. J.* **C75**, 290 (2015), [arXiv:1503.03098](#). [pp 2, 5]
- [17] S. Borowka, T. Hahn, S. Heinemeyer, G. Heinrich, W. Hollik, “Renormalization scheme dependence of the two-loop QCD corrections to the neutral Higgs-boson masses in the MSSM”, *Eur. Phys. J.* **C75**, 424 (2015), [arXiv:1505.03133](#). [p 2]
- [18] F. Staub, P. Athron, U. Ellwanger, R. Gröber, M. Mühlleitner, P. Slavich, A. Voigt, “Higgs mass predictions of public NMSSM spectrum generators”, *Comput. Phys. Commun.* **202**, 113 (2016), [arXiv:1507.05093](#). [p 2]
- [19] T. Hahn, S. Paßehr, “Implementation of the $\mathcal{O}(\alpha_t^2)$ MSSM Higgs-mass corrections in FeynHiggs”, *Comput. Phys. Commun.* **214**, 91 (2017), [arXiv:1508.00562](#). [p 2]
- [20] G. Lee, C. E. M. Wagner, “Higgs bosons in heavy supersymmetry with an intermediate m_A ”, *Phys. Rev.* **D92**, 075032 (2015), [arXiv:1508.00576](#). [p 2]
- [21] M. D. Goodsell, K. Nickel, F. Staub, “The Higgs Mass in the MSSM at two-loop order beyond minimal flavour violation”, *Phys. Lett.* **B758**, 18 (2016), [arXiv:1511.01904](#). [p 2]
- [22] P. Drechsel, L. Galeta, S. Heinemeyer, G. Weiglein, “Precise Predictions for the Higgs-Boson Masses in the NMSSM”, *Eur. Phys. J.* **C77**, 42 (2017), [arXiv:1601.08100](#). [p 2]
- [23] M. D. Goodsell, F. Staub, “The Higgs mass in the CP violating MSSM, NMSSM, and beyond”, *Eur. Phys. J.* **C77**, 46 (2017), [arXiv:1604.05335](#). [p 2]
- [24] J. Braathen, M. D. Goodsell, P. Slavich, “Leading two-loop corrections to the Higgs boson masses in SUSY models with Dirac gauginos”, *JHEP* **09**, 045 (2016), [arXiv:1606.09213](#). [p 2]
- [25] H. Bahl, W. Hollik, “Precise prediction for the light MSSM Higgs boson mass combining effective field theory and fixed-order calculations”, *Eur. Phys. J.* **C76**, 499 (2016), [arXiv:1608.01880](#). [p 2]
- [26] P. Athron, J.-h. Park, T. Stuedtner, D. Stöckinger, A. Voigt, “Precise Higgs mass calculations in (non-)minimal supersymmetry at both high and low scales”, *JHEP* **01**, 079 (2017), [arXiv:1609.00371](#). [pp 2, 5, 13]
- [27] J. Braathen, M. D. Goodsell, “Avoiding the Goldstone Boson Catastrophe in general renormalisable field theories at two loops”, *JHEP* **12**, 056 (2016), [arXiv:1609.06977](#). [pp 2, 3, 4, 5, 6, 8, 16]
- [28] P. Drechsel, R. Gröber, S. Heinemeyer, M. Mühlleitner, H. Rzehak, G. Weiglein, “Higgs-Boson Masses and Mixing Matrices in the NMSSM: Analysis of On-Shell Calculations”, *Eur. Phys. J.* **C77**, 366 (2017), [arXiv:1612.07681](#). [p 2]
- [29] F. Staub, W. Porod, “Improved predictions for intermediate and heavy Supersymmetry in the MSSM and beyond”, *Eur. Phys. J.* **C77**, 338 (2017), [arXiv:1703.03267](#). [pp 2, 13]
- [30] E. Bagnaschi, J. Pardo Vega, P. Slavich, “Improved determination of the Higgs mass in the MSSM with heavy superpartners”, *Eur. Phys. J.* **C77**, 334 (2017), [arXiv:1703.08166](#). [p 2]
- [31] S. Paßehr, G. Weiglein, “Two-loop top and bottom Yukawa corrections to the Higgs-boson masses in the complex MSSM”, *Eur. Phys. J.* **C78**, 222 (2018), [arXiv:1705.07909](#). [p 2]

- [32] H. Bahl, S. Heinemeyer, W. Hollik, G. Weiglein, “Reconciling EFT and hybrid calculations of the light MSSM Higgs-boson mass”, *Eur. Phys. J. C* **78**, 57 (2018), [arXiv:1706.00346](#). [pp 2, 13]
- [33] J. Braathen, M. D. Goodsell, F. Staub, “Supersymmetric and non-supersymmetric models without catastrophic Goldstone bosons”, *Eur. Phys. J. C* **77**, 757 (2017), [arXiv:1706.05372](#). [pp 2, 3, 5]
- [34] R. V. Harlander, J. Klappert, A. Voigt, “Higgs mass prediction in the MSSM at three-loop level in a pure $\overline{\text{DR}}$ context”, *Eur. Phys. J. C* **77**, 814 (2017), [arXiv:1708.05720](#). [p 2]
- [35] P. Athron, M. Bach, D. Harries, T. Kwasnitza, J.-h. Park, D. Stöckinger, A. Voigt, J. Ziebell, “FlexibleSUSY 2.0: Extensions to investigate the phenomenology of SUSY and non-SUSY models”, *Comput. Phys. Commun.* **230**, 145 (2018), [arXiv:1710.03760](#). [p 2]
- [36] T. Biekötter, S. Heinemeyer, C. Muñoz, “Precise prediction for the Higgs-boson masses in the $\mu\nu$ SSM”, *Eur. Phys. J. C* **78**, 504 (2018), [arXiv:1712.07475](#). [p 2]
- [37] S. Borowka, S. Paßehr, G. Weiglein, “Complete two-loop QCD contributions to the lightest Higgs-boson mass in the MSSM with complex parameters”, *Eur. Phys. J. C* **78**, 576 (2018), [arXiv:1802.09886](#). [p 2]
- [38] D. Stöckinger, J. Unger, “Three-loop MSSM Higgs-boson mass predictions and regularization by dimensional reduction”, *Nucl. Phys. B* **935**, 1 (2018), [arXiv:1804.05619](#). [p 2]
- [39] H. Bahl, W. Hollik, “Precise prediction of the MSSM Higgs boson masses for low M_A ”, *JHEP* **07**, 182 (2018), [arXiv:1805.00867](#). [p 2]
- [40] R. V. Harlander, J. Klappert, A. D. Ochoa Franco, A. Voigt, “The light CP-even MSSM Higgs mass resummed to fourth logarithmic order”, *Eur. Phys. J. C* **78**, 874 (2018), [arXiv:1807.03509](#). [p 2]
- [41] J. Braathen, M. D. Goodsell, P. Slavich, “Matching renormalisable couplings: simple schemes and a plot”, *Eur. Phys. J. C* **79**, 669 (2019), [arXiv:1810.09388](#). [pp 2, 3, 13]
- [42] M. Gabelmann, M. Mühlleitner, F. Staub, “Automatised matching between two scalar sectors at the one-loop level”, *Eur. Phys. J. C* **79**, 163 (2019), [arXiv:1810.12326](#). [p 2]
- [43] H. Bahl, T. Hahn, S. Heinemeyer, W. Hollik, S. Paßehr, H. Rzehak, G. Weiglein, “Precision calculations in the MSSM Higgs-boson sector with FeynHiggs 2.14”, *Comput. Phys. Commun.* **249**, 107099 (2020), [arXiv:1811.09073](#). [p 2]
- [44] H. Bahl, “Pole mass determination in presence of heavy particles”, *JHEP* **02**, 121 (2019), [arXiv:1812.06452](#). [p 2]
- [45] T. N. Dao, R. Gröber, M. Krause, M. Mühlleitner, H. Rzehak, “Two-loop $\mathcal{O}(\alpha_t^2)$ corrections to the neutral Higgs boson masses in the CP-violating NMSSM”, *JHEP* **08**, 114 (2019), [arXiv:1903.11358](#). [p 2]
- [46] E. Bagnaschi, G. Degrandi, S. Paßehr, P. Slavich, “Full two-loop QCD corrections to the Higgs mass in the MSSM with heavy superpartners”, *Eur. Phys. J. C* **79**, 910 (2019), [arXiv:1908.01670](#). [p 2]
- [47] M. D. Goodsell, S. Paßehr, “All two-loop scalar self-energies and tadpoles in general renormalisable field theories”, *Eur. Phys. J. C* **80**, 417 (2020), [arXiv:1910.02094](#). [p 2]
- [48] R. V. Harlander, J. Klappert, A. Voigt, “The light CP-even MSSM Higgs mass including $\text{N}^3\text{LO}+\text{N}^3\text{LL}$ QCD corrections”, *Eur. Phys. J. C* **80**, 186 (2020), [arXiv:1910.03595](#). [p 2]
- [49] H. Bahl, S. Heinemeyer, W. Hollik, G. Weiglein, “Theoretical uncertainties in the MSSM Higgs boson mass calculation”, *Eur. Phys. J. C* **80**, 497 (2020), [arXiv:1912.04199](#). [p 2]
- [50] H. Bahl, I. Sobolev, G. Weiglein, “Precise prediction for the mass of the light MSSM Higgs boson for the case of a heavy gluino”, *Phys. Lett. B* **808**, 135644 (2020), [arXiv:1912.10002](#). [p 2]
- [51] T. Kwasnitza, D. Stöckinger, A. Voigt, “Improved MSSM Higgs mass calculation using the 3-loop FlexibleEFTHiggs approach including x_t -resummation”, *JHEP* **07**, 197 (2020), [arXiv:2003.04639](#). [pp 2, 13]
- [52] H. Bahl, I. Sobolev, G. Weiglein, “The light MSSM Higgs boson mass for large $\tan\beta$ and complex input parameters”, *Eur. Phys. J. C* **80**, 1063 (2020), [arXiv:2009.07572](#). [p 2]
- [53] H. Bahl, I. Sobolev, “Two-loop matching of renormalizable operators: general considerations and applications”, *JHEP* **03**, 286 (2021), [arXiv:2010.01989](#). [p 2]
- [54] H. Bahl, N. Murphy, H. Rzehak, “Hybrid calculation of the MSSM Higgs boson masses using the complex THDM as EFT”, *Eur. Phys. J. C* **81**, 128 (2021), [arXiv:2010.04711](#). [p 2]
- [55] G. Degrandi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori, A. Strumia, “Higgs mass and vacuum stability in the Standard Model at NNLO”, *JHEP* **08**, 098 (2012), [arXiv:1205.6497](#). [p 2]
- [56] D. Buttazzo, G. Degrandi, P. P. Giardino, G. F. Giudice, F. Sala, A. Salvio, A. Strumia, “Investigating the near-criticality of the Higgs boson”, *JHEP* **12**, 089 (2013), [arXiv:1307.3536](#). [p 2]
- [57] B. A. Kniehl, A. F. Pikelner, O. L. Veretin, “Two-loop electroweak threshold corrections in the Standard Model”, *Nucl. Phys. B* **896**, 19 (2015), [arXiv:1503.02138](#). [pp 2, 4]
- [58] B. A. Kniehl, A. F. Pikelner, O. L. Veretin, “mr: a C++ library for the matching and running of the Standard Model parameters”, *Comput. Phys. Commun.* **206**, 84 (2016), [arXiv:1601.08143](#). [p 2]
- [59] S. P. Martin, D. G. Robertson, “Standard Model parameters in the tadpole-free pure $\overline{\text{MS}}$ scheme”, (2019), [arXiv:1907.02500](#). [p 2]
- [60] C. Coriano, L. Delle Rose, C. Marzo, “Constraints on abelian extensions of the Standard Model from two-loop vacuum stability and $U(1)_{B-L}$ ”, *JHEP* **02**, 135 (2016), [arXiv:1510.02379](#). [p 2]
- [61] J. Braathen, M. D. Goodsell, M. E. Krauss, T. Opferkuch, F. Staub, “ N -loop running should be combined with N -loop matching”, *Phys. Rev. D* **97**, 015011 (2018), [arXiv:1711.08460](#). [p 2]
- [62] M. E. Krauss, T. Opferkuch, F. Staub, “The Ultraviolet Landscape of Two-Higgs Doublet Models”, *Eur. Phys. J. C* **78**, 1020 (2018), [arXiv:1807.07581](#). [p 2]
- [63] W. G. Hollik, S. Liebler, G. Moortgat-Pick, S. Paßehr, G. Weiglein, “Phenomenology of the inflation-inspired NMSSM at the electroweak scale”, *Eur. Phys. J. C* **79**, 75 (2019), [arXiv:1809.07371](#). [pp 2, 5, 18, 20]

- [64] J.-W. Wang, X.-J. Bi, P.-F. Yin, Z.-H. Yu, “Impact of Fermionic Electroweak Multiplet Dark Matter on Vacuum Stability with One-loop Matching”, *Phys. Rev. D* **99**, 055009 (2019), [arXiv:1811.08743](#). [p 2]
- [65] W. G. Hollik, G. Weiglein, J. Wittbrodt, “Impact of Vacuum Stability Constraints on the Phenomenology of Supersymmetric Models”, *JHEP* **03**, 109 (2019), [arXiv:1812.04644](#). [p 2]
- [66] S. P. Martin, D. G. Robertson, “Higgs boson mass in the Standard Model at two-loop order and beyond”, *Phys. Rev.* **D90**, 073010 (2014), [arXiv:1407.4336](#). [p 2]
- [67] S. P. Martin, “Complete Two Loop Effective Potential Approximation to the Lightest Higgs Scalar Boson Mass in Supersymmetry”, *Phys. Rev.* **D67**, 095012 (2003), [arXiv:hep-ph/0211366](#). [p 3]
- [68] S. P. Martin, “Taming the Goldstone contributions to the effective potential”, *Phys. Rev.* **D90**, 016013 (2014), [arXiv:1406.2355](#). [p 3]
- [69] J. Elias-Miro, J. R. Espinosa, T. Konstandin, “Taming Infrared Divergences in the Effective Potential”, *JHEP* **08**, 034 (2014), [arXiv:1406.2652](#). [p 3]
- [70] N. Kumar, S. P. Martin, “Resummation of Goldstone boson contributions to the MSSM effective potential”, *Phys. Rev.* **D94**, 014013 (2016), [arXiv:1605.02059](#). [pp 3, 16]
- [71] M. Sperling, D. Stöckinger, A. Voigt, “Renormalization of vacuum expectation values in spontaneously broken gauge theories”, *JHEP* **07**, 132 (2013), [arXiv:1305.1548](#). [p 3]
- [72] M. Sperling, D. Stöckinger, A. Voigt, “Renormalization of vacuum expectation values in spontaneously broken gauge theories: Two-loop results”, *JHEP* **01**, 068 (2014), [arXiv:1310.7629](#). [p 3]
- [73] G. Belanger, K. Benakli, M. Goodsell, C. Moura, A. Pukhov, “Dark Matter with Dirac and Majorana Gaugino Masses”, *JCAP* **08**, 027 (2009), [arXiv:0905.1043](#). [p 3]
- [74] K. Benakli, M. D. Goodsell, A.-K. Maier, “Generating μ and $B\mu$ in models with Dirac Gauginos”, *Nucl. Phys. B* **851**, 445 (2011), [arXiv:1104.2695](#). [p 3]
- [75] K. Benakli, M. D. Goodsell, F. Staub, “Dirac Gauginos and the 125 GeV Higgs”, *JHEP* **06**, 073 (2013), [arXiv:1211.0552](#). [p 3]
- [76] M. D. Goodsell, S. Kraml, H. Reyes-González, S. L. Williamson, “Constraining Electroweakinos in the Minimal Dirac Gaugino Model”, *SciPost Phys.* **9**, 047 (2020), [arXiv:2007.08498](#). [p 3]
- [77] J. Fleischer, F. Jegerlehner, “Radiative Corrections to Higgs Decays in the Extended Weinberg-Salam Model”, *Phys. Rev. D* **23**, 2001 (1981). [p 3]
- [78] F. Jegerlehner, M. Yu. Kalmykov, O. Veretin, “MS versus pole masses of gauge bosons: Electroweak bosonic two loop corrections”, *Nucl. Phys.* **B641**, 285 (2002), [arXiv:hep-ph/0105304](#). [p 4]
- [79] F. Jegerlehner, M. Yu. Kalmykov, O. Veretin, “Full two loop electroweak corrections to the pole masses of gauge bosons”, *Nucl. Phys. Proc. Suppl.* **116**, 382 (2003), [arXiv:hep-ph/0212003](#), [382(2002)]. [p 4]
- [80] F. Jegerlehner, M. Yu. Kalmykov, O. Veretin, “MS-bar versus pole masses of gauge bosons. 2. Two loop electroweak fermion corrections”, *Nucl. Phys.* **B658**, 49 (2003), [arXiv:hep-ph/0212319](#). [p 4]
- [81] F. Jegerlehner, M. Y. Kalmykov, “O($\alpha\alpha(s)$) correction to the pole mass of the t quark within the standard model”, *Nucl. Phys. B* **676**, 365 (2004), [arXiv:hep-ph/0308216](#). [p 4]
- [82] F. Bezrukov, M. Yu. Kalmykov, B. A. Kniehl, M. Shaposhnikov, “Higgs Boson Mass and New Physics”, *JHEP* **10**, 140 (2012), [arXiv:1205.2893](#), [275(2012)]. [p 4]
- [83] M. Krause, R. Lorenz, M. Mühlleitner, R. Santos, H. Ziesche, “Gauge-independent Renormalization of the 2-Higgs-Doublet Model”, *JHEP* **09**, 143 (2016), [arXiv:1605.04853](#). [p 4]
- [84] A. Denner, L. Jenniches, J.-N. Lang, C. Sturm, “Gauge-independent \overline{MS} renormalization in the 2HDM”, *JHEP* **09**, 115 (2016), [arXiv:1607.07352](#). [p 4]
- [85] L. Altenkamp, S. Dittmaier, H. Rzehak, “Renormalization schemes for the Two-Higgs-Doublet Model and applications to $h \rightarrow WW/ZZ \rightarrow 4$ fermions”, *JHEP* **09**, 134 (2017), [arXiv:1704.02645](#). [p 4]
- [86] M. Krause, M. Mühlleitner, “Impact of Electroweak Corrections on Neutral Higgs Boson Decays in Extended Higgs Sectors”, *JHEP* **04**, 083 (2020), [arXiv:1912.03948](#). [p 4]
- [87] P. Chankowski, S. Pokorski, J. Rosiek, “Complete on-shell renormalization scheme for the minimal supersymmetric Higgs sector”, *Nucl. Phys.* **B423**, 437 (1994), [arXiv:hep-ph/9303309](#). [p 4]
- [88] A. Dabelstein, “The One loop renormalization of the MSSM Higgs sector and its application to the neutral scalar Higgs masses”, *Z. Phys.* **C67**, 495 (1995), [arXiv:hep-ph/9409375](#). [p 4]
- [89] A. Freitas, D. Stöckinger, “Gauge dependence and renormalization of tan beta in the MSSM”, *Phys. Rev. D* **66**, 095014 (2002), [arXiv:hep-ph/0205281](#). [p 4]
- [90] S. Kanemura, Y. Okada, E. Senaha, C. P. Yuan, “Higgs coupling constants as a probe of new physics”, *Phys. Rev. D* **70**, 115002 (2004), [arXiv:hep-ph/0408364](#). [p 4]
- [91] M. Farina, D. Pappadopulo, A. Strumia, “A modified naturalness principle and its experimental tests”, *JHEP* **08**, 022 (2013), [arXiv:1303.7244](#). [pp 4, 5]
- [92] W. Porod, “SPheno, a program for calculating supersymmetric spectra, SUSY particle decays and SUSY particle production at e^+e^- colliders”, *Comput. Phys. Commun.* **153**, 275 (2003), [arXiv:hep-ph/0301101](#). [p 5]
- [93] W. Porod, F. Staub, “SPheno 3.1: Extensions including flavour, CP-phases and models beyond the MSSM”, *Comput. Phys. Commun.* **183**, 2458 (2012), [arXiv:1104.1573](#). [p 5]
- [94] F. Staub, “SARAH”, (2008), [arXiv:0806.0538](#). [p 5]
- [95] F. Staub, “From Superpotential to Model Files for FeynArts and CalcHep/CompHep”, *Comput. Phys. Commun.* **181**, 1077 (2010), [arXiv:0909.2863](#). [p 5]
- [96] F. Staub, “Automatic Calculation of supersymmetric Renormalization Group Equations and Self Energies”, *Comput. Phys. Commun.* **182**, 808 (2011), [arXiv:1002.0840](#). [p 5]

- [97] F. Staub, “SARAH 3.2: Dirac Gauginos, UFO output, and more”, [Comput. Phys. Commun.](#) **184**, 1792 (2013), [arXiv:1207.0906](#). [p 5]
- [98] F. Staub, “SARAH 4: A tool for (not only SUSY) model builders”, [Comput. Phys. Commun.](#) **185**, 1773 (2014), [arXiv:1309.7223](#). [p 5]
- [99] S. P. Martin, “Two loop scalar self energies in a general renormalizable theory at leading order in gauge couplings”, [Phys. Rev.](#) **D70**, 016005 (2004), [arXiv:hep-ph/0312092](#). [pp 6, 8]
- [100] F. Domingo, S. Paßehr, “Towards Higgs masses and decay widths satisfying the symmetries in the (N)MSSM”, [Eur. Phys. J. C](#) **80**, 1124 (2020), [arXiv:2007.11010](#). [p 8]
- [101] U. Ellwanger, C. Hugonie, A. Teixeira, “The Next-to-Minimal Supersymmetric Standard Model”, [Phys. Rept.](#) **496**, 1 (2010), [arXiv:0910.1785](#). [p 15]
- [102] G. G. Ross, K. Schmidt-Hoberg, F. Staub, “The Generalised NMSSM at One Loop: Fine Tuning and Phenomenology”, [JHEP](#) **08**, 074 (2012), [arXiv:1205.1509](#). [p 15]