

DEUTSCHES ELEKTRONEN-SYNCHROTRON
Ein Forschungszentrum der Helmholtz-Gemeinschaft



DESY 21-092
DO-TH 21/20
TTP 21-023
SAGEX-21-11
RISC Report Series 21-14
arXiv:2107.09350
July 2021

New 2- und 3-Loop Heavy Flavor Corrections to Unpolarized and Polarized Deep-Inelastic Scattering

J. Ablinger, C. Schneider

RISC, Johannes Kepler Universität Linz, Austria

J. Blümlein, A. De Freitas, M. Saragnese

Deutsches Elektronen-Synchrotron DESY, Zeuthen

K. Schönwald

Institut für Theoretische Teilchenphysik, Karlsruher Institut für Technologie (KIT), Karlsruhe

ISSN 0418-9833

PLATANENALLEE 6 – 15738 ZEUTHEN

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your reports and preprints are promptly included in the
HEP literature database
send them to (if possible by air mail):

DESY Zentralbibliothek Notkestraße 85 22607 Hamburg Germany	DESY Bibliothek Platanenallee 6 15738 Zeuthen Germany
---	---

DESY 21–092, DO–TH 21/20, TTP 21–023, SAGEX–21–11, RISC Report Series 21–14,
arXiv: 2107.09350 [hep-ph]

New 2– and 3–loop heavy flavor corrections to unpolarized and polarized deep-inelastic scattering¹

J. Ablinger¹, J. Blümlein^{2*}, A. De Freitas², M. Saragnese², C. Schneider¹, K. Schönwald³,

1 Johannes Kepler University Linz, Research Institute for Symbolic Computation (RISC),
Altenberger Straße 69, A-4040, Linz, Austria

2 Deutsches Elektronen-Synchrotron DESY,
Platanenallee 6, D-15738 Zeuthen, Germany

3 Institut für Theoretische Teilchenphysik Campus Süd,
Karlsruher Institut für Technologie (KIT), D-76128 Karlsruhe, Germany

* Johannes.Blumlein@desy.de

July 21, 2021



*Proceedings for the XXVIII International Workshop
on Deep-Inelastic Scattering and Related Subjects,
Stony Brook University, New York, USA, 12-16 April 2021
doi:10.21468/SciPostPhysProc.?*

Abstract

A survey is given on the new 2– and 3–loop results for the heavy flavor contributions to deep–inelastic scattering in the unpolarized and the polarized case. We also discuss related new mathematical aspects applied in these calculations.

Contents

1	Introduction	2
2	Status of the massive OME calculations	2
3	Calculation of the 3–loop OMEs	3
4	Single mass contributions	5
5	Two-mass contributions	6
6	Conclusions	7
	References	8

¹Also contribution to the Proceedings of RADCOR 2021, Tallahassee, FL, May 2021.

1 Introduction

The scaling violations of deep–inelastic structure functions provide a precise way to measure the strong coupling constant $\alpha_s(M_Z)$ [1]. This requires to calculate their Q^2 dependence both due to the massless and massive contributions at highest precision. Moreover, the structure functions also allow the precise measurement of the charm quark mass, m_c , [2]. At future facilities, like the EIC [3] or the LHeC [4], operating at high luminosity, unpolarized and polarized structure functions can be measured at high precision to supplement and extend the present deep–inelastic world data. The massless QCD corrections are available to 3–loop order [5–13]. In the case of massive corrections, analytic results at the 3–loop level can currently only be obtained in the asymptotic approximation, $Q^2 \gg m_q^2$, with Q^2 the virtuality of the process and m_q the heavy quark mass, cf. [14], which, however, are accurate to $\sim 1\%$ for the structure function $F_2(x, Q^2)$ in the heavy quark region of smaller values of x for $Q^2/m_q^2 \gtrsim 10$ already. This region should be chosen also to avoid higher twist effects, requiring at least the cuts $W^2 > 15 \text{ GeV}^2, Q^2 > 10 \text{ GeV}^2$, [15].

After having obtained a series of Mellin moments for massive 3–loop operator matrix elements (OMEs) in 2009 [16] the systematic calculation of the different OMEs contributing in the unpolarized and polarized case for the massive Wilson coefficients and the OMEs contributing to the matching conditions in the variable flavor number scheme (VFNS) has been started for the single mass [7, 17–30] and the two–mass contributions [31–34].

At 2–loop order the unpolarized and polarized non–singlet and pure singlet contributions have been calculated for the full kinematic region and analytic results have been derived for the power corrections in Refs. [14, 35–38] to the structure functions F_2, F_L and g_1 .

In Section 2 we describe the status of the calculation of the massive 3–loop OMEs and discuss the calculation methods used in Section 3. Recent results in the single mass and two–mass cases are reported in Sections 4 and 5 and Section 6 contains the conclusions.

2 Status of the massive OME calculations

In the leading twist approximation deep–inelastic structure functions have the representation

$$F_{2,L}(x, Q^2) = \sum_{i,q} \mathbb{C}_{2,L}^{(i)} \left(x, \frac{Q^2}{\mu^2}, \frac{m_q^2}{\mu^2} \right) \otimes f_{(i)}(x, \mu^2), \quad (1)$$

with \otimes the Mellin convolution. A corresponding relation holds for the polarized structure function $g_1(x, Q^2)$. Here $\mathbb{C}^{(i)}$ denotes the Wilson coefficient related to the parton density $f_{(i)}$, with μ the factorization scale and m_q the heavy quark masses $m_q = m_c, m_b$. x denotes the Bjorken variable. The Wilson coefficients can be decomposed into the massless, C , and massive contributions, H ,

$$\mathbb{C}_{2,L}^{(i)} \left(x, \frac{Q^2}{\mu^2}, \frac{m_q^2}{\mu^2} \right) = C_{2,L}^{(i)} \left(x, \frac{Q^2}{\mu^2} \right) + H_{2,L}^{(i)} \left(x, \frac{Q^2}{\mu^2}, \frac{m_q^2}{\mu^2} \right). \quad (2)$$

At large scales Q^2 the heavy flavor Wilson coefficients have the representation [14]

$$H_{2,L}^{(i)} \left(x, \frac{Q^2}{\mu^2}, \frac{m_q^2}{\mu^2} \right) = \sum_{j,q} C_{(j),2,L} \left(x, \frac{Q^2}{\mu^2} \right) \otimes A^{ij} \left(x, \frac{m_q^2}{\mu^2} \right). \quad (3)$$

The leading order (LO) heavy flavor corrections were calculated for neutral and charged current interactions in [39] and numerically at next-to-leading order (NLO) in [40]. Analytic

results at NLO have been obtained in complete form for the non-singlet and pure singlet corrections in [14, 35–37] and in the asymptotic case $Q^2 \gg m_q^2$ in Refs. [14, 23, 35, 41–43].

At 3-loop order at present only results in the asymptotic case have been calculated. In the single mass case the unpolarized OMEs for all OMEs $\propto N_F$ and the complete results for $A_{qq,Q}^{\text{NS}}$, $A_{qg,Q}$, $A_{qq,Q}^{\text{PS}}$, A_{Qq}^{PS} , $A_{gq,Q}$ ² have been calculated in Refs. [7, 17, 18, 21, 44] and those $\propto T_F^2$ for $A_{gg,Q}$ in [19]. All logarithmic corrections were computed in [22]. For the pure N_F contributions and both in the non-singlet case and for $A_{gq,Q}$ the OMEs can be expressed by harmonic sums [45] or harmonic polylogarithms [46] only. In the pure singlet case also generalized harmonic sums [47, 48] contribute. In z space harmonic polylogarithms of argument z do not span the result, unless one allows also for the argument $1 - 2z$. For the T_F^2 terms of $A_{gg,Q}$ also nested finite binomial sums contribute, leading to root-valued iterated integrals [49]. Rather involved binomial structures also occur in the case of A_{Qg} [50]. The OME A_{Qg} also contains iterative non-iterative integrals [51], containing complete elliptic integrals. The first order factorizing contributions to A_{Qg} have been calculated in [27]. Phenomenological predictions in the non-singlet case have been given for the the structure functions $x F_3$ and $F_L^{W^+ - W^-}(x, Q^2)$ and $F_2^{W^+ - W^-}(x, Q^2)$ in [24, 26].

In the polarized case single mass contributions have been computed for the OMEs $A_{qq,Q}^{\text{NS}}$, $A_{qg,Q}$, $A_{qq,Q}^{\text{PS}}$, A_{Qq}^{PS} and $A_{gq,Q}$ in Refs. [18, 28–30] and for all logarithmic contributions in [29]. Here the same mathematical structures as in the unpolarized case contribute. Phenomenological predictions in the non-singlet case for $g_1(x, Q^2)$ were given in [25].

The two-mass corrections contributing from 3-loop order onward were calculated in the unpolarized case for $A_{qq,Q}^{\text{NS}}$, $A_{qg,Q}$ in [31], $A_{gg,Q}$ [32] and A_{Qq}^{PS} [31] and analogously in the polarized case [30, 33, 34, 34]. The analytic results in z space can be expressed by iterative integrals over root-valued letters, also parameterized by the mass ratio m_c^2/m_b^2 .

Because of the missing hierarchy between m_c^2 and m_b^2 , one has to decouple both these heavy quark effects together in a variable flavor number scheme [42], generalizing the single-mass variable flavor number scheme [31, 52].

Further calculations concern the missing terms for the OMEs $A_{gg,Q}$ in the single mass case and for A_{Qg} in the single and two-mass case. For A_{Qg} there is a first phenomenological representation in [53] based on only five Mellin moments calculated by us in Ref. [16].

3 Calculation of the 3-loop OMEs

A survey on the calculation methods for the 3-loop massive OMEs has been given in Ref. [54]. Massive OMEs contain, beyond the usual Feynman rules, those for the twist-2 local operators, cf. [16]. To apply integration-by-parts (IBP) techniques [55] one needs to resum these operators into propagators [56]. In the topological simple cases we perform the calculation of the 3-loop integrals directly, using hypergeometric techniques [57]. More complex integrals are calculated using the method of first order factorizing differential and difference equations [50, 58]. From the IBP reductions one obtains systems of differential equations which can be mapped into systems of difference equations, allowing to calculate a large number of Mellin moments for the master integrals and the OMEs by using the method of arbitrary large moments [59]. Having generated a sufficient number of moments the method of guessing [60, 61] allows to find the difference equations for the respective color and zeta factors

²In the non-singlet case we have also calculated the OMEs for transversity.




Figure 1: The ratios of the power expanded structure function to the complete structure function, $R_{2,q}^{(1)}$ (left) and $R_{L,q}^{(1)}$ (right), as a function of $\chi = 1/\kappa = Q^2/m_q^2$ for different values of z gradually improved with κ suppressed terms. Dotted lines: asymptotic result; dashed lines: $O(m_q^2/Q^2)$ improved; solid lines : $O((m_q^2/Q^2)^2)$ improved. From Ref. [37].

of the different OMEs. The number of these moments in the case of $A_{Qg}^{(3)}$ is very large. Already for the T_F^2 terms one needs ~ 7500 moments [27].

If the difference equations obtained factorize to first order the difference ring techniques [62] implemented in the package Sigma [63] are sufficient to find the final solution in Mellin N space. The z space solutions can be obtained by the techniques implemented in the package HarmonicSums [45, 46, 48, 49, 64] in terms of iterated integrals over certain alphabets, which are found algorithmically. All massive 3-loop OMEs except those in the single and two-mass case of $A_{Qg}^{(3)}$ belong to this class and have been solved by now.

In the following we describe some recent results in calculating single and two-mass contributions at the 2- and 3-loop level.

4 Single mass contributions

We consider first the successive inclusion of power corrections in m_q^2/Q^2 to the asymptotic result for the heavy flavor contributions to the structure functions $F_2(x, Q^2)$ and $F_L(a, Q^2)$ at NLO. The power corrections are of clear importance for F_L , since the pure asymptotic terms describe the heavy flavor corrections for $Q^2/m_q^2 \gtrsim 800$ only. Also in the case of F_2 for larger values of z an improved description is obtained. The results for the pure-singlet contribution to the polarized structure function $g_1(x, Q^2)$ [38] are similar. The inclusion of the power corrections in the pure-singlet case in the unexpanded expressions leads to new iterative integrals over a corresponding alphabet, cf. [37, 38]. Expanding in the ratio m_q^2/Q^2 leads to harmonic polylogarithms again. The situation is simpler in the non-singlet case, where classical polylogarithms with more complicated arguments suffice in the general case, cf. e.g. [36]. Note that the corresponding Wilson coefficients are not the ones given in [14], but need an extension.

The most important calculation for the future consist in the analytic computation of the OME $A_{Qg}^{(3)}$. The terms $\propto N_F$ were calculated in [17]. Based on 1000 Mellin moments all contributions, but the pure rational and ζ_3 terms were calculated, since these can be obtained from difference equations which are factorizing at first order, which is not the case for the former terms, [27]. Based on 8000 moments we obtained the difference equations for all missing terms $\propto T_F^2$. In the T_F -case one will need more moments to find the corresponding difference equations. In the T_F^2 case the difference equations have the following characteristics for degree d and order o :

$$\begin{aligned} T_F^2 C_A & : (d; o) = (1407; 46) \\ T_F^2 C_A \zeta_3 & : (d; o) = (323; 24) \\ T_F^2 C_F & : (d; o) = (654; 27) \\ T_F^2 C_F \zeta_3 & : (d; o) = (283; 14). \end{aligned}$$

The first difference equation is more voluminous than the largest occurring in guessing the largest contribution to the 3-loop massless Wilson coefficient in Ref. [61]. For the 3-loop massive form factor, [65], one difference equation of $(d, o) = (1324; 55)$ has been obtained.

The separate analysis of the difference equations for the rational and $\zeta_3 T_F^2$ cases showed, that the associated differential equations develop exponential singularities in the region $z \in [0, 1]$, although the complete solution is regular. Indeed, the differential equation for the purely rational term develops a ζ_3 factor asymptotically such, that both the singularities cancel, which requires to deal with both difference equations at once, despite the fact, that one can establish them only separately.

Starting from the difference equations in Mellin N space one may find Laurent series solutions of the associated differential equations around $z_0 = 1$ up to a finite upper power in z . This expansion is possible also around any other regular point $z_0 \in [0, 1]$, [66]. All these expansions have a finite convergence radius and several expansions are necessary to map out the interval $z \in [0, 1]$ in terms of overlapping expansions. One obtains an approximate analytic solution, containing high-precision numerical constants in part, in this way, which may be tuned to any accuracy. Given the fact that also all the other special functions need numerical representations, the numerical representation in this case is already obtained. The generation of a high number of moments for the T_F terms is underway.

In Figure 2 we summarize the different contributions to the currently known charm quark QCD corrections up to 3-loop order to the structure function F_2 at the scale $Q^2 = 100 \text{ GeV}^2$.




Figure 2: The different single mass heavy flavor contributions to the structure function F_2 at $Q^2 = 100\text{GeV}^2$. From [67].

5 Two-mass contributions

Irreducible two-mass contributions emerge first at 3-loop order and lead to a change of the variable flavor number scheme [31]. Reducible contributions imply two-mass contributions already at NLO [52]. In Figure 3 we illustrate the 2-mass effects on the singlet and bottom quark contribution. In the singlet case the effect amounts to 1% and in the case of the b -quark distribution of 4–5%.




Figure 3: The ratio of the 2-mass contributions to the singlet parton distribution $\Sigma(x, Q$ in the on-shell scheme. Dash-dotted line: $Q^2 = 30 \text{ GeV}^2$; Dotted line: $Q^2 = 30 \text{ GeV}^2$; Dashed line: $Q^2 = 100 \text{ GeV}^2$; Dash-dotted line: $Q^2 = 1000 \text{ GeV}^2$; Full line: $Q^2 = 10000 \text{ GeV}^2$. For the PDFs the NNLO variant of ABMP16 with $N_f = 3$ flavors was used [68]. Form [52].

We also have calculated the next-to-next-to leading order (NNLO) heavy flavor corrections to the flavor non-singlet structure functions $F_2^{\text{NS}}(x, Q^2)$ and $g_1^{\text{NS}}(x, Q^2)$ in the case of scheme-invariant evolution [69] shown in Figure 4. Here one considers the evolution

$$F_2^{\text{NS}}(x, Q^2) = E_{\text{NS}}(x, Q^2, Q_0^2) \otimes F_2^{\text{NS}}(x, Q_0^2) \quad (4)$$

from a starting scale Q_0^2 to Q^2 , where $E_{\text{NS}}(x, Q^2, Q_0^2)$ denotes a scheme-invariant evolution operator and the input distribution function $F_2^{\text{NS}}(x, Q_0^2)$ is measured experimentally. The evolution operator also depends on the masses m_c and m_b .




Figure 4: Left: The relative contribution of the heavy flavor contributions due to c and b quarks to the structure function F_2^{NS} at N³LO; dashed lines: 100 GeV²; dashed-dotted lines: 1000 GeV²; dotted lines: 10000 GeV². Right: The same for the structure function xg_1^{NS} at N³LO. From [69].

These corrections amount to $O(1\%)$ and are important in future factorization-scheme invariant measurements from the scaling violations of these structure functions at high luminosity facilities like the EIC or the LHeC.

6 Conclusions

Most of the massive 3-loop OMEs have been calculated in the single and two-mass case. They contribute to the (two-mass) variable flavor number scheme and the heavy flavor Wilson coefficients in unpolarized and polarized deep-inelastic scattering in the asymptotic region $Q^2 \gg m_q^2$. All quantities which can be described by first order factorizing difference equations have been computed. For the remaining OMEs the determination of their difference equations is underway. A method has recently been developed that can solve these equations as well. In all the computations extensive use has been made of the methods of arbitrary high Mellin moments, the method of guessing, and of difference ring theory to solve the respective physical problems. In course of these computations a series of different function spaces has been found to perform different intermediary steps of the calculation and to find a minimal analytic representation of the final results.

At two-loop order analytic results have been derived for the non-singlet and pure-singlet Wilson coefficients in the whole kinematic region. The 2-mass corrections are quantitatively as important as the $O(T_F^2)$ contributions in the single mass case. The $O(T_F)$ contributions to the 3-loop anomalous dimensions have been calculated as by-product of the calculation of the massive OMEs and they agree with the results of previous calculations. The calculation methods have also being applied to higher order massive QED corrections for the initial state radiation to observables in e^+e^- annihilation [70].

Acknowledgment. This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 764850, SAGEX and from the Austrian Science Fund (FWF) grant SFB F50 (F5009-N15).

References

- [1] S. Bethke et al. Proceedings of the Workshop on Precision Measurements of α_s (2011), arXiv:1110.0016 [hep-ph];
S. Alekhin, J. Blümlein and S.O. Moch, Mod. Phys. Lett. A **31** (2016) no.25, 1630023;
S. Moch, S. Weinzierl, et al. *High precision fundamental constants at the TeV scale*, arXiv:1405.4781 [hep-ph].
- [2] S. Alekhin, J. Blümlein, K. Daum, K. Lipka and S. Moch, Phys. Lett. B **720** (2013) 172–176 [arXiv:1212.2355 [hep-ph]].
- [3] D. Boer, M. Diehl, R. Milner, R. Venugopalan, W. Vogelsang, D. Kaplan, H. Montgomery, S. Vidor, A. Accardi and E. C. Aschenauer, et al. *Gluons and the quark sea at high energies: Distributions, polarization, tomography*, [arXiv:1108.1713 [nucl-th]].
- [4] P. Agostini et al. [LHeC and FCC-he Study Group], [arXiv:2007.14491 [hep-ex]];
J.L. Abelleira Fernandez et al. [LHeC Study Group], J. Phys. G **39** (2012) 075001 [arXiv:1206.2913 [physics.acc-ph]].
- [5] S. Moch, J.A.M. Vermaseren and A. Vogt, Nucl. Phys. B **688** (2004) 101–134 [arXiv:hep-ph/0403192 [hep-ph]].
- [6] A. Vogt, S. Moch and J.A.M. Vermaseren, Nucl. Phys. B **691** (2004) 129–181 [arXiv:hep-ph/0404111 [hep-ph]].
- [7] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel and C. Schneider, Nucl. Phys. B **890** (2014) 48–151 [arXiv:1409.1135 [hep-ph]].
- [8] S. Moch, J.A.M. Vermaseren and A. Vogt, Nucl. Phys. B **889** (2014) 351–400 [arXiv:1409.5131 [hep-ph]].
- [9] S. Moch, J.A.M. Vermaseren and A. Vogt, Phys. Lett. B **748** (2015) 432–438 [arXiv:1506.04517 [hep-ph]].
- [10] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel and C. Schneider, Nucl. Phys. B **922** (2017), 1-40 [arXiv:1705.01508 [hep-ph]].
- [11] A. Behring, J. Blümlein, A. De Freitas, A. Goedicke, S. Klein, A. von Manteuffel, C. Schneider and K. Schönwald, Nucl. Phys. B **948** (2019), 114753 [arXiv:1908.03779 [hep-ph]].
- [12] J. Blümlein, P. Marquard, C. Schneider and K. Schönwald, *The three-loop unpolarized and polarized non-singlet anomalous dimensions from off shell operator matrix elements*, [arXiv:2107.06267 [hep-ph]].
- [13] J.A.M. Vermaseren, A. Vogt and S. Moch, Nucl. Phys. B **724** (2005) 3–182 [arXiv:hep-ph/0504242 [hep-ph]].
- [14] M. Buza, Y. Matiounine, J. Smith, R. Migneron and W.L. van Neerven, Nucl. Phys. B **472** (1996) 611–658 [arXiv:hep-ph/9601302].
- [15] S. Alekhin, J. Blümlein and S. Moch, Phys. Rev. D **86** (2012) 054009 [arXiv:1202.2281 [hep-ph]].

- [16] I. Bierenbaum, J. Blümlein and S. Klein, Nucl. Phys. B **820** (2009) 417–482 [arXiv:0904.3563 [hep-ph]].
J. Blümlein, S. Klein and B. Tödtli, Phys. Rev. D **80** (2009) 094010 [arXiv:0909.1547 [hep-ph]].
- [17] J. Ablinger, J. Blümlein, S. Klein, C. Schneider and F. Wißbrock, Nucl. Phys. B **844** (2011) 26–54 [arXiv:1008.3347 [hep-ph]].
- [18] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel, M. Round, C. Schneider and F. Wißbrock, Nucl. Phys. B **886** (2014) 733–823 [arXiv:1406.4654 [hep-ph]].
- [19] J. Ablinger, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel, M. Round and C. Schneider, Nucl. Phys. B **885** (2014) 280–317 [arXiv:1405.4259 [hep-ph]].
- [20] J. Ablinger et al., DESY 15–112.
- [21] J. Ablinger, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel, M. Round, C. Schneider and F. Wissbrock, Nucl. Phys. B **882** (2014) 263–288 [arXiv:1402.0359 [hep-ph]].
- [22] A. Behring, I. Bierenbaum, J. Blümlein, A. De Freitas, S. Klein and F. Wißbrock, Eur. Phys. J. C **74** (2014) no.9, 3033 [arXiv:1403.6356 [hep-ph]].
- [23] J. Blümlein, A. Hasselhuhn and T. Pföh, Nucl. Phys. B **881** (2014) 1–41 [arXiv:1401.4352 [hep-ph]].
- [24] A. Behring, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel and C. Schneider, Phys. Rev. D **92** (2015) no.11, 114005 [arXiv:1508.01449 [hep-ph]].
- [25] A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel and C. Schneider, Nucl. Phys. B **897** (2015) 612–644 [arXiv:1504.08217 [hep-ph]].
- [26] A. Behring, J. Blümlein, G. Falcioni, A. De Freitas, A. von Manteuffel and C. Schneider, Phys. Rev. D **94** (2016) no.11, 114006 [arXiv:1609.06255 [hep-ph]].
- [27] J. Blümlein, J. Ablinger, A. Behring, A. De Freitas, A. von Manteuffel, and C. Schneider, PoS (QCDEV2017) 031 [arXiv:1711.07957 [hep-ph]].
- [28] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel, C. Schneider and K. Schönwald, Nucl. Phys. B **953** (2020) 114945 [arXiv:1912.02536 [hep-ph]].
- [29] J. Blümlein, A. De Freitas, M. Saragnese, C. Schneider and K. Schönwald, [arXiv:2105.09572 [hep-ph]].
- [30] A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel, K. Schönwald and C. Schneider, Nucl. Phys. B **964** (2021), 115331 [arXiv:2101.05733 [hep-ph]].
- [31] J. Ablinger, J. Blümlein, A. De Freitas, C. Schneider and K. Schönwald, Nucl. Phys. B **927** (2018) 339–367 [arXiv:1711.06717 [hep-ph]].
- [32] J. Ablinger, J. Blümlein, A. De Freitas, A. Goedicke, C. Schneider and K. Schönwald, Nucl. Phys. B **932** (2018) 129–240 [arXiv:1804.02226 [hep-ph]].
- [33] J. Ablinger, J. Blümlein, A. De Freitas, A. Goedicke, M. Saragnese, C. Schneider and K. Schönwald, Nucl. Phys. B **955** (2020) 115059 [arXiv:2004.08916 [hep-ph]].

- [34] J. Ablinger, J. Blümlein, A. De Freitas, M. Saragnese, C. Schneider and K. Schönwald, Nucl. Phys. B **952** (2020) 114916 [arXiv:1911.11630 [hep-ph]].
- [35] M. Buza, Y. Matiounine, J. Smith and W.L. van Neerven, Nucl. Phys. B **485** (1997), 420–456 [arXiv:hep-ph/9608342 [hep-ph]].
- [36] J. Blümlein, G. Falcioni and A. De Freitas, Nucl. Phys. B **910** (2016) 568–617 [arXiv:1605.05541 [hep-ph]].
- [37] J. Blümlein, A. De Freitas, C.G. Raab and K. Schönwald, Nucl. Phys. B **945** (2019) 114659 [arXiv:1903.06155 [hep-ph]].
- [38] J. Blümlein, C. Raab and K. Schönwald, Nucl. Phys. B **948** (2019) 114736 [arXiv:1904.08911 [hep-ph]].
- [39] E. Witten, Nucl. Phys. B **104** (1976) 445–47;
J. Babcock, D.W. Sivers, and S. Wolfram, Phys. Rev. D **18** (1978) 162–181;
M.A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. B **136** (1978) 157–176;
J.P. Leveille and T.J. Weiler, Nucl. Phys. **B147** (1979) 147–173;
M. Glück and E. Reya, Phys. Lett. B **83** (1979) 98–102;
M. Glück, E. Hoffmann, and E. Reya, Z. Phys. C **13** (1982) 119–130;
M. Glück, S. Kretzer and E. Reya, Phys. Lett. B **380** (1996) 171–176 [Erratum: Phys. Lett. B **405** (1997), 391] [arXiv:hep-ph/9603304 [hep-ph]];
J. Blümlein, A. Hasselhuhn, P. Kovacikova and S. Moch, Phys. Lett. B **700** (2011) 294–304 [arXiv:1104.3449 [hep-ph]];
A.D. Watson, Z. Phys. C **12** (1982) 123–125;
W. Vogelsang, Z. Phys. C **50** (1991) 275–284.
- [40] E. Laenen, S. Riemersma, J. Smith, W.L. van Neerven, Nucl. Phys. **B392** (1993) 162–228; 229–250.
S. Riemersma, J. Smith, W. L. van Neerven, Phys. Lett. **B347** (1995) 143–151 [hep-ph/9411431];
Precise representations in Mellin space to $O(\alpha_s^2)$ were derived in : S.I. Alekhin and J. Blümlein, Phys. Lett. B **594** (2004) 299–307 [arXiv:hep-ph/0404034];
F. Hekhorn and M. Stratmann, Phys. Rev. D **98** (2018) no.1, 014018 [arXiv:1805.09026 [hep-ph]].
- [41] M. Buza and W. L. van Neerven, Nucl. Phys. B **500** (1997), 301-324 doi:10.1016/S0550-3213(97)00327-1 [arXiv:hep-ph/9702242 [hep-ph]].
- [42] M. Buza, Y. Matiounine, J. Smith and W. L. van Neerven, Eur. Phys. J. C **1** (1998) 301–320 [hep-ph/9612398].
- [43] I. Bierenbaum, J. Blümlein and S. Klein, Nucl. Phys. B **780** (2007) 40–75 [arXiv:hep-ph/0703285];
I. Bierenbaum, J. Blümlein, S. Klein and C. Schneider, Nucl. Phys. B **803** (2008) 1–41 [arXiv:0803.0273 [hep-ph]];
I. Bierenbaum, J. Blümlein and S. Klein, Phys. Lett. B **672** (2009) 401–406 [arXiv:0901.0669 [hep-ph]]; *Two-loop massive operator matrix elements for polarized and unpolarized deep-inelastic scattering*, [arXiv:0706.2738 [hep-ph]].
- [44] J. Blümlein, A. Hasselhuhn, S. Klein and C. Schneider, Nucl. Phys. B **866** (2013), 196–211 [arXiv:1205.4184 [hep-ph]].

- [45] J.A.M. Vermaseren, *Int. J. Mod. Phys. A* **14** (1999) 2037–2076 [hep-ph/9806280];
J. Blümlein and S. Kurth, *Phys. Rev. D* **60** (1999) 014018 [hep-ph/9810241].
- [46] E. Remiddi and J. A. M. Vermaseren, *Int. J. Mod. Phys. A* **15** (2000) 725–754 [hep-ph/9905237].
- [47] S. Moch, P. Uwer and S. Weinzierl, *J. Math. Phys.* **43** (2002) 3363–3386 [arXiv:hep-ph/0110083 [hep-ph]].
- [48] J. Ablinger, J. Blümlein and C. Schneider, *J. Math. Phys.* **54** (2013) 082301 [arXiv:1302.0378 [math-ph]].
- [49] J. Ablinger, J. Blümlein, C.G. Raab and C. Schneider, *J. Math. Phys.* **55** (2014) 112301 [arXiv:1407.1822 [hep-th]].
- [50] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel and C. Schneider, *Comput. Phys. Commun.* **202** (2016), 33–112 [arXiv:1509.08324 [hep-ph]].
- [51] J. Ablinger, J. Blümlein, A. De Freitas, M. van Hoeij, E. Imamoglu, C.G. Raab, C.S. Radu and C. Schneider, *J. Math. Phys.* **59** (2018) no.6, 062305 [arXiv:1706.01299 [hep-th]].
- [52] J. Blümlein, A. De Freitas, C. Schneider and K. Schönwald, *Phys. Lett. B* **782** (2018), 362–366 [arXiv:1804.03129 [hep-ph]].
- [53] H. Kawamura, N. A. Lo Presti, S. Moch and A. Vogt, *Nucl. Phys. B* **864** (2012) 399–468 [arXiv:1205.5727 [hep-ph]].
- [54] J. Blümlein, *Large scale analytic calculations in quantum field theories*, in: *Algorithmic Combinatorics: Enumerative Combinatorics, Special Functions and Computer Algebra*, Eds. V. Pillwein and C. Schneider, pp. 63–88, [arXiv:1905.02148 [hep-ph]].
- [55] A. von Manteuffel and C. Studerus, arXiv:1201.4330 [hep-ph];
C. Studerus, *Comput. Phys. Commun.* **181** (2010) 1293–1300 [arXiv:0912.2546 [physics.comp-ph]].
- [56] J. Ablinger, J. Blümlein, C. Raab, C. Schneider and F. Wißbrock, *Nucl. Phys. B* **885** (2014) 409–447 [arXiv:1403.1137 [hep-ph]].
- [57] W.N. Bailey, *Generalized Hypergeometric Series*, (Cambridge University Press, Cambridge, 1935);
L.J. Slater, *Generalized Hypergeometric Functions*, (Cambridge University Press, Cambridge, 1966);
P. Appell and J. Kampé de Fériet, *Fonctions Hypergéométriques et Hyperspériques, Polynomes D’ Hermite*, (Gauthier-Villars, Paris, 1926);
P. Appell, *Les Fonctions Hypergéométriques de Plusieurs Variables*, (Gauthier-Villars, Paris, 1925);
J. Kampé de Fériet, *La fonction hypergéométrique*, (Gauthier-Villars, Paris, 1937);
H. Exton, *Multiple Hypergeometric Functions and Applications*, (Ellis Horwood, Chichester, 1976).
H. Exton, *Handbook of Hypergeometric Integrals*, (Ellis Horwood, Chichester, 1978).
H.M. Srivastava and P.W. Karlsson, *Multiple Gaussian Hypergeometric Series*, (Ellis Horwood, Chichester, 1985).
- [58] J. Ablinger, J. Blümlein, P. Marquard, N. Rana and C. Schneider, *Nucl. Phys. B* **939** (2019) 253–291 [arXiv:1810.12261 [hep-ph]].

- [59] J. Blümlein and C. Schneider, Phys. Lett. B **771** (2017) 31–36 [arXiv:1701.04614 [hep-ph]].
- [60] M. Kauers, *Guessing Handbook*, JKU Linz, Technical Report RISC 09–07; M. Kauers, M. Jaroschek, and F. Johansson, in: *Computer Algebra and Polynomials*, Editors: J. Gutierrez, J. Schicho, Josef, M. Weimann, Eds.. Lecture Notes in Computer Science **8942** (Springer, Berlin, 2015) 105–125 [arXiv:1306.4263 [cs.SC]].
- [61] J. Blümlein, M. Kauers, S. Klein and C. Schneider, Comput. Phys. Commun. **180** (2009) 2143–2165 [arXiv:0902.4091 [hep-ph]].
- [62] M. Karr, J. ACM **28** (1981) 305–350;
M. Bronstein, J. Symbolic Comput. **29** (2000) no. 6, 841–877;
C. Schneider, *Symbolic Summation in Difference Fields*, Ph.D. Thesis RISC, Johannes Kepler University, Linz technical report 01–17 (2001);
C. Schneider, An. Univ. Timisoara Ser. Mat.-Inform. **42** (2004) 163–179;
C. Schneider, J. Differ. Equations Appl. **11** (2005) 799–821;
C. Schneider, Appl. Algebra Engrg. Comm. Comput. **16** (2005) 1–32;
C. Schneider, J. Algebra Appl. **6** (2007) 415–441;
C. Schneider, Clay Math. Proc. **12** (2010) 285–308 [arXiv:0904.2323 [cs.SC]];
C. Schneider, Ann. Comb. **14** (2010) 533–552 [arXiv:0808.2596];
C. Schneider, in: *Computer Algebra and Polynomials, Applications of Algebra and Number Theory*, J. Gutierrez, J. Schicho, M. Weimann (ed.), Lecture Notes in Computer Science (LNCS) 8942 (2015) 157–191 [arXiv:1307.7887 [cs.SC]];
C. Schneider, J. Symb. Comput. **72** (2016) 82–127 [arXiv:1408.2776 [cs.SC]].
C. Schneider, J. Symb. Comput. **80** (2017) 616–664 [arXiv:1603.04285 [cs.SC]]. S.A. Abramov, M. Bronstein, M. Petkovšek, C. Schneider, J. Symb. Comput. **107** (2021) 23–66 [arXiv:2005.04944 [cs.SC]].
- [63] C. Schneider, Sémin. Lothar. Combin. **56** (2007) 1–36 article B56b; Simplifying Multiple Sums in Difference Fields, in: *Computer Algebra in Quantum Field Theory: Integration, Summation and Special Functions Texts and Monographs in Symbolic Computation* eds. C. Schneider and J. Blümlein (Springer, Wien, 2013) 325–360 [arXiv:1304.4134 [cs.SC]].
- [64] J. Ablinger, J. Blümlein and C. Schneider, J. Phys. Conf. Ser. **523** (2014) 012060 [arXiv:1310.5645 [math-ph]];
J. Ablinger, PoS (LL2014) 019 [arXiv:1407.6180[cs.SC]]; *A Computer Algebra Toolbox for Harmonic Sums Related to Particle Physics*, Diploma Thesis, JKU Linz, 2009, arXiv:1011.1176[math-ph]; *Computer Algebra Algorithms for Special Functions in Particle Physics*, Ph.D. Thesis, Linz U. (2012) arXiv:1305.0687[math-ph]; PoS (LL2016) 067; Experimental Mathematics **26** (2017) [arXiv:1507.01703 [math.CO]]; PoS (RADCOR2017) 001 [arXiv:1801.01039 [cs.SC]]; arXiv:1902.11001 [math.CO]; PoS (LL2018) 063;
J. Ablinger, J. Blümlein and C. Schneider, J. Math. Phys. **52** (2011) 102301 [arXiv:1105.6063 [math-ph]];
J. Blümlein, Comput. Phys. Commun. **180** (2009) 2218–2249 [arXiv:0901.3106 [hep-ph]].
- [65] J. Blümlein, P. Marquard, N. Rana and C. Schneider, Nucl. Phys. B **949** (2019) 114751 [arXiv:1908.00357 [hep-ph]].
- [66] J. Ablinger, J. Blümlein, and C. Schneider, in preparation.

- [67] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel, M. Round, C. Schneider and F. Wißbrock, PoS (QCDEV2016) 052 [arXiv:1611.01104 [hep-ph]].
- [68] S. Alekhin, J. Blümlein, S. Moch and R. Placakyte, Phys. Rev. D **96** (2017) no.1, 014011 [arXiv:1701.05838 [hep-ph]].
- [69] J. Blümlein and M. Saragnese, *The N^3LO Scheme-invariant QCD Evolution of the Non-singlet Structure Functions $F_2^{NS}(x, Q^2)$ and $g_1^{NS}(x, Q^2)$* , [arXiv:2107.01293 [hep-ph]].
- [70] J. Blümlein, A. De Freitas, C. G. Raab and K. Schönwald, Phys. Lett. B **791** (2019) 206–209 [arXiv:1901.08018 [hep-ph]]; Phys. Lett. B **801** (2020) 135196 [arXiv:1910.05759 [hep-ph]]; Nucl. Phys. B **956** (2020) 115055 [arXiv:2003.14289 [hep-ph]]; J. Ablinger, J. Blümlein, A. De Freitas and K. Schönwald, Nucl. Phys. B **955** (2020) 115045 [arXiv:2004.04287 [hep-ph]]; J. Blümlein, A. De Freitas and K. Schönwald, Phys. Lett. B **816** (2021) 136250 [arXiv:2102.12237 [hep-ph]].