

**SEARCH FOR NEW FLAVOUR PRODUCTION AT PETRA**

JADE Collaboration

W. BARTEL, T. CANZLER<sup>1</sup>, D. CORDS, P. DITTMANN, R. EICHLER, R. FELST, D. HAIDT,  
S. KAWABATA, H. KREHBIEL, B. NAROSKA, L.H. O'NEILL, J. OLSSON, P. STEFFEN and W.L. YEN<sup>2</sup>  
*Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany*

E. ELSEN, M. HELM, A. PETERSEN, P. WARMING and G. WEBER  
*II. Institut für Experimentalphysik der Universität Hamburg, Germany*

H. DRUMM, J. HEINTZE, G. HEINZELMANN, R.D. HEUER, J. von KROGH, P. LENNERT,  
H. MATSUMURA, T. NOZAKI, H. RIESEBERG and A. WAGNER  
*Physikalisches Institut der Universität Heidelberg, Germany*

J. ALLISON, J. ARMITAGE, I. DUERDOTH, F. LOEBINGER, A. MACBETH, H. MILLS,  
P.G. MURPHY, H. PROSPER and K. STEPHENS  
*University of Manchester, England*

D.C. DARVILL, F. FOSTER, G. HUGHES and H. WRIEDT  
*University of Lancaster, England*

D. CLARKE, M.C. GODDARD, R. HEDGECOCK, R. MARSHALL and G.F. PEARCE  
*Rutherford Laboratory, Chilton, England*

M. IMORI, T. KOBAYASHI, S. KOMAMIYA, M. KOSHIBA, M. MINOWA, S. ORITO, A. SATO, T. SUDA<sup>3</sup>  
H. TAKEDA, Y. TOTSUKA, Y. WATANABE, S. YAMADA and C. YANAGISAWA<sup>4</sup>  
*Lab. of Int. Coll. on Elementary Particle Physics and Department of Physics, University of Tokyo, Japan*

Received 29 October 1979

The topological distribution of hadrons from the reaction  $e^+e^- \rightarrow$  multihadrons has been studied at PETRA energies between  $\sqrt{s} = 22$  and 31.6 GeV. No evidence is seen for spherical events which would be expected if massive particles bearing new flavours were produced.

In a previous paper [1] we reported on measurements of the cross section for the reaction  $e^+e^- \rightarrow$  multihadrons using the JADE detector at PETRA

energies between 22 and 31.6 GeV. The  $R$  values, i.e. the hadron cross section in units of the theoretical  $\mu^+\mu^-$  cross section, were found to be consistent with the production of the known quark flavours (u, d, s, c, b) only. The energy dependence of the  $R$  value does not show any significant discontinuity, so that there is no apparent indication for a threshold of new flavour

<sup>1</sup> Present address: Siemens AG, Munich, Germany.

<sup>2</sup> Present address: Purdue University, Indianapolis, USA.

<sup>3</sup> Present address: Cosmic Ray Lab., University of Tokyo, Japan.

<sup>4</sup> Partly supported by the Yamada Science Foundation.

production in the energy range in question. However, the existence of a sixth quark, *t* (top), which would be the doublet partner of the *b* quark, has been conjectured in the standard SU(2) × U(1) model of quarks and leptons [2]. The electric charge of the *t* quark is predicted to be 2/3 but its mass is essentially unknown.

In order to obtain a better sensitivity for new flavour detection, the topological distribution of final state hadrons has been studied as reported in this letter <sup>#1</sup> Heavy particles with new flavours are expected to decay predominantly into multihadron final states, which near threshold would yield a relatively wide angled distribution in contrast to the narrow jet-like distributions from known low-mass flavours. For the *t* quark the decay sequence *t* → *b* → *c* → *s*, which is expected to be favoured [5], would yield such wide distributions [8]. A comparison of the measured hadron distributions with the QCD predictions and evidence for three-jet events will be presented in a subsequent letter.

The JADE detector has a solid angle of 97% and 90% of the full solid angle for detecting charged particles and photons, respectively. The detector, the trigger conditions and the criteria for the selection of hadronic events have been described in ref. [1]. The final multihadron sample for this analysis consists of 21, 81, 198 and 49 events at  $\sqrt{s} = 22, 27.6, 30$  and 31.6 GeV, respectively, with a residual background estimated to be less than 3%.

For a detailed study of the event shape the sphericity tensor <sup>#2</sup> was constructed and diagonalized for each event:

$$T_{\alpha\beta} = \sum_i P_{i\alpha} P_{i\beta} / \sum_i P_i^2,$$

where  $P_{i\alpha}$  is the  $\alpha$ -component ( $\alpha = x, y, z$ ) of the momentum of the *i*th particle. The sum runs over all charged particles and shower energy clusters. The resulting eigenvalues  $Q_1, Q_2, Q_3$  ( $Q_1 < Q_2 < Q_3$ ) correspond to the principal axes of the momentum ellipsoid and satisfy the constraint  $Q_1 + Q_2 + Q_3 = 1$ . Each event is then represented by a point in a Dalitz plot as shown

<sup>#1</sup> Preliminary results were presented at the 1979 Intern. Symp. on Lepton and photon interactions at high energies (Fermilab National Accelerator Laboratory, Batavia, IL, USA).

<sup>#2</sup> Our definition of  $T_{\alpha\beta}$  is slightly different from but equivalent to the one proposed in ref. [3].




Fig. 1. The distribution of the eigenvalues  $Q_1, Q_2, Q_3$  of the sphericity tensor  $T$  described in the text.  $Q_1$  is plotted versus  $(Q_3 - Q_2)/\sqrt{3}$ . (a) Data from all energies combined and (b) model prediction at  $\sqrt{s} = 30$  GeV including top-meson ( $m_t = 14$  GeV) production. The cuts  $Q_1 > 0.075$  and sphericity  $> 0.55$  are indicated.

in fig. 1a. The perpendicular sides of the triangle are  $Q_1$  and  $(Q_3 - Q_2)/\sqrt{3}$  so that the hypotenuse is the sphericity

$$S = \frac{3}{2}(Q_1 + Q_2) \equiv \frac{3}{2} \frac{\sum_i p_{i\perp}^2}{\sum_i |p_i|^2},$$

where  $p_{i\perp}$  is the momentum component of particle *i* perpendicular to the principle three-axis.

The plot shown in fig. 1a contains all events from the various c.m. energies. No significant accumulation

of "spherical" events is observed in the upper part of the triangle.

In order to assess the significance of this observation we compare the data with model calculations. This model is based on virtual photon decays into a quark-antiquark pair  $q\bar{q}$  ( $q = u, d, c, b, t$ ) with a probability proportional to  $(e_q)^2$ , where  $e_q$  is the electric charge of the quark  $q$ . The primary quarks are then allowed to fragment into hadrons according to a cascade mechanism described in ref. [4]. For the  $u, d, s$  quarks the fragmentation function  $f(z) = 1 - a + 3a(1 - z)^2$  with  $a = 0.77$  was used, whereas  $f(z) = \text{const.}$  was used for the heavier quarks. Particles containing a  $t$  quark were assumed to decay [5] through the chain  $t \rightarrow b \rightarrow c \rightarrow s$ . To generate the final state particles we use the Monte Carlo program of ref. [6] where the emission of hard photons by the initial electron and positron was taken into account. The final particles were then followed through the detector, taking into account decays and multiple scattering, conversion of photons and nuclear interaction in the detector material. We then calculate the quantities actually measured such as drift times and pulse heights, and bin them according to the resolution of the apparatus. These data were afterwards processed by the same chain of pattern recognition as our real data.

Fig. 1b shows the results of this model calculation for a top meson mass of 14 GeV and a c.m. energy of 30 GeV. For quantitative comparison we list in table 1 the number of observed and expected events inside the region  $S > 0.55$  and  $Q_1 > 0.075$  for three different values of the top meson mass. The normalization is based on the accumulated luminosity and a quark production cross section of  $\sigma_{q\bar{q}} = 3e_q^2\sigma_{\mu^+\mu^-}$ . Monte Carlo simulation without top meson production yields no events in this region of the Dalitz plot.

Furthermore, if the top particles are allowed to decay according to a uniform invariant phase space distribution the same number of expected events is obtained to within 10%.

In conclusion, no evidence has been observed for the production of heavy mesons with new flavour. Our data are inconsistent with the production of top mesons with masses between 11 and 14 GeV provided  $\sigma_{t\bar{t}} \approx \frac{4}{3}\sigma_{\mu^+\mu^-}$ , although it should be remarked that the conclusion is based on the assumption of multihadron decays and a short life time of any hypothetical heavy particle. Similar deductions were made in ref. [7].

Table 1

The observed numbers of events with  $Q_1 > 0.075$  and sphericity  $> 0.55$  and the numbers expected from top-meson production for assumed top-meson masses of 8, 11 and 14 GeV.

	$m_t$ (GeV)	$\sqrt{s}$ (GeV)			
		22	27.7	30	31.6
Number of events	8	4	3	2	0
expected	11	—	15	22	4
Number of events observed	14	—	—	25	5
Number of events observed		0	1	1	1

We acknowledge the efforts of the PETRA machine group, who provided us with the opportunity of doing this experiment, and also the efforts of the technical support groups of the participating institutes in the construction and maintenance of our apparatus. This experiment was supported by the Bundesministerium für Forschung und Technologie, by the Education Ministry of Japan and by the U.K. Science Research Council through the Rutherford Laboratory. The visiting groups at DESY wish to thank the DESY directorate for their hospitality.

### References

- [1] JADE Collaboration, W. Bartel et al., Phys. Lett. 88B (1979) 171.
- [2] M. Kobayashi and K. Maskawa, Prog. Theor. Phys. 49 (1973) 652.
- [3] J.D. Bjorken and S.J. Brodsky, Phys. Rev. D1 (1970) 1416; C. Hanson et al., Phys. Rev. Lett. 35 (1975) 1609.
- [4] R.D. Field and R.P. Feynman, Nucl. Phys. B136 (1978) 1; B. Anderson, G. Gustafson and C. Peterson, Nucl. Phys. B135 (1978) 273.
- [5] J. Ellis, M.K. Gaillard, D. Nanopoulos and S. Rudaz, Nucl. Phys. B131 (1977) 252.
- [6] T. Sjöstrand, Lund report LU TP 79-8, and references therein.
- [7] See contributions by the Mark J, Pluto and Tasso Collaborations to the 1979 Intern. Symp. on Lepton and photon interactions (Batavia, IL, USA); D.P. Barber et al., Phys. Lett. 85B (1979) 463; Ch. Berger et al., DESY 79/56.
- [8] A. Ali, J.G. Körner, G. Kramer and J. Willrodt, Z. Phys. C1 (1979) 203.