

# Heavy-quark and Quarkonia production in high-energy heavy-ion collisions

*André Mischke*<sup>†</sup>

ERC Starting Independent Research Group, Institute for Subatomic Physics, Utrecht University,  
Princetonplein 5, 3584 CC Utrecht, the Netherlands.

## Abstract

Relativistic heavy-ion collisions provide the unique conditions to produce and study a novel state of QCD matter, the Quark-Gluon Plasma, in the laboratory. Heavy-quarks are a powerful probe for the detailed investigation of the QGP properties. In this paper we review recent results from RHIC on open heavy-flavor and Quarkonia production and their interaction with the QCD matter on the partonic level.

## 1 Introduction

High-energy nucleus-nucleus collisions at the Relativistic Heavy Ion Collider (RHIC) at Brookhaven National Laboratory allow exploring strongly interacting matter at very high temperatures and energy density. QCD matter at these conditions is expected to form a system of deconfined quarks and gluons, the so-called Quark-Gluon Plasma (QGP), if the critical energy density ( $\epsilon_c \sim 0.7 \text{ GeV/fm}^3$ ) is exceeded. The goal of relativistic heavy-ion physics is to study the properties of the QGP under laboratory controlled conditions [1, 2].

The results from RHIC have given evidence that the nuclear matter created in such collisions exhibits properties consistent with the QGP formation [3]. In particular, measurements of the momentum distribution of emitted particles and comparison with hydro-dynamic model calculations have shown that the outward steaming particles move collectively, with the patterns arising from variations of pressure gradients early after the collision. This phenomenon, called elliptic flow, is analogous to the properties of fluid motion. The flow results suggest that colour degrees of freedom carried by quarks and gluons are present in the produced medium, which flow with negligible shear viscosity. Thus, the QCD matter at RHIC behaves like a perfect liquid. Moreover, it has been found that the matter remaining in the collision zone is extremely opaque to the passage of partons from hard scattering processes in the initial state of the collisions. These traversing partons are believed to lose energy via gluon Bremsstrahlung in the medium before fragmenting into hadrons.

A detailed and quantitative understanding of the parton energy loss in the medium is one of the intriguing issues which need to be addressed. The study of heavy-flavor (charm, bottom) production in heavy-ion collisions provides key tests of the parton energy loss mechanisms and offers important information on the properties of the produced medium [4]. Due to their large mass ( $m > 1 \text{ GeV}/c^2$ ), heavy quarks are expected to be primarily produced in the early stage of the collision and, therefore, probe the complete space-time evolution of the medium. Theoretical

---

<sup>†</sup>E-mail: a.mischke@uu.nl




Fig. 1: (a) Nuclear modification factor  $R_{AA}$  (averaged above  $p_T > 3$  GeV/c) of heavy-flavor decay electrons as a function of collision centrality (quantified in  $N_{part}$ ) in Au+Au and minimum bias Cu+Cu collisions at  $\sqrt{s_{NN}} = 200$  GeV. (b) Relative bottom contribution to the total yield of heavy-flavor decay electrons derived from  $e-D^0$  and  $e$ -hadron azimuthal angular correlations, compared to the uncertainty band from a FONLL calculation.

models predicted that heavy quarks should experience smaller energy loss than light quarks while propagating through the QCD medium due to the suppression of small angle gluon radiation, the so-called *dead-cone effect* [5, 6].

## 2 Probing the QCD medium with heavy quarks

Nuclear effects are typically quantified using the nuclear modification factor  $R_{AA}$  where the particle yield in Au+Au collisions is divided by the yield in  $pp$  reactions scaled by the number of binary collisions.  $R_{AA} = 1$  would indicate that no nuclear effects, such as Cronin effect, shadowing or gluon saturation, are present and that nucleus-nucleus collisions can be considered as a incoherent superposition of nucleon-nucleon interactions. Charm and bottom quarks can be identified by assuming that isolated electrons in the event stem from semi-leptonic decays of heavy-quark mesons. At high transverse momentum ( $p_T$ ), this mechanism of electron production is dominant enough to reliably subtract other sources of electrons like conversions from photons and  $\pi^0$  Dalitz decays. Fig. 1 (a) shows the average  $R_{AA}$  for heavy-flavor decay electrons in Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV as a function of participating nucleons ( $N_{part}$ ) measured by the STAR and PHENIX experiments [7, 8]. The data are consistent with each other and the  $R_{AA}$  shows an increasing suppression from peripheral to central Au+Au collisions. The minimum bias Cu+Cu data fit into this systematics. The strong suppression for the most central Au+Au collisions indicates an unexpectedly large energy loss of heavy quarks in the medium in contradiction to expectations from the dead-cone effect. Surprisingly, the  $R_{AA}$  of 0.2 is similar to the one observed for light-quark hadrons. Current models with reasonable model parameters overpredict the observed suppression [7, 8]. The data is described reasonably well if the bottom contribution to the electrons is assumed to be small. Therefore, the observed discrepancy could




Fig. 2: (Color online) (a) The centrality dependence of the nuclear modification factor  $R_{AA}$  of  $J/\psi$ , measured for different collisions energies and rapidity regions. For Au+Au collisions, the forward rapidity region (full circles) shows more suppression than the mid-rapidity region (open symbols). (b)  $R_{AA}$  of  $J/\psi$  in the 20 and 60% most central Cu+Cu collisions at  $\sqrt{s_{NN}} = 200$  GeV. The boxes in the right indicate the normalization uncertainty. The horizontal line represents a fit to the data in the  $p_T$  range 5-10 GeV/c. The curves are model predictions from the *Two Component Approach* model.

indicate that the  $B$  dominance over  $D$  mesons starts at higher  $p_T$  than expected. A possible scenario for  $B$  meson suppression invokes collisional dissociation in the medium [9].

The measurement of the relative charm and bottom contributions to the heavy-flavor decay electrons (also called non-photonic electrons) is essential for the interpretation of the electron spectra and nuclear modification factor. Azimuthal angular correlations between non-photonic electrons and hadrons allow to identify the underlying production process [10]. The relative bottom contribution  $B/(B+D)$  to the non-photonic electrons is extracted from the  $e$ -hadron and  $e$ - $D^0$  azimuthal correlation distributions [11]. Figure 1(b) shows the  $B/(B+D)$  ratio together with predictions from calculations of heavy-flavor production in  $pp$  collisions at Fixed-Order plus Next-to-Leading Logarithm (FONLL) level [12]. These data provide convincing evidence that bottom contributes significantly ( $\sim 50\%$ ) to the non-photonic electron yields above  $p_T = 5$  GeV/c. Further studies have to show whether these results imply substantial suppression of bottom production at high  $p_T$  in the produced medium. An important step to answer this question will be the direct measurement of open charmed mesons and the identification of  $B$  mesons via displaced electrons using the detector upgrades of the STAR and PHENIX experiments.

### 3 Dissociation of quarkonium states in the hot and dense QCD medium

The dissociation of quarkonia due to color-screening in a QGP is a classic signature of deconfinement in relativistic heavy-ion collisions [13, 14], where the sequential suppression of the quarkonia states, such as  $\Upsilon$ ,  $\Upsilon'$  and  $\Upsilon''$ , depends on the temperature of the surrounding medium, thus providing a QCD thermometer.




Fig. 3: (a) The  $e^+e^-$  invariant mass distribution in  $pp$  (upper panel) and  $\text{Cu}+\text{Cu}$  collisions (lower panel) at  $\sqrt{s_{NN}} = 200 \text{ GeV}$ . The solid and dashed histograms represent the distribution of unlike and like-sign pair combinations, respectively. (b)  $J/\psi$ -hadron azimuthal angular correlations in  $pp$  collisions after background subtraction. The dashed curve shows the  $J/\psi$ -hadron contribution from  $B$  decays obtained from PYTHIA simulations.

### 3.1 $J/\psi$ measurements

Results from the PHENIX experiment have shown that the centrality dependence of the suppression of the  $J/\psi$  yield in  $\sqrt{s_{NN}} = 200 \text{ GeV}$   $\text{Au}+\text{Au}$  collisions is similar to that observed at the CERN-SPS accelerator ( $\sqrt{s_{NN}} = 17.3 \text{ GeV}$ ) [16], even though the energy density reached in collisions at RHIC is about a factor of 2-3 higher (cf. Fig. 2 (a)). It has also been found that the yield in the forward rapidity region is more suppressed than the one at mid-rapidity. Theoretical prediction based on string theory application of AdS/CFT suggests that the effective  $J/\psi$  dissociation temperature is expected to decrease with  $p_T$  [15]. This conjecture is different from the predictions of more traditional screening models where the suppression due to screening vanishes towards higher  $p_T$ . Recent  $R_{AA}$  measurements for  $J/\psi$  in  $\text{Cu}+\text{Cu}$  collisions at  $\sqrt{s_{NN}} = 200 \text{ GeV}$  from the STAR [17] and PHENIX experiments [18] are compared in Fig. 2 (b). The  $R_{AA}$  is suppressed at low  $p_T$  (around  $1 \text{ GeV}/c$ ), and the data suggest that  $R_{AA}$  increases with increasing  $p_T$  and reaches unity around  $5 \text{ GeV}/c$ , although the large errors currently preclude strong conclusions. However, this result is in contradiction with expectations from the *Two Component Approach* model [19], which predicts a suppression at high  $p_T$ . These results could indicate that other  $J/\psi$  production mechanisms that counter the suppression such as recombination and formation-time effects might play an increasingly dominant role at higher  $p_T$ .

The relative large signal-to-background ratio of the  $J/\psi$  in  $pp$  collisions (cf. Fig. 3(a)) makes it possible studying  $J/\psi$ -hadron correlations at high trigger- $p_T$ , which provide important information on the underlying  $J/\psi$  production mechanisms. Figure 3(b) illustrates the azimuthal angular correlation distribution of  $J/\psi$  and charged hadrons. Notably, no significant correlation yield on the near-side ( $\Delta\phi \sim 0 \text{ rad}$ ) is observed, which is not in line with earlier results from di-hadron correlation measurements [3]. Since corresponding PYTHIA simulations (also depicted in Fig. 3(b)) show a strong near-side correlation peak from  $J/\psi$  from  $B$  decays ( $B \rightarrow J/\psi + X$ ),

the experimental results can be used to estimate the  $B$  feed-down contribution to the inclusive  $J/\psi$  yield at  $p_T > 5$  GeV/ $c$ . It was found to be  $17 \pm 3\%$  in the studied  $p_T$  range [17].

### 3.2 First $\Upsilon$ measurements in nuclear collisions

The golden decay channel for the  $\Upsilon$  reconstruction is the decay into electron pairs  $\Upsilon \rightarrow e^+e^-$ . The STAR detector with its large acceptance ( $|\eta| < 1$  and  $0 < \phi < 2\pi$ ) and excellent trigger capabilities combined with a very good electron identification is very good suited for  $\Upsilon$  measurements in nuclear collisions. The first preliminary measurements of the  $\Upsilon$  invariant mass in Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV are presented in [20] and shows a significant  $\Upsilon$  signal. The  $\Upsilon$  production cross-section is also measured in  $pp$  collisions and is found to be  $BR_{ee} \times \frac{d\sigma}{dy}_{y=0} = 91 \pm 28(stat.) \pm 22(sys.)$ pb. The measurement follows the world data trend and shows, within uncertainties, very good agreement with NLO calculations [21]. The analysis of the full  $pp$  and Au+Au data-sets will allow to extract the  $\Upsilon$  nuclear modification factor in the near future.

## 4 Summary

The observed strong suppression of the yield of heavy-flavor decay electrons at high  $p_T$  in central Au+Au collisions together with the measurement of the azimuthal angular correlation of electrons and hadrons in  $pp$  collisions imply that  $B$  production is stronger suppressed in nuclear collisions than expected. The nuclear modification factor ( $R_{AA}$ ) of  $J/\psi$  in Cu+Cu collisions increases from low to high  $p_T$  and reaches unity for  $p_T > 5$  GeV/ $c$ . This result is about  $2\sigma$  above the  $R_{AA}$  at low  $p_T$  ( $< 4$  GeV/ $c$ ) and is consistent with no  $J/\psi$  suppression. First RHIC results on the  $\Upsilon$  production in nuclear collisions are promising and show that the suppression measurements will be possible in the near future.

## Acknowledgments

The author would like to thank the organizers for the intellectual stimulating atmosphere during the conference. The European Research Council has provided financial support under the European Community's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement no 210223. This work was supported in part by a Veni grant from the Netherlands Organisation for Scientific Research (project number 680-47-109).

## References

- [1] P. Braun-Munzinger and J. Stachel, *Nature* **448**, 302 (2007).
- [2] B. Muller, *Rep. Prog. Phys.* **58**, 611 (1995) and references therein.
- [3] I. Arsene *et al.*, *Nucl. Phys.* **A757**, 1 (2005).
- [4] A.D. Frawley, T. Ullrich and R. Vogt, *Phys. Rep.* **462**, 125 (2008).
- [5] Y.L. Dokshitzer and D.E. Kharzeev, *Phys. Lett.* **B519**, 199 (2001).
- [6] M. Djordjevic, M. Gyulassy and S. Wicks, *Phys. Rev. Lett.* **94**, 112301 (2005).
- [7] B.I. Abelev *et al.*, *Phys. Rev. Lett.* **98**, 192301 (2007).
- [8] A. Adare *et al.*, *Phys. Rev. Lett.* **98**, 172301 (2007); S.S. Adler *et al.*, *Phys. Rev. Lett.* **96**, 032301 (2006).

- [9] A. Adil and I. Vitev, Phys. Lett. **B649**, 139 (2007).
- [10] A. Mischke, Phys. Lett. **B671**, 361 (2009).
- [11] A. Mischke *et al.*, J. Phys. G: Nucl. Part. Phys. **35**, 104117 (2008).
- [12] M. Cacciari, P. Nason, and R. Vogt, Phys. Rev. Lett. **95**, 122001 (2005).
- [13] T. Matsui and H. Satz, Phys. Lett. **B178**, 416 (1986).
- [14] F. Karsch, D. Kharzeev and H. Satz, Phys. Lett. **B637**, 75 (2006) and arXiv:hep-ph/0512239 (2005).
- [15] H. Liu, K. Rajagopal and U.A.Wiedemann, Phys. Rev. Lett. **98** (2007) 182301.
- [16] A. Adare *et al.*, Phys. Rev. Lett. **98**, 232301 (2007).
- [17] Z. Xu, T. Ullrich *et al.*, arXiv:0809.2288 (2008).
- [18] A. Adare *et al.*, Phys. Rev. Lett. **101**, 122301 (2008).
- [19] X. Zhao and R. Rapp, Phys. Lett. **B664**, 253 (2008), private communication for Cu+Cu predictions.
- [20] D. Das *et al.*, J. Phys. G: Nucl. Part. Phys. **35**, 104153 (2008).
- [21] P. Djawotho *et al.*, J. Phys. G: Nucl. Part. Phys. **34**, S947 (2007).