

Status of the BMV experiment

Franck Bielsa¹, Rémy Battesti¹, Mathilde Fouché^{2,3}, Paul Berceau¹, Cécile Robilliard^{2,3}, Gilles Bailly^{2,3}, Sébastien Batut¹, Julien Mauchain¹, Marc Nardone¹, Oliver Portugall¹, Carlo Rizzo^{2,3}

¹Laboratoire National des Champs Magnétiques Intenses (UPR 3228, CNRS-INSU-UJF-UPS), F-31400 Toulouse Cedex, France.

²Université de Toulouse, UPS, Laboratoire Collisions Agrégats Réactivité, IRSAMC, F-31062 Toulouse, France.

³CNRS, UMR 5589, F-31062 Toulouse, France.

DOI: http://dx.doi.org/10.3204/DESY-PROC-2009-05/rizzo_carlo

In this contribution we present the status of the BMV experiment whose goal is to measure the vacuum magnetic birefringence.

1 Introduction

In this contribution we present the status of the BMV (Biréfringence Magnétique du Vide) experiment [1] whose goal is to measure the vacuum magnetic birefringence i.e. the birefringence induced in vacuum by the presence of an intense magnetic field. Linearly polarized light passing through a region where a magnetic field B perpendicular to the direction of propagation is present will acquire an ellipticity Ψ because of the vacuum magnetic birefringence. Ψ can be written as $\Psi = \frac{\pi}{\lambda} \Delta n B^2 L$, where λ is the light wavelength, Δn is the difference between the index of refraction of the light polarized parallel to the magnetic field and the index of refraction of the light polarized perpendicular to the magnetic field and L is the length of the magnetic field region. When B is given in Tesla Δn is expected to be about 4×10^{-24} . It is clear looking to the previous equation that the critical parameter for experiments looking for vacuum magnetic birefringence is $B^2 L$. Our choice, since the beginning in 2001, has been to reach a $B^2 L$ approaching 1000 T²m having a B as high as possible with an L as small as possible to set-up a table-top optical experiment which, we believe, has the best chances of success.

The value of Δn has been first calculated in the seventies [2] starting from the Heisenberg-Euler Lagrangian established in 1935 [3] to describe the photon-photon interactions in the framework of Quantum Electrodynamics based on Dirac's model of the vacuum. At the lower orders in α , the fine structure constant, Δn can be written as

$$\Delta n = \frac{2}{15} \frac{\alpha^2 \hbar^3}{m_e^4 c^5} \left(1 + \frac{25}{4\pi} \alpha\right) \frac{B^2}{\mu_0} \quad (1)$$

where \hbar is the Planck constant over 2π , m_e is the electron mass, c the speed of light in vacuum, and μ_0 is the magnetic constant. The α^2 term is given in ref. [2]. The α^3 term has been first reported in ref. [4] and it corresponds to the lower order radiative correction to the main term. Its value is about 1.5% of the first order term.

Using the 2006 CODATA recommended values [5] for the fundamental constants, equation (1) gives $\Delta n = (4.031699 \pm 0.000005) \times 10^{-24} B(\text{T})^2$.

As we see, the error due to the knowledge of fundamental constants is negligible compared with the error coming from the fact that only first order QED radiative correction has been calculated. QED α^4 radiative correction should affect the fourth digit and the QED α^5 radiative correction the sixth digit. Thus measurement of Δn up to a precision of a few ppm remains a pure QED test.

It is known that an experiment designed to measure the vacuum magnetic birefringence can also give limits on the mass and the coupling constant of pseudoscalar particles that couples with two photons like axions or axionlike particles [6]. Using formulas given for example in ref. [6] on the ellipticity induced because of the existence of axionlike particles on polarized light propagating in the presence of a transverse magnetic field, one can infer the limits that can be given by an experiment like BMV [1]. If we suppose that the QED test has been successfully performed e.g. the Δn value given previously has been found experimentally with a precision of 10%, a coupling constant lower than about 5×10^7 GeV will be excluded for axionlike particles of a mass between 10^{-3} and 10^{-2} eV. A QED test of a precision of 1% i.e. at the level of the first radiative correction, will give a coupling constant limit of a few 10^8 GeV. Vacuum magnetic birefringence experiments are intrinsically different from astrophysics searches because they are terrestrial and all the experimental parameters are under control. Their discovery potential is therefore important. On the other hand, as far as we understand, their capacity to give interesting limits is reduced, and unless quoting unreasonable precisions in the QED test i.e. the measurement of Δn , astrophysical existing limits [7] are already better than the ones that can be obtained via the vacuum magnetic birefringence.

The BMV experiment is a collaboration between the Laboratoire Collision Agrégats Réactivité (LCAR) of Toulouse, France [8], the Laboratoire National Champs Magnétiques Intenses (LNCMI) of Toulouse and Grenoble, France [9], and the Laboratoire Matériaux Avancés (LMA) of Lyon [10], France. The apparatus is set up at the Toulouse site of the LNCMI, which is a laboratory specialized in pulsed magnetic fields.

To produce very high magnetic fields the only way is to have a strong current flowing into a coil. There are two main problems : heating and magnetic pressure. Pulsed fields have the advantage compared with continuous fields that coils do not have the time to heat and very high fields can be reached. On the other hand, magnetic pressure which is proportional to $\frac{B^2}{\mu_0}$ becomes very important and ultra strong conductors and special external reinforcement are needed. At the LNCMI of Toulouse, thanks to a 14 MJ bank of capacitors, pulsed fields of more than 80 T have been obtained using coils of solenoid type.

2 Present status

The BMV experiment is detailed in ref. [1]. We need a transverse magnetic field, since 2002 we have designed and tested a new geometry of coils, namely X-coils, which have reached more than 14 T over 0.25 m length corresponding to a $28 \text{ T}^2\text{m}$. The coils are operated at liquid nitrogen temperature and the repetition rate is about 5 pulses per hour. In 2006 a clean room to host the experiment was realised. To increase the optical path of light in the magnetic field region we developed a Fabry-Perot cavity to which a laser is locked. We plan to use for such a cavity mirrors provided by LMA. Mirror losses are of the order of a few ppm and they always have to be kept in a dust free environment. A first version of the experiment mounted in




Figure 1: 1 atm Helium Cotton-Mouton effect : raw experimental data

our clean room consists of two X-coils with their cryostats surrounding the vacuum pipe and four vacuum chambers where polarizing prisms and cavity mirrors are located. Optics under vacuum is all placed on a 3.6 m table which satisfies our project requirement to have a table top experiment. As for data analysis, at the exit of the cavity we measure both the light intensity corresponding to light polarized like the light entering in the magnetic field region I_t and the light intensity corresponding to light polarized perpendicularly to the polarization of the light entering in the cavity I_{ext} . The ellipticity to be measured $\Psi(t)$ can be written as

$$\Psi(t) = \Gamma \sqrt{1 + \frac{I_{ext} - I_t(\sigma^2 + \Gamma^2)}{I_t \Gamma^2}} - \Gamma \quad (2)$$

where Γ is the ellipticity due to the cavity, and σ^2 is the polarizer extinction. When no magnetic field is present, and therefore $\psi(t) = 0$ one can obtain the value of Γ as a function of σ^2 , I_t and I_{ext} . Since in principle $\Psi(t) = k B^2(t)$, for each pulse we calculate the correlation between $\Psi(t)$ and $B^2(t)$, and finally a statistical analysis will give the mean value of k and its error.

We have recently taken data to measure the Cotton-Mouton effect of different gases [11], like air, molecular nitrogen and helium. The Fabry-Perot cavity used for such measurements is 2.2 meter long, corresponding to a free spectral range of 68 MHz. Once the laser is locked to the cavity we infer the cavity finesse by the measurement of the intensity decay time τ following a sudden stop of the light entering in the cavity. Typically τ is about 300 μ s corresponding to a finesse of about 130 000, a cavity linewidth of about 520 Hz (FWHM), and a quality factor Q of about 5.4×10^{11} . Magnetic pulse duration is about 4 ms, which is comparable with the photon lifetime in the cavity. Actually, we have observed that ellipticity pulse is deformed by the cavity acting as a low pass filter of about 260 Hz cut-off frequency, as predicted in ref. [1]. In figure 1 we show the raw data corresponding to the Cotton-Mouton effect of Helium gas, which is the smallest that one can find in nature (except vacuum effect).

We have performed some measurements at different pressure between 0.1 to 1 atm. We have obtained a preliminary value for the Δn per Tesla of a 1 atm of Helium gas of $\Delta n = (2.1 \pm 0.4) \times 10^{-16}$ in agreement with the theoretical prediction $\Delta n = 2.4 \times 10^{-16}$ and the other

three experimental values published [12]. We have also performed measurements in vacuum that are compatible with zero within the errors. Thanks to the pulse duration our frequency working point is around 500 Hz, current sensitivity is about $10^{-7} \text{ 1}/\sqrt{\text{Hz}}$, mostly limited by the photodiode noise equivalent power. We are upgrading the detection system. We are also working to decrease Γ and σ^2 which also limit the sensitivity.

3 Short term and long term perspectives

In the near future, we plan to measure helium Cotton-Mouton effect precisely with a sensitivity better than $\Delta n = 10^{-19}$ per Tesla per pulse, which also will give a precise calibration of our instrument. Vacuum measurements will follow. Using LMA mirrors (the expected cavity finesse is about 600 000) we plan to reach $\Delta n = 10^{-22}$ per Tesla and therefore to give new terrestrial limits on the oscillations of photons into massive particles in 2010.

Long terms perspectives depend on the possibility to have higher magnetic fields. We have designed a new pulsed coil, named XXL-Coil, which should reach a field higher than 25 T when a current higher than 27 000 A is injected. An XXL-Coil should provide more than 200 T²m. One XXL-Coil is under construction, and winding started in July 2009. Tests will follow as soon as possible. Eventually, three of them will be installed in the final set up. We hope in the next few years to finally reach our goal that is to measure the vacuum magnetic birefringence.

This work is supported by the *ANR-Programme non thématique* (ANR-BLAN06-3-139634), and by the *CNRS-Programme National Astroparticules*.

References

- [1] R. Battesti et al., Eur. Phys. J. D **46**, 323 (2008).
- [2] Z. Bialynicka-Birula and I. Bialynicki-Birula, Phys. Rev. D **2**, 2341 (1970).
- [3] W. Heisenberg and H. Euler, Z. Phys. **38**, 714 (1936).
- [4] V.I.Ritus, Sov. Phys. JETP, **42** 774 (1975).
- [5] Constants <http://physics.nist.gov/cuu/Constants/index.html>
- [6] R. Cameron et al., Phys. Rev. D **47**, (1993) 3707.
- [7] K.Zioutas et al. (CAST Collaboration), Phys. Rev. Lett. **94**, 121301 (2005).
- [8] <http://www.lcar.ups-tlse.fr/?lang=en>
- [9] <http://lncmi.cnrs.fr/>
- [10] <http://lma.in2p3.fr/Lmagb.htm>
- [11] C. Rizzo, A. Rizzo and D. M. Bishop, Int. Rev. Phys. Chem. **16**, 81 (1997).
- [12] R. Cameron et al., Phys. Lett. A **157**, 125 (1991); K. Muroo et al., J. Opt. Soc. Am. B **20**, 2249 (2003) ; M. Bregant et al., Chem. Phys. Lett. **471**, 322 (2009).