

NLO and parton showers: the POWHEG-BOX

Simone Alioli

DESY, Platanenallee 6, 15738 Zeuthen, Germany

DOI: <http://dx.doi.org/10.3204/DESY-PROC-2010-01/238>

We describe the POWHEG-BOX package, a general computer code framework for implementing NLO calculations in Shower Monte Carlo programs according to the POWHEG method. The program can be downloaded from <http://powhegbox.mib.infn.it/~nason/POWHEG>.

1 The POWHEG method

Next-to-leading order (NLO) perturbative QCD computations as well as *Shower Monte Carlo* (SMC) programs are fundamental tools for the present-days particle physics phenomenology. In particular, SMC programs incorporate the description of a generic high-energy hadronic collision process, starting from the collision between constituents and developing the parton shower, that increases the number of final-state particles by means of strongly ordered subsequent emissions. Eventually, the interface with a phenomenological hadronization model, enables the comparison with experimental data. For these reasons, they are routinely used by experimentalists to simulate signal and backgrounds processes in physics searches. Nevertheless, the demand for better and better predictions from high energy experiments calls for improving the precision of existing SMC's, including NLO corrections. The MC@NLO [1] method has shown first how to reach NLO accuracy for inclusive quantities, implementing the hard subprocess at NLO and developing showers within the *leading logarithmic approximation*, avoiding double counting of radiation. In this way one achieves benefits of both approaches: exclusive final states generation of SMC's and accuracy of NLO calculations.

The POWHEG method is a different prescription for interfacing NLO calculations with parton shower generators. It was first suggested in Ref. [2], and was described in great detail in Ref. [3]. This method is independent from the Monte Carlo program used for subsequent showering and generates positive weighted events only. In these respects it improves the MC@NLO approach. Until now, the POWHEG method has been successfully applied to several processes, both at lepton [4, 5] and hadron colliders [6, 7, 8, 9, 10, 11, 12, 13, 14]. In these implementations, it has been interfaced to the HERWIG [15, 16], PYTHIA [17] and HERWIG++ [18] SMC programs.

In the POWHEG method the hardest radiation¹ is generated first, independently from the following ones. Schematically², the hardest radiation is distributed according to

$$d\sigma = \bar{B}(\Phi_B) d\Phi_B \left[\Delta_R(p_T^{\min}) + \frac{R(\Phi_R)}{B(\Phi_B)} \Delta_R(k_T(\Phi_R)) d\Phi_{\text{rad}} \right], \quad (1)$$

¹By hardest we mean the radiation with the highest transverse momentum, either with respect to the beam for initial state radiation (ISR), either with respect to another parton for final state radiation (FSR).

²Here we avoid entering into the details concerning the radiation regions and the correct treatment of the associated flavour configurations [2, 3].

where $B(\Phi_B)$ is the Born contribution and

$$\bar{B}(\Phi_B) = B(\Phi_B) + \left[V(\Phi_B) + \int d\Phi_{\text{rad}} R(\Phi_R) \right] \quad (2)$$

is the NLO differential cross section at fixed underlying Born kinematics and integrated over the radiation variables. The transverse momentum of the emitted parton, with respect to the beam or to another particle, depending on the region of singularity, is denoted by $k_T(\Phi_R)$. The lower cutoff p_T^{min} is necessary in order to avoid the coupling constant to reach unphysical values. $V(\Phi_B)$ and $R(\Phi_R)$ are the virtual and the real corrections and in the expression within the square bracket in Eq. (2) a procedure that takes care of the cancellation of soft and collinear singularities is understood, *e.g.* Frixione-Kunszt-Signer (FKS) [19] or Catani-Seymour (CS) [20] dipole subtraction. Then,

$$\Delta_R(p_T) = \exp \left[- \int d\Phi_{\text{rad}} \frac{R(\Phi_R)}{B(\Phi_B)} \theta(k_T(\Phi_R) - p_T) \right] \quad (3)$$

is the POWHEG Sudakov, that is the probability of not having an emission harder than p_T . Equation (1) can be seen as an improvement on the original SMC hardest-emission formula, since the Born cross section is replaced with $\bar{B}(\Phi_B)$ which is normalized to NLO by construction. At small transverse momenta the POWHEG Sudakov becomes equal to a standard SMC one. However, the NLO accuracy of Eq. (1) is maintained for inclusive quantities. Moreover, the high- p_T radiation region is correctly described by the real contributions

$$d\sigma \approx \bar{B}(\Phi_B) d\Phi_B \frac{R(\Phi_R)}{B(\Phi_B)} d\Phi_{\text{rad}} \approx R(\Phi_R) d\Phi_B d\Phi_{\text{rad}}, \quad (4)$$

since $\Delta_R \approx 1$ and $\bar{B}/B \approx 1 + \mathcal{O}(\alpha_s)$. After having generated the hardest radiation, one can interface with any available shower generator, in order to develop the rest of the shower. To avoid the double-counting, the SMC is required to be either p_T -ordered or to have the ability to veto emissions with a p_T harder than the first one³.

2 The POWHEG-BOX

In a real collision process several colored massless partons are present, either in the initial or the final state. One thus should repeat the procedure outlined in Sec. 1 for every possible singular region, associated with any massless colored leg becoming collinear to another one, or soft. In order to do this, the full real emission cross section is decomposed into a sum of terms, each of which has at most one collinear and one soft singularities. The radiation is then generated independently in each of this regions, but only the hardest radiation is retained and the event is generated according to the flavour and kinematics associated to it. Because of this complexity, an automatic tool, the POWHEG-BOX, has been built [22], in order to help the inclusion of new processes. On the other hand, the POWHEG-BOX may also be seen as a library, where previously implemented processes are available in a common framework. The processes implemented so far and already available in the public version comprise: $W, Z/\gamma$ single vector boson production, Higgs boson through gluon and Vector Boson Fusion, single-top in s - and t -channel.

³All modern SMC generators compliant with the *Les Houches Interface for User Processes* [21] should implement this last feature.

The user wishing to include a new NLO calculation must only know how to communicate the needed information to the POWHEG-BOX. This happens either defining the appropriate variables, either providing the necessary Fortran routines. The required inputs ⁴ are:

1. The number of legs in Born process, *e.g.* `nlegborn= 5` for $pp \rightarrow (Z \rightarrow e^+e^-) + j$.
2. The list of Born and Real processes flavours, according to PDG [23] conventions⁵. Flavor is defined incoming (outgoing) for incoming (outgoing) fermion lines, *e.g.* `[5,2,23,6,3,0]` for $bu \rightarrow Ztsq$.
3. The Born phase space routine that, given random numbers in the unit `ndims`-dimensional hypercube, set the Born phase space Jacobian and returns the momenta in lab. and CM frames and the Bjorken x 's.
4. The routines that performs the initialization of the couplings, and the setting of the factorization and renormalization scales.
5. The Born squared amplitude routine that, for a given set of momenta and a flavour configuration, returns $\mathcal{B} = |\mathcal{M}|^2$, summed and averaged over color and helicities as well as the color-ordered Born squared amplitudes \mathcal{B}_{jk} and the helicity correlated Born squared amplitudes $\mathcal{B}_{k,\mu\nu}$, where k runs over all external gluons.
6. The real emission squared amplitude routine, that returns \mathcal{R} for a given momenta and flavour list.
7. The finite part of the interference of Born and virtual amplitude contributions $\mathcal{V}_b = 2\text{Re}\{\mathcal{B} \times \mathcal{V}\}$, after factorizing out the common factor $\mathcal{N} = \frac{(4\pi)^\epsilon}{\Gamma(1-\epsilon)} \left(\frac{\mu^2}{Q^2}\right)^\epsilon$. The routine is again defined with momenta and flavour list as input.
8. The Born color structures in the large N_c limit, set through the Les Houches interface [21].

We remark that items (1-7) are the usual ingredients needed to perform a NLO calculation in any subtraction method. Item (8) is instead needed to provide a defined color structure to the SMC generator. Internally, the POWHEG-BOX implements the FKS subtraction procedure in a general way. At the beginning, it automatically evaluates the combinatorics, identifying all the singular regions and the corresponding underlying Born contributions. It also performs the projection of real contributions over the singular region and computes the subtraction counterterms, from soft and collinear approximations of real emissions. Then, it builds the ISR and FSR phase space, according to the FKS parametrization of the singular region and performs the integration. Eventually, one gets the NLO differential cross section. At this stage, one can also interface to some analysis routine to obtain NLO differential distributions as a byproduct. After the integration stage, it performs the calculation of upper bounds for an efficient generation of Sudakov-suppressed events and then the generation of hardest radiation, according to the POWHEG Sudakov. At this point, the generated event, which contains at most only one extra radiation, has to be passed to a standard SMC program, for developing the rest

⁴For the precise definition of the following routines, we refer the reader to Ref. [22].

⁵Internally gluons are labelled 0 instead of the PDG value of 21. At the moment of writing the event on the Les Houches common block, the PDG value is restored.

of the shower. This can be done either on-the-fly or storing the events on a Les Houches events file [24]. Standard analysis routines, at partonic and hadronic level, are provided for included processes, as well as drivers for common SMC generators. Users can modify them or implement new ones.

2.1 Recent developments

Recently, thanks to this framework, the relatively complex process of $Z + 1$ jet production has been implemented [25]. This is a promising processes for jet calibration with the early LHC data. It is also an important source of missing energy signal as well as a background to many new physics searches. In experimental studies carried up until now, the NLO theoretical calculations were supplemented by correction factors for shower, hadronization and underlying event effects. However, these factors were evaluated by means of standard LO SMC programs. It is clear the advantage to have a SMC program which is NLO accurate, in order to ease and improve the comparisons with experimental results. We have tried a simple approach [25] to merge consistently Z and $Z + 1$ jet samples, in order to obtain a description as smooth and accurate as possible both in the low and high-transverse momentum regions. The results are showed in Figs. 1 and 2.




Figure 1: The p_T distribution of the Z boson in single Z production (blue dashed curve), in $Z + 1$ jet (black solid) and in the merged sample (red solid).




Figure 2: The p_T distribution of the *next-to-hardest* jet in single Z production (blue dashed curve), in $Z + 1$ jet (black solid) and in the merged sample (red solid).

From the two figures, one can see how the merged sample models both the single Z Sudakov form factor, that plays an important role in resumming collinear/soft logarithms in the low- p_T region and the high- p_T behaviour of the next-to-hardest jet, which follows the $Z + 1$ jet distribution. In this last figure, jets are reconstructed according to the k_T algorithm, imposing also an angular separation $R_{jj} > 3$.

Acknowledgments

All results presented in this talk have been obtained in collaboration with P. Nason, C. Oleari and E. Re. This work has been supported in part by the Deutsche Forschungsgemeinschaft in SFB/TR 9.

References

- [1] S. Frixione and B. R. Webber, JHEP **0206** (2002) 029 [arXiv:hep-ph/0204244].
- [2] P. Nason, JHEP **0411** (2004) 040 [arXiv:hep-ph/0409146].
- [3] S. Frixione, P. Nason and C. Oleari, JHEP **0711** (2007) 070 [arXiv:0709.2092 [hep-ph]].
- [4] O. Latunde-Dada, S. Gieseke and B. Webber, JHEP **0702** (2007) 051 [arXiv:hep-ph/0612281].
- [5] O. Latunde-Dada, Eur. Phys. J. C **58** (2008) 543 [arXiv:0806.4560 [hep-ph]].
- [6] P. Nason and G. Ridolfi, JHEP **0608** (2006) 077 [arXiv:hep-ph/0606275].
- [7] S. Frixione, P. Nason and G. Ridolfi, JHEP **0709** (2007) 126 [arXiv:0707.3088 [hep-ph]].
- [8] S. Alioli, P. Nason, C. Oleari and E. Re, JHEP **0807** (2008) 060 [arXiv:0805.4802 [hep-ph]].
- [9] K. Hamilton, P. Richardson and J. Tully, JHEP **0810** (2008) 015 [arXiv:0806.0290 [hep-ph]].
- [10] A. Papaefstathiou and O. Latunde-Dada, JHEP **0907** (2009) 044 [arXiv:0901.3685 [hep-ph]].
- [11] S. Alioli, P. Nason, C. Oleari and E. Re, JHEP **0904** (2009) 002 [arXiv:0812.0578 [hep-ph]].
- [12] K. Hamilton, P. Richardson and J. Tully, JHEP **0904** (2009) 116 [arXiv:0903.4345 [hep-ph]].
- [13] S. Alioli, P. Nason, C. Oleari and E. Re, JHEP **0909** (2009) 111 [Erratum-ibid. **1002** (2010) 011] [arXiv:0907.4076 [hep-ph]].
- [14] P. Nason and C. Oleari, JHEP **1002** (2010) 037 [arXiv:0911.5299 [hep-ph]].
- [15] G. Corcella *et al.*, JHEP **0101** (2001) 010 [arXiv:hep-ph/0011363].
- [16] G. Corcella *et al.*, arXiv:hep-ph/0210213.
- [17] T. Sjostrand, S. Mrenna and P. Z. Skands, JHEP **0605** (2006) 026 [arXiv:hep-ph/0603175].
- [18] M. Bahr *et al.*, Eur. Phys. J. C **58**, 639 (2008) [arXiv:0803.0883 [hep-ph]].
- [19] S. Frixione, Z. Kunszt and A. Signer, Nucl. Phys. B **467** (1996) 399 [arXiv:hep-ph/9512328].
- [20] S. Catani and M. H. Seymour, Nucl. Phys. B **485** (1997) 291 [Erratum-ibid. B **510** (1998) 503] [arXiv:hep-ph/9605323].
- [21] E. Boos *et al.*, arXiv:hep-ph/0109068.
- [22] S. Alioli, P. Nason, C. Oleari and E. Re, JHEP **1006** (2010) 043 [arXiv:1002.2581 [hep-ph]].
- [23] <http://pdg.lbl.gov>
- [24] J. Alwall *et al.*, Comput. Phys. Commun. **176** (2007) 300 [arXiv:hep-ph/0609017].
- [25] S. Alioli, P. Nason, C. Oleari, and E. Re, In preparation.