Photon-axion oscillations and the transparency of the universe

Alessandro Mirizzi¹, Daniele Montanino²

¹Institut für Theoretische Physik, Universität Hamburg, Luruper Chaussee 149, 22761 Hamburg, Germany

²Dipartimento di Fisica, Università del Salento, Italy & Istituto Nazionale di Fisica Nucleare sez. di Lecce, Italy

DOI: http://dx.doi.org/10.3204/DESY-PROC-2010-03/montanino_daniele

Universe should be opaque to photons with energy \geq TeV due scattering on the extragalactic background light during their propagation. However, a surprisingly high degree of transparency of the universe has been observed. In order to explain this fact, the conversion between photons and hypothetical axion-like particles in the turbulent extragalactic magnetic field has been invoked. We have derived new equations to calculate the mean survival probability of the photons. We have also found that the photon transfer functions on different lines of sight could have relevant deviations with respect to the mean value, producing both an enhancement or a suppression in the observable photon flux.

1 Introduction

Axion-like particles (ALP's) with a two-photon vertex are predicted in many extensions of the Standard Model. Pseudoscalar ALP's couple with photons through the effective coupling $g_{a\gamma}\tilde{\mathbf{F}}\mathbf{F}a$, where *a* is the ALP field with mass m_a , \mathbf{F} and $\tilde{\mathbf{F}}$ are the electromagnetic field-strength tensor and its dual, and $g_{a\gamma}$ the ALP-photon coupling. As a consequence of this coupling, ALP's and photons do oscillate into each other in an external magnetic field.

ALP's could play an intriguing role in astrophysics. Indeed, photons emitted by distant sources and propagating through cosmic magnetic fields can oscillate into ALP's. In particular, in the last recent years photon-ALP conversions have been proposed as a mechanism to avoid the opacity of the extragalactic sky to high-energy radiation due to pair production on the Extragalactic Background Light (EBL) [1, 2, 3]. At this regard, recent observations of cosmologically distant gamma-ray sources by ground-based gamma-ray telescopes have revealed a surprising degree of transparency of the universe to very high-energy (VHE) photons (E > 100 GeV). Oscillations between very high-energy photons and ALP's could represent an intriguing possibility to explain this puzzle. In fact, if VHE photons are converted into ALP's and then regenerated, they should not suffer absorption effects while they propagate as ALP's. We have worked out simple equations that describe the mean and the variance of the photon transfer function in the random structure of the intergalactic magnetic field without a montecarlo simulation [4].

2 VHE $\gamma - a$ mixing

The evolution equation for photon moving in the x_3 direction can be written as

$$\frac{\partial}{\partial x_3} \left(\begin{array}{c} A_1 \\ A_2 \\ a \end{array} \right) = -i\mathcal{H} \left(\begin{array}{c} A_1 \\ A_2 \\ a \end{array} \right) \,,$$

where, in the presence of absorption and in the high energy limit ($E \ge 100 \text{ GeV}$) the (no longer hermitian) hamiltonian \mathcal{H} can be written as

$$\mathcal{H} = \begin{bmatrix} -i\frac{\Gamma_{\gamma}(E)}{2} & 0 & \frac{g_{a\gamma}B_T}{2}c_{\phi} \\ 0 & -i\frac{\Gamma_{\gamma}(E)}{2} & \frac{g_{a\gamma}B_T}{2}s_{\phi} \\ \frac{g_{a\gamma}B_T}{2}c_{\phi} & \frac{g_{a\gamma}B_T}{2}s_{\phi} & 0 \end{bmatrix} \,.$$

Here $\mathbf{B}_T = \mathbf{B} - B_3 \mathbf{e}_3$ is the transverse component of the external magnetic field, $c_{\phi} \equiv \cos \phi = \mathbf{B}_T \cdot \mathbf{e}_1 / B_T$, and Γ_{γ} is the absorption rate for the pair production process $\gamma^{\text{VHE}} \gamma^{\text{bkg}} \rightarrow e^+ e^-$, where γ^{bkg} is a background (EBL) photon. We stick in the hypothesis that $m_a < 10^{-10}$ eV so that the axion mass term in the hamiltonian can be neglected.

3 Mean photon transfer function

Very high-energy gamma-rays propagate in the extragalactic magnetic fields during their route to the Earth which presumably have a turbulent structure. Let us now consider the propagation of photons in many domains of equal size $l \simeq 1$ Mpc in our case) in which the magnetic field has (constant) random values and directions. Along a given line of sight, the angles ϕ are randomly distributed in $[0, 2\pi)$. During their path with a total length L, photons cross $k = 1, \ldots n$ domains (n = L/l) representing a given random realization of B_k and ϕ_k . Since we cannot know this particular configuration, we perform an ensemble average over all the possible realizations on the $1, \ldots n$ domains. Since we have to perform averages on the random configurations of the intergalactic magnetic field, it is convenient to stick in the formalism of the density matrix. Defining this ensemble average as $\bar{\rho}_n = \langle \rho_n \rangle_{1...n}$, we have

$$\bar{\rho}_n = \langle e^{-i\mathcal{H}_n l} \cdot \bar{\rho}_{n-1} \cdot e^{i\mathcal{H}_n^{\dagger} l} \rangle_n$$

Expanding \mathcal{H} at second order and performing the ensemble average, we finally arrive at a system of two coupled differential equations $(dy = P_{a\gamma} dx_3/l)$ [4]

$$\frac{d}{dy} \begin{pmatrix} T_{\gamma} \\ T_{a} \end{pmatrix} = \begin{bmatrix} -\alpha - \frac{1}{2} & 1\\ \frac{1}{2} & -1 \end{bmatrix} \begin{pmatrix} T_{\gamma} \\ T_{a} \end{pmatrix}, \qquad (1)$$

where we have used the approximation $\bar{\rho}_n - \bar{\rho}_{n-1} \simeq l\partial_{x_3}\bar{\rho}(x_3)$. Here $T_{\gamma} \equiv I_{\gamma}(y)/I_{\gamma}(0) = \bar{\rho}_{11} + \bar{\rho}_{22}$ (since the two polarization states are indistinguishable) and $T_a = \bar{\rho}_{aa}$ are the mean transfer functions for the photon and for the ALP respectively; $P_{a\gamma} = g_{a\gamma}^2 |\mathbf{B}|^2 l^2/6$ is the average photon-ALP conversion probability in each domain (in absence of absorption) and finally $\alpha = \Gamma_{\gamma} l/P_{a\gamma}$ is the ratio between the absorption probability and the conversion probability. In realistic astrophysical situations both $P_{a\gamma}$ and α are functions of the distance, due to the redshift dependence of the extragalactic magnetic field and of the EBL. PHOTON-AXION OSCILLATIONS AND THE TRANSPARENCY OF THE UNIVERSE



Figure 1: Photon transfer function $T_{\gamma}(y)$ for $\alpha = 0$ (the upper panels) and $\alpha = 1$ (lower panels) for a given random realization of the magnetic field (left) and for 20 realizations (right).

In particular, in the case of strong absorption $(\alpha \gg 1)$ from Equation (1) we obtain $T_{\gamma} \propto (\Gamma_{\gamma})^{-2}$. Using the approximate power–law spectrum for the EBL we observe that the transfer function would drop as a power of the energy (rather than exponentially as expected without ALP mixing). Moreover, also the attenuation of the transfer function with the distance is less than in the case of absence of conversions. In fact, in this case we have $T_{\gamma} \propto e^{-\Gamma_{\gamma} x/\alpha}$, in which the argument of the exponential is suppressed by a factor α with respect to the no-conversion case. Thus this effect would explain the high transparency of the universe.

In [4] we have also calculated the root mean square δT_{γ} for the distribution of the transfer function in different random realizations of the magnetic field. This result is useful to estimate the uncertainty associated with the averaging procedure. In Figure 1 we compare the transfer function $T_{\gamma}(y)$ (continuous lines) calculated from Eq. (1) with those calculated for a given random realization of the magnetic field along the photon line of sight, with and without absorption (left panels) and for 20 realizations (right panels). For comparison we have also shown the dispersion around the mean $T_{\gamma} \pm \delta T_{\gamma}$ (dashed lines).

4 Results

In Figure 2 we show the photon transfer function as function of the energy for four different values of the redshift of the emitting source. We have used a realistic EBL model [5] which provides a strict lower-limit flux for the extragalactic background light which gives us the maximal possible transparency compatible with the standard expectations, so that an evidence of a greater transparency would have to be attributed to nonstandard effects in the photon propagation. We see that in absence of ALP conversions the photon transfer function would be strongly suppressed at energies above $E \geq 100$ GeV, the stronger the suppression the larger



Figure 2: VHE photons transfer function T_{γ} in function of the observed photon energy E, for different values of the redshift z.

the redshift. We also realize that the spread in the possible values of T_{γ} would make difficult to infer strong conclusions about ALP mixing observing only few sources. To test this effect we would need to collect data from sources along different directions in the sky in order to perform a study of the photon energy distributions, from which we could hope to infer possible hints of ALP's. A further signature of these stochastic conversions would be the detection of peculiar direction-dependent dimming effects in the diffuse photon radiation observable in GeV range, testable with the FERMI (previously called GLAST) experiment.

References

- A. De Angelis, O. Mansutti and M. Roncadelli, "Evidence for a new light spin-zero boson from cosmological gamma-ray propagation?," Phys. Rev. D 76, 121301 (2007).
- [2] A. De Angelis, O. Mansutti and M. Roncadelli, "Axion-Like Particles, Cosmic Magnetic Fields and Gamma-Ray Astrophysics," Phys. Lett. B 659, 847 (2008).
- [3] A. De Angelis, O. Mansutti, M. Persic and M. Roncadelli, "Photon propagation and the VHE gamma-ray spectra of blazars: how transparent is the Universe?," Mon. Not. R. Astron. Soc. 394, L21 (2009).
- [4] A. Mirizzi and D. Montanino, "Stochastic conversions of TeV photons into axion-like particles in extragalactic magnetic fields," JCAP 0912 (2009) 004.
- [5] T. M. Kneiske and H. Dole, "A strict lower-limit EBL: Applications on gamma-ray absorption," AIP Conf. Proc. 1085, 620 (2009).