Limit on Lorentz-Invariance- and CPT-Violating Neutron Spin Interactions Using a ³He-¹²⁹Xe Comagnetometer

Fabian Allmendinger¹, Ulrich Schmidt¹, Werner Heil², Sergei Karpuk², Yury Sobolev², Kathlynne Tullney²

¹Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Germany ²Institut für Physik, Johannes Gutenberg-Universität Mainz, Germany

DOI: http://dx.doi.org/10.3204/DESY-PROC-2014-04/41

We performed a search for a Lorentz-invariance- and CPT-violating coupling of the ³He and ¹²⁹Xe nuclear spins to posited background fields. Our experimental approach is to measure the free precession of nuclear spin polarized ³He and ¹²⁹Xe atoms using SQUIDs as detectors. As the laboratory reference frame rotates with respect to distant stars, we look for a sidereal modulation of the Larmor frequencies of the co-located spin samples. As a result we obtain an upper limit on the equatorial component of the background field $\tilde{b}_{\perp}^{1} < 8.4 \cdot 10^{-34}$ GeV (68% C.L.). This experiment is currently the most precise test of spin anisotropy due to the excellent long spin-coherence time.

1 Introduction and Experimental Setup

In the context of the Standard-Model Extension (SME)[1, 2, 3], couplings of the neutron or proton spin $\sigma^{n,p}$ to relic background fields $\tilde{b}^{n,p}$ are discussed. The background fields have distinct directions in space and correspond to preferred spin directions. These couplings with the potential $V = \tilde{b}^{n,p} \cdot \sigma^{n,p}$ are purely non-magnetic, but change the energy levels of spins in a magnetic field, which can be detected by changes in the Larmor frequency of precessing spins.

Nuclear spin clocks, based on the detection of free spin precession of gaseous nuclear polarized ³He and ¹²⁹Xe atoms with LT_C SQUIDs as low-noise magnetic flux detectors are used as ultra-sensitive probe for nonmagnetic spin interactions, since the magnetic interaction (Zeeman term) drops out in the case of co-located spin samples (comagnetometry). Measurements of uninterrupted precession of $T \sim 1$ day can be achieved at the present stage of investigation due to long spin-coherence times. The principle of measurement is to search for sidereal variations of the precession frequency of co-located spin species while the Earth and hence the laboratory reference frame rotates with respect to distant stars.

To give a short overview of the setup (details are given in Ref. [4]): The two polarized gas species (and N₂ as a buffer gas) are filled into a low-relaxation spherical glass cell with radius R = 5 cm. Typically, the optimum conditions in terms of long transverse relaxation times (T_2^*) and high Signal-to-Noise Ratio are met at a gas mixture with pressures of $p_{He} = 3$ mbar, $p_{Xe} = 5$ mbar, $p_{N2} = 25$ mbar. The cell is positioned in a homogeneous static magnetic field (about 400 nT)

LIMIT ON LORENTZ-INVARIANCE- AND CPT-VIOLATING NEUTRON SPIN ...

that is generated by Helmholtz coils mounted inside the strongly magnetically shielded room BMSR-2 at the Physikalisch-Technische Bundesanstalt in Berlin. At that field strength, the Larmor frequencies of ³He and ¹²⁹Xe are about $\omega_{He} \approx 2\pi \cdot 13$ Hz and $\omega_{Xe} \approx 2\pi \cdot 4.7$ Hz, respectively. To measure these precession frequencies very precisely, low-noise low-temperature DC-SQUID gradiometers are used as magnetic flux detectors. Due to the very low field gradients in the order of pT/cm at the location of the cell, the transverse relaxation times reached $T_2^* = 8.5$ h for ¹²⁹Xe and up to $T_2^* = 100$ h for ³He [4]. The measured signal amplitudes at the beginning of the measurement were up to $A_{\text{He}} = 20$ pT and $A_{\text{Xe}} = 8$ pT for ³He and ¹²⁹Xe, respectively. The noise level (combination of four gradiometers) was $\rho = 3$ fT/ $\sqrt{\text{Hz}}$. Due to the long spin-coherence time and the high initial Signal-to-Noise Ratio, the spin precession could be monitored for more than one day, which improves the sensitivity remarkably (see below).

2 Data Evaluation and Results

To be sensitive to tiny nonmagnetic interactions, one has to consider the weighted difference of the respective Larmor frequencies of the co-located spin samples, or the corresponding time integral (weighted phase difference), which are defined by

$$\Delta \omega = \omega_{\rm He} - \frac{\gamma_{\rm He}}{\gamma_{\rm Xe}} \omega_{\rm Xe} \quad \text{and} \quad \Delta \Phi = \Phi_{\rm He} - \frac{\gamma_{\rm He}}{\gamma_{\rm Xe}} \Phi_{\rm Xe} \quad . \tag{1}$$

In doing so, magnetic field fluctuations are canceled, i.e. in principle $\Delta \omega = 0$ and $\Delta \Phi = \text{const.}$ if there are no further interactions. However, on a closer look, $\Delta \Phi$ is not constant in time, as higher order effects have to be take into account. These can be parameterized by

$$\Delta\Phi(t) = c_0 + c_1 t + E_{\rm He} e^{-t/T_{2,\rm He}^*} + E_{\rm Xe} e^{-t/T_{2,\rm Xe}^*} + F_{\rm He} e^{-2t/T_{2,\rm He}^*} + F_{\rm Xe} e^{-2t/T_{2,\rm Xe}^*} \quad . \tag{2}$$

The linear contribution stems from Earth's rotation and from chemical shift. The four exponential terms account for the Ramsey-Bloch-Siegert shift [5, 6]. These effects are discussed in Ref. [4]. Finally, the function in Eq. (2) together with the appropriate parameterization of the Lorentz-invariance-violating effect - in this case a sidereal modulation $\propto \tilde{b} \cdot \sin(\Omega_S \cdot t + \varphi_0)$ - is fitted to the combined weighted phase difference data of all measurement runs (7 in total). The resulting estimate on sidereal modulation is compatible with zero within the correlated and uncorrelated uncertainties and can be expressed as an upper limit on the magnitude of the hypothetical background field:

$$\tilde{b}_{\perp}^{n} < 6.7 \cdot 10^{-34} \text{ GeV} (68\% \text{ C.L.})$$
 . (3)

3 Frequency Stability

The benefit of long spin-coherence times can be explained on the basis of the Cramer-Rao Lower Bound (CRLB). The CRLB gives the minimum variance of an unbiased estimator of a deterministic parameter. In this case, the sensitivity of frequency measurements depends on the measurement time with coherent spin precession T, the signal amplitude at the beginning of the measurement A decaying with a time constant T_2^* and the noise level ρ (assuming white noise)[7]. The frequency uncertainty is:

$$\sigma_f \geq \frac{\sqrt{3}}{\pi} \frac{\rho}{A} \cdot T^{-3/2} \cdot \sqrt{C(T, T_2^*)}$$
(4)

PANIC2014

with the dimensionless factor $C(T, T_2^*)$ accounting for the exponential decay of the Signal-to-Noise Ratio. The remarkable result is that the frequency uncertainty decreases with $T^{-3/2}$ for white (Gaussian) noise. The Allan Standard Deviation (ASD) is the appropriate measure to study the temporal characteristics of the ${}^{3}\text{He}{}^{-129}\text{Xe}$ comagnetometer with respect to phase or frequency determination. The ASD method is an established analysis technique for studying the low-frequency component of a time series and is a measure of phase and frequency stability of clocks and oscillators. In the case of the ³He-¹²⁹Xe comagnetometer, one compares the ³He precession frequency to the "scaled" ¹²⁹Xe precession frequency. These two oscillator frequencies are supposed to be the same if all deterministic phase shifts (chemical shift, Earth's rotation, Ramsey-Bloch-Siegert shift etc.) have been incorporated correctly. The behavior of the frequency uncertainty in the ASD plot is shown in Fig. 1 (black symbols) for two different runs. With increasing integration times τ the uncertainty in frequency decreases with $\sigma_{ASD} \propto \tau^{-\frac{3}{2}}$ as expected by the CRLB in Eq. (4). This indicates the presence of pure white (Gaussian) noise. In other words: All deterministic phase shifts (chemical shift, Earth's rotation, Ramsey-Bloch-Siegert shift etc.) have been incorporated correctly. This has been tested for all measurement runs with different experimental conditions (e.g. different sample cells, partial pressures, magnetic field directions, T_2^*) and the ASD plot shows the expected behavior, leading to the conclusion that the fit model in Eq. (2) describes the behavior of the comagnetometer correctly. An incorrect fit model would immediately lead to a deviation from the CRLB power law. This is also demonstrated in Fig. 1 (gray symbols) showing the ASD as a function of τ for the residuals of a fit model where the $F_{\text{He, Xe}}$ terms (called "cross-talk") of Eq. (2) are omitted. For run 6 (the run with largest cross-talk effect), the large deviation from the CRLB power law above $\tau \approx 500$ s is impressive. For run 3 (the run with the smallest cross-talk effect) the deviation in the ASD plot is smaller.

4 Conclusion

Freely precessing gaseous, nuclear polarized ³He and ¹²⁹Xe samples can be used as ultra-sensitive probe for nonmagnetic spin interactions, since the magnetic interaction (Zeeman term) drops out in the case of co-located spin samples. Long spin-coherence times are highly beneficial as the uncertainty in frequency estimation decreases with $T^{-3/2}$ for Gaussian noise according to the CRLB. The ASD method can be used to verify the CRLB power law and thereby rule out the presence of further noise sources (i.e. further deterministic frequency shifts). With a similar setup, upper limits on interactions mediated by axion-like particles were obtained [8]. The next challenging step is to apply this method in the search for an electric dipole moment of ¹²⁹Xe.

References

- [1]~ V. A. Kostelecký and S. Samuel, Phys. Rev. D ${\bf 39},\,683$ (1989).
- [2] D. Colladay and V. A. Kostelecký, Phys. Rev. D 58, 116002 (1998).
- [3] V. A. Kostelecký and C. D. Lane, Phys. Rev. D 60, 116010 (1999).
- [4] F. Allmendinger et. al, Phys. Rev. Lett. 112, 110801 (2014).
- [5] F. Bloch and A. Siegert, Phys. Rev. 57, 522 (1940).
- [6] N. F. Ramsey, Phys. Rev. 100, 1191 (1955).
- [7] W. Heil et al., Ann. Phys. (Berlin) 525, 539 (2013).
- [8] K. Tullney et al., Phys. Rev. Lett. 111, 100801 (2013).



Figure 1: ASD plot of the residual frequency noise of two independent runs (Top: Run 3, Bottom: Run 6). Black symbols: ASD for the correct fit model (Eq. (2)). With increasing integration times τ the uncertainty in frequency decreases as $\sigma \propto \tau^{-\frac{3}{2}}$ indicating the presence of white (Gaussian) noise. Gray symbols: ASD for an incorrect fit model (omitting the Cross-Talk term). For run 6 (the run with largest cross-talk effect), the large deviation from the CRLB power law above a few hundred seconds is impressive. For run 3 (the run with the smallest cross-talk effect) the deviation in the ASD plot is smaller.

PANIC2014