

D2-84/192

# DEUTSCHES ELEKTRONEN-SYNCHROTRON **DESY**

DESY 84-099  
October 1984



RECENT RESULTS ON THE REACTIONS  $\gamma\gamma \rightarrow \eta'$ ,  $A_2$ ,  $(5\pi)^0$  AND UPPER LIMITS

ON  $\gamma\gamma \rightarrow \rho^0\omega$  AND  $\gamma\gamma \rightarrow \omega\omega$  PRODUCTION



by

Jacob Grunhaus

*School of Physics and Astronomy, Tel Aviv University, Tel Aviv, Israel*

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your preprints are promptly included in the  
HIGH ENERGY PHYSICS INDEX ,  
send them to the following address ( if possible by air mail ) :

DESY  
Bibliothek  
Notkestrasse 85  
2 Hamburg 52  
Germany

Recent Results on the Reactions

$\gamma\gamma \rightarrow \eta', A_2, (5\pi)^0$  and Upper Limits  
on  $\gamma\gamma \rightarrow \rho^0\omega$  and  $\gamma\gamma \rightarrow \omega\omega$  Production

PLUTO COLLABORATION

presented by

Jacob Grunhaus\*

School of Physics and Astronomy  
Tel Aviv University, Tel Aviv, Israel

Abstract

Values are presented for the radiative widths,  $\Gamma_{\gamma\gamma}(\eta')$  and  $\Gamma_{\gamma\gamma}(A_2)$  and for the transition form factor of the  $\eta'$  meson. The cross section for the process  $\gamma\gamma \rightarrow 2\pi^+ 2\pi^- \pi^0$  is given. Upper limits for the reactions  $\gamma\gamma \rightarrow \rho^0\omega$  and  $\gamma\gamma \rightarrow \omega\omega$  are set and the relevance of these limits to models which explain the enhancement at threshold for the process  $\gamma\gamma \rightarrow \rho^0\rho^0$  is discussed.

THE  $\eta'$  MESON

We have studied  $\eta'$  meson formation by 2 "quasireal" photons,  $\langle Q^2 \rangle \sim 0.0 \text{ GeV}^2$  via the reaction  $e^+e^- \rightarrow e^+\eta'$ . The  $\eta'$  meson was identified by its decay into  $\rho^0\gamma$ . The transition form factor was also determined using "single tag" events up to a  $Q^2 = 1.0 \text{ GeV}^2$ .

The study is based on an integrated luminosity of  $45 \text{ pb}^{-1}$  at an average  $e^+e^-$  CM energy of 34.7 GeV. A sample of events of 2 unlike charged tracks and 1 photon having  $E_\gamma \geq 100 \text{ MeV}$  was selected and the standard cuts were applied in order to remove background from beam gas interactions, cosmic rays and the  $\gamma\gamma$  annihilation process. The momenta of the 3 final state particles, for the

\* Talk given at the XV International Symposium on Multiparticle Dynamics, Lund, Sweden, June 10-16, 1984.

no tag sample, were required to be consistent with momentum conservation in the  $R-\phi$  plane by satisfying the condition that the angle between the transverse vectors  $\vec{P}_\perp(\pi^+, \pi^-)$  and  $\vec{P}_\perp(\gamma)$  deviate from  $180^\circ$  by at most  $28^\circ$ . The photon energy was then scaled using the recipe,  $E_\gamma = E_\gamma(\text{seen}) \cdot \vec{P}_\perp(\pi^+, \pi^-) / \vec{P}_\perp(\gamma)$  seen. The  $M(\pi^+\pi^-\gamma)$  plot shown in Fig. 1a exhibits a prominent  $\eta'$  signal and an enhancement at the  $A_2$  mass which arises from the  $A_2 \rightarrow \pi^+\pi^-\pi^0$  decay, where only 1 photon was detected. A fit to this mass plot yielded  $243 \pm 17 \eta'$  events, which corresponds to the product  $\Gamma_{\gamma\gamma} B(\eta' \rightarrow \rho^0\gamma) = 1.14 \pm 0.08 \text{ keV}$ . Using the value of  $B(\eta' \rightarrow \rho^0\gamma) = 30.0 \pm 1.6\%$  (1) we get a partial width value of,

$$\Gamma_{\gamma\gamma}(\eta') = 3.80 \pm 0.26 \text{ (stat)} \pm 0.43 \text{ (syst)} \text{ keV.}$$

Combining our value of  $\Gamma_{\gamma\gamma}(\eta')$  with the world value of  $B(\eta' \rightarrow \gamma\gamma) = 1.9 \pm 0.2\%$  (1) we get the total width of the  $\eta'$  to be  $200 \pm 34 \text{ keV}$




Fig. 1. The distribution  $M(\pi^+\pi^-\gamma)$  arising from the collision of a) quasireal photons (no tag) and b) from single tag events with  $\langle Q^2 \rangle = 0.4 \text{ GeV}^2$ .

which is the most accurate determination to date of this parameter. Our value of  $\Gamma_{\gamma\gamma}(\eta')$  is appreciably lower than the world average value of  $5.3 \pm 0.6$  keV<sup>(2)</sup>. We attribute our lower value to our including in the Monte Carlo acceptance studies the "p-wave" nature of the  $\eta' \rightarrow \rho^0 \gamma$  decay. Without the inclusion of the proper dynamics of the  $\eta'$  decay we get a value of  $\sim 5$  keV. In the framework of flavor SU(3) assuming nonet symmetry for the  $0^{-+}$  mesons, one obtains the following mixing angle independent relationship:

$$\frac{\Gamma_{\gamma\gamma}(\eta')}{m^3(\eta')} = \frac{3\Gamma_{\gamma\gamma}(\eta^0)}{m^3(\eta^0)} - \frac{\Gamma_{\gamma\gamma}(\eta)}{m^3(\eta)}$$

Using the recent values of  $\Gamma_{\gamma\gamma}(\eta) = 0.56 \pm 0.12 \pm 0.10$  keV<sup>(3)</sup> and  $\Gamma_{\gamma\gamma}(\eta^0) = 7.56 \pm 0.09$  eV<sup>(4)</sup> this relationship predicts  $\Gamma_{\gamma\gamma}(\eta') = 5.13 \pm 0.84$  keV, a value which is consistent with our determination, although, it is on the high side. It has been suggested that a value of  $\Gamma_{\gamma\gamma}(\eta')$  smaller than the SU(3) expectation might indicate a glueball admixture in the  $\eta'$  (5). Using the definition of the transition form factor in terms of  $\sigma(\gamma\gamma \rightarrow \eta')$  and the width and mass of the  $\eta'$  as given by G. Köpp, T. Walsch and P. Zerwas (6), we have determined its value in the  $Q^2$  range up to 1 GeV<sup>2</sup>. Fig. 2 shows the  $\eta'$  form factor along with the expected  $\rho^0$  pole contribution normalized to  $Q^2 = 0$ .




Fig. 2. The transition form factor for  $\eta' + \gamma\gamma$ . The expected  $\rho^0$  pole contribution normalized to  $Q^2 = 0$  is shown.

$\eta'$  Form Factor 37059

THE  $A_2$  MESON

We have studied the formation of the  $A_2^0$  meson by "quasireal" photons in  $\gamma\gamma$  collisions. The  $A_2^0$  meson was detected by its decay into  $\rho^+\pi^-$ . In addition to the usual background reduction cuts, the sample of events consisting of 2 charged unlike tracks and 2 shower clusters was also kinematically fitted. Imposition of momentum conservation in the R- $\phi$  plane and the requirement that the 2 shower clusters form a  $\pi^0$  meson made possible a 3 constraint fit. Events satisfying the kinematic fit with a  $\chi^2$ -probability greater than 2% were accepted. The  $M(\pi^+\pi^-\pi^0)$  distribution shown in Fig. 3 exhibits a clear  $A_2$  signal which is strongly correlated with the  $\pi^+\pi^0$  mass combination being in the  $\rho^\pm$  meson mass band. The fit shown in Fig. 3 yielded  $97 \pm 18$   $A_2$  events.




Fig. 3. The  $M(\pi^+\pi^-\pi^0)$  distribution for the channel  $\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$ . Entries for which  $M(\pi^+\pi^0) = M(\rho^\pm) \pm 200$  MeV are indicated by the shaded regions. The fit to the  $A_2$  meson is shown.

The detector acceptance was determined from Monte Carlo studies where the events were generated taking into consideration constructive interference between the 2 decay chains,  $A_2^0 \rightarrow \rho^+\pi^-\pi^0$  and  $A_2^0 \rightarrow \rho^-\pi^+\pi^0$  (7).

$$|M|^2 \propto (2J+1) \frac{M_{A_2}}{S} \frac{\Gamma_{YY}(A_2)}{(M_{A_2}^2 - S)^2 + M_{A_2}^2 \Gamma_{A_2}^2} *$$

$$* \frac{\Sigma_{\lambda} \rho_{\lambda\lambda} \left| \frac{T_{\lambda}(A_2 \rightarrow \rho^+ \pi^-)}{M_{\rho}^2 - S_{\rho^+} - i M_{\rho} \Gamma_{\rho^+}} + \frac{T_{\lambda}(A_2 \rightarrow \rho^- \pi^+)}{M_{\rho}^2 - S_{\rho^-} - i M_{\rho} \Gamma_{\rho^-}} \right|^2}{2}$$

The spin (helicity) of the  $A_2$  is denoted by  $J(\lambda)$ ,  $S_{\rho^{\pm}}$  is the  $(\pi^{\pm} \pi^0)$  mass system and  $T_{\lambda}(A_2 \rightarrow \rho^{\pm} \pi)$  is the transition amplitude for the decay  $A_2 \rightarrow \rho^{\pm} \pi$ . The widths for the  $A_2$  and  $\rho$  meson were described by momentum dependent forms.

In order to ascertain the spin-parity of the 3 pion final state we have calculated the Dalitz plot parameter,

$$A = \frac{|\vec{p}_+ \times \vec{p}_-|^2}{\pi \max(|\vec{p}_+ \times \vec{p}_-|^2)_{\pi}}$$




Fig. 4. Distribution of the acceptance corrected value of the Dalitz plot parameter  $A$ . The expected behaviour for a  $2^+$  resonance is shown by the full line, for  $2^-$  by the dashed line, and for  $0^-$  by the point-dashed line.

and this is plotted in Fig. 4. The expected behaviour for the  $J^P$  values which are consistent with a  $3\pi$  final state initiated by 2 photons, namely  $2^+$ ,  $2^-$ ,  $0^-$  are also indicated in Fig. 4. The data is best described by the  $2^+$  curve. We denote the angle between the photon photon axis and the normal to the  $3\pi$  plane (in the  $3\pi$  CM) by the symbol  $\alpha$ . The distribution of  $\cos \alpha$  along with the expected behaviour for the  $J^P(\lambda)$  values,  $2^+(0)$ ,  $2^+(2)$  and  $2^-(0)$  are shown in Fig. 5. The data is well described by  $|\lambda| = 2$  with some admixture of  $\lambda = 0$ . A fit to  $\lambda = 2$  and 0 yields an admixture of  $39 \pm 3\%$  of  $\lambda = 0$ . In our Monte Carlo studies we have generated events with  $\lambda = 2$  and 0, in the same proportion found in the fit of the data. We have found the radiative width of the  $A_2$  meson,  $\Gamma_{YY}(A_2) = 1.06 \pm 0.18 \pm 0.19$  keV, a value consistent with previous measurements of  $0.77 \pm 0.18 \pm 0.27$  (8),  $0.81 \pm 0.19 \pm 0.27$  (9), and  $0.84 \pm 0.07 \pm 0.15$  (10).

In the framework of SU(3) flavour, assuming nonet symmetry, we have the following expression for the radiative widths of the  $2^{++}$  mesons, which is mixing angle independent:

$$\frac{3 \Gamma_{YY}(A_2)}{m_{A_2}^5} = \frac{\Gamma_{YY}(f^+)}{m_{f^+}^3} + \frac{\Gamma_{YY}(f^0)}{m_{f^0}^3}$$




Fig. 5 The acceptance corrected distribution of  $|\cos \alpha|$  and the expected behaviour for the  $J^P(\lambda)$  values  $2^+(2)$  (full line),  $2^-(0)$  (-----),  $2^+(0)$  (.....) and  $0^-$  (-.-.-).

Using the world average value  $\Gamma_{\gamma\gamma}(f^0) = 2.8 \pm 0.2$  keV and the TASSO (11) measurement of  $B(f' \rightarrow K\bar{K}) \Gamma_{\gamma\gamma}(f') = 0.11 \pm 0.02 \pm 0.04$  keV, and if we set  $B(f' \rightarrow K\bar{K}) = 1$ , we obtain a lower limit prediction of  $\Gamma_{\gamma\gamma}(A_2) = 1.06 \pm 0.08$  keV in agreement with our measurement.

UPPER LIMITS ON  $\gamma\gamma \rightarrow \rho^0\omega$  and  $\gamma\gamma \rightarrow \omega\omega$

In view of the sizeable cross section measured for the channel  $\gamma\gamma \rightarrow \rho^0\rho^0$  near threshold (12,13,14) which cannot be accounted for by simple VDM expectations, searches have been carried out for other vector meson final states -  $\rho^+\rho^-$ ,  $\rho^0\omega$  and  $\omega\omega$ . Models proposed to explain the  $\rho^0\rho^0$  enhancement also make predictions for the production of the other vector meson states. Table I below summarizes these predictions.

The JADE Collaboration (15) has set upper limits on  $\gamma\gamma \rightarrow \rho^+\rho^-$  which rule out the interpretation of the  $\rho^0\rho^0$  enhancement as a non-exotic I=0,1 resonance. The proposed models 4) and 5) in Table I are inconsistent with the  $\rho^+\rho^-$  upper limit.

TABLE I

Model	Rate Relative to $\rho^0\rho^0$				
	$\rho^0\rho^0$	$\rho^+\rho^-$	$\omega\omega$	$\rho^0\omega$	$\rho^+\omega$
* 1) VDM (YV-Coupling)	1	0	1/81	1/9	
2) VDM + Regge, Ref. 16	1		5.8	0.5	
3) Alexander et al., Ref. 17	1		-1	-0.2	
4) Quark Model, Ref. 18	1	2	1	-1.4	
5) Quark Model, Ref. 19	1	~1.3			
6) qq $\bar{q}\bar{q}$ , Ref. 20	1	~0	~0.03	~0.6	
7) qq $\bar{q}\bar{q}$ , Ref. 21	1	~0	~0.06	~0.03	

\* does not explain the  $\rho^0\rho^0$  enhancement.

We have carried out a search for  $\gamma\gamma \rightarrow \rho^0\omega$  using a topological sample of 2 positive tracks, 2 negative tracks and 2 neutral clusters, with  $E \geq 100$  MeV. The energies of the 2 photons were scaled so that momentum was conserved in the R- $\phi$  plane. Also the probability that the 2 photons originated from a spin zero particle had

to be greater than 40% as determined from their opening angle (22). A sample of  $52 \pm 8$  events of the exclusive channel  $\gamma\gamma \rightarrow 2\pi^+ 2\pi^-\pi^0$  was found. The cross section for this exclusive 5 pion final state was determined as a function of  $W_{\gamma\gamma}$ , and these values are shown in Fig. 6 along with preliminary values from CELLO (23).

No  $\rho^0$  or  $\omega$  meson signal was detected in this exclusive channel. The acceptance of the PLUTO detector for the reaction  $\gamma\gamma \rightarrow \rho^0\omega \rightarrow 2\pi^+ 2\pi^-\pi^0$  was studied as a function of  $W_{\gamma\gamma}$  using the Monte Carlo simulation program. An isotropic distribution was assumed for the  $\rho^0\omega$  production and the subsequent decays  $\rho^0 \rightarrow \pi^+\pi^-$  and  $\omega \rightarrow \pi^+\pi^-\pi^0$ . The Monte Carlo events when processed through the physics analysis programs yield a clear  $\omega$  signal recoiling from the  $\rho^0$  meson. Upper limits at the 95% confidence level are given in Fig. 7 along with the upper limits set by the JADE Collaboration (15).




Fig. 6. The value of the cross section of the process  $\gamma\gamma \rightarrow 2\pi^+ 2\pi^-\pi^0$  as a function of the  $\gamma\gamma$  CM energy,  $W_{\gamma\gamma}$ . Preliminary results from the CELLO Collaboration are also shown.




Fig. 7. Upper limits (95% conf. level) for the reaction  $\gamma\gamma \rightarrow \rho^0\omega$ . Expectations of Achasov, Ref. 20 and Alexander, Ref. 16 and 17 are shown.

THE CHANNEL  $\gamma\gamma \rightarrow \omega\omega$

In our study of the channel  $\gamma\gamma \rightarrow \omega\omega$  we have considered the sample of events having 2 positive tracks, 2 negative tracks and from one to four neutral clusters with  $E_\gamma > 40$  MeV. We have searched for the  $\omega$  meson through its dominant decay mode  $\gamma_+ \pi^- \pi^0$ . The rather low acceptance of the detector for the complete final state  $2\pi^+ 2\pi^- 2\pi^0$  (4 photons) precluded the use of momentum conservation constraints to fit the events. We have, therefore, taken the approach of maximizing the  $\omega$ -meson signal by reconstructing one  $\omega$  meson using only 1 photon. We have studied the mass system  $\pi^+ \pi^- \gamma$ , where we approximate the  $\pi^0$  by one energetic photon,  $E_\gamma > 135$  MeV, which carries almost all the energy and retains quite well the direction of its parent  $\pi^0$ . Extensive Monte Carlo studies show that such a reconstruction yields a mass for the  $\omega$  meson which is  $\sim 30$  MeV lower. The detector simulation programs give a clear  $\omega$  signal at a mass of  $\sim 753$  MeV.

The above outlined reconstruction method has been previously used by the CELLO Collaboration in their study of the  $A_2$  meson (9).

Our data show no  $\omega$  signal. Upper limits at the 95% confidence level are given in Fig. 8 along with previously determined upper limits from the JADE Collaboration (15).




Fig. 8. Upper limits (95% conf. level) for the reaction  $\gamma\gamma \rightarrow \omega\omega$ . Expectations of Alexander, Ref. 16 and 17 are shown.

References

- 1) Particle Data Group, Phys. Lett. 111B (1982).
- 2) J.H. Field, Proc. of Int. Europhysics Conf. on H.E. Phys., Brighton (1983) 168.
- 3) Crystal Ball Coll., S. Cooper, 2 $\gamma$  parallel session, Int. Europhysics Conf. on H.E. Phys., Brighton (1983).
- 4) Cronin et al., result reported at  $\gamma\gamma$  Workshop, Paris, (1984)
- 5) J. Olsson, Proc. Fifth Int. Workshop on Photon-Photon Interactions, Aachen 1983, see Ref. 38.
- 6) G. Köpp, T. Walsh, P. Zerwas, Nucl. Phys. B70 (1974) 461.
- 7) W.R. Frazer et al., Phys. Rev. 136 (1964) B1207.
- 8) Crystal Ball Coll., C. Edwards et al., Phys. Lett. 105B (1981) 304.
- 9) CELLO Coll., H.J. Behrend et al., Phys. Lett. 114B (1982) 378, Erratum 125B (1983) 518.
- 10) JADE Coll., J. Olsson, Proc. Fifth Int. Workshop on Photon-Photon Interactions, Aachen 1983.
- 11) TASSO Coll., M. Althoff et al., Phys. Lett. 121B (1983) 216.
- 12) TASSO Coll., R. Brandelik et al., Phys. Lett. 97B (1980) 448.
- 13) D.L. Burke et al, Phys. Lett. 103B (1981) 153.
- 14) CELLO Coll., H.J. Behrend et al., Z. Phys. C21 (1984) 205.
- 15) JADE Coll., J. Olsson contributed paper to Int. Europhysics Conf. on H.E. Physics, Brighton 1983.
- 16) G. Alexander et al., Phys. Rev. D26 (1982) 205.
- 17) G. Alexander et al., private communication.
- 18) H. Kolanoski, Bonn Report HE-84-06, see Ref. 149.
- 19) K. Biswal and S.P. Misra, Phys. Rev. D26 (1982) 3020.
- 20) N.N. Achasov et al., Phys. Lett. 108B (1982); Z. Phys. C16 (1982) 55.
- 21) Bing An Li and K.F. Liu, Phys. Lett. 118B (1982) 435 and Erratum, Phys. Lett. 124B (1983) 550.
- 22) J. Grunhaus, Nevis Report 156 ( 1966).
- 23) CELLO Coll., preliminary results, given at German Physics Society Meeting, Bielefeld 1984.