

DESY 85-010
February 1985



HAVE COLOR OCTET VECTOR BOSONS BEEN FOUND?

by

F. Cornet

Deutsches Elektronen-Synchrotron DESY, Hamburg

and

Dept. de Física Teórica, Univ. Autònoma de Barcelona, Bellaterra, Spain

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX ;
send them to the following address (if possible by air mail) :

DESY
Bibliothek
Notkestrasse 85
2 Hamburg 52
Germany

HAVE COLOR OCTET VECTOR BOSONS BEEN FOUND?

F. Cornet

Deutsches Elektronen-Synchrotron, DESY
Hamburg, GERMANY

and

Departament de Física Teòrica
Universitat Autònoma de Barcelona
BELLATERRA (Barcelona), SPAIN

ABSTRACT: We study the interpretation of the CERN monojet and e jet + missing energy events in terms of a new triplet of color octet vector bosons, \vec{W}_8 . We show that they should also appear in photon-jet final states that will provide a clear signature on the existence of the \vec{W}_8 .

The discovery of the W^\pm , Z at CERN¹⁾ have represented an impressive advance in the experimental test of the standard model. However, the appearance of some unexpected events in the same experiments where the W^\pm , Z have been discovered²⁾ suggests that new physics might also be present. In this sense, models where the W^\pm , Z are composite particles offer an attractive scenario where these strange events could arise in a simple way. In these models, the $SU(2) \times U(1)$ theory of electroweak interactions is nothing but an effective theory describing an interaction remnant of the hypercolor force that bounds the constituents to form a W^\pm or a Z. Until now, however, the problems encountered when building a realistic composite model have been insurmountable. For this reason, the study of the phenomenological implications of the idea of compositeness, without relying on any specific model, is of great interest.

In this note we discuss the recent paper of Gounaris and Nicolaïdis³⁾ where an interpretation of the CERN monojet and e jet+missing energy events is proposed in terms of the production of a new colored vector boson (\vec{W}_8). This particle appears naturally in models where the constituents of the W^\pm , Z carry color. In particular, if we assume that the constituents are color triplets⁴⁾, they can be bounded in color singlet states, corresponding to the usual weak bosons, or in a color octet state corresponding to the new colored weak boson. In the absence of color and electromagnetic interactions the W^\pm , Z and \vec{W}_8 are expected to be degenerate in mass, but color forces split the masses causing the octet to be more massive than the singlet. Using a nonrelativistic potential model one obtains that the mass of the \vec{W}_8 is roughly given by^{5,6)}

$$M(\vec{W}_8) = M(\vec{W}) + \frac{3}{2} \alpha_s \Lambda_h, \quad (1)$$

where α_s is the strong coupling constant and Λ_h is the scale of the hypercolor interaction. Using $\alpha_s \sim 0.1$ and $\Lambda_h \sim 0.5-1 \text{ TeV}$ ⁶⁾ one finds

$$M(\vec{W}_8) = 150 - 230 \text{ GeV}. \quad (2)$$

In order to reproduce the low energy neutral-current phenomenology we assume a global SU(2) symmetry broken by a γ - W^3 mixing (analogous to the γ - ρ^0 mixing that appears in hadronic phenomenology) described by the Lagrangian^{8,9,10)}

$$L_{\gamma W^3} = -\frac{1}{2} \lambda F_{\mu\nu} W_{\mu\nu}^3, \quad (3)$$

where $F_{\mu\nu} (W_{\mu\nu}^3)$ is the photon (W^3) field and λ is a mixing parameter that has to be determined. Using Q^2 -duality¹¹⁾ and assuming that the mass of the first excited state of the \vec{W} is much larger than the mass of the \vec{W} ($M_{W^*}^2 \gg M_W^2$), so that one can safely take into account only the \vec{W} contribution, the mass and coupling constant of the neutral vector boson to fermion pairs are shifted according to

$$M_Z = \frac{M_W}{1-\lambda^2} \quad (4)$$

$$\text{and} \quad g_3 = \frac{g}{\sqrt{1-\lambda^2}} \quad (5)$$

which are exactly the same expressions as those of the standard model if we take

$$\lambda = \sin\theta_W \quad (6)$$

It is interesting to note that, due to the fact that the W^3 - γ and W^3 - g mixings are forbidden by color conservation and the assumed global SU(2) symmetry, respectively, the \vec{W}_8 are degenerate in mass.

The main couplings of the \vec{W}_8 will be to $q\bar{q}$ pairs and to \vec{W}_8 . Following Gounaris and Nicolaïdis³⁾ we shall take for the Lagrangians describing these couplings

$$L_1 = -g_8 \vec{W}_8^{\mu\alpha} \vec{J}_\mu^\alpha \quad (7)$$

$$L_2 = -g_B \vec{W}_8^{\mu\nu} \vec{W}_8^{\alpha\beta} - g_B^{\prime\prime} \epsilon^{\mu\nu\lambda\alpha} \vec{W}_8^{\mu\nu} \vec{W}_8^{\lambda\alpha}, \quad (8)$$



where $\alpha=1, \dots, 8$ is the color index and the quark current \vec{J}_μ^α is given by

$$\vec{J}_\mu^\alpha = a \vec{q}_L^{\dagger} \frac{\lambda_\alpha}{2} \vec{\gamma} \cdot \vec{q}_L, \quad (9)$$

where a is a model dependent constant of order 1. For definiteness we will take in all our numerical calculations $a=\sqrt{6}$ ^{3,6)}.

There is also a coupling $\vec{W}_8 \vec{W}_8 g$, but it will be ignored because it is not relevant for present energy phenomenology.

We can now interpret the CERN monojet and e jet+missing energy events in terms of \vec{W}_8 production by $e\bar{e}$ collisions through the chains



respectively (Fig. 1a). The total cross sections for these processes are given by

$$\sigma_a = \frac{1}{9} \int_0^1 dx \int_0^1 dt [q_i(x) q_j(\tau/x) + (i \leftrightarrow j)] \sigma_a^0(\tau s), \quad (11)$$

where $q_i(x)$ is the distribution function of quarks i in the proton (we have taken the same distribution function for quark i in the proton as for antiquark i in the antiproton), q_a^0 are the cross sections at the parton level and $\tau = \frac{\hat{s}}{s}$ is the ratio between the squared C.M. energy of the colliding quarks to the squared C.M. energy of the colliding protons. In our calculation we have used the distribution functions given by Glück, Hoffmann and Reya¹²⁾, that are valid until very high energies and take into account scaling violation effects.

The evolution of the cross-sections is straightforward using eqs. (7), (8) and (11). The results obtained for $\sqrt{s} = 540$ GeV are

$$\begin{aligned} \sigma(p\bar{p} \rightarrow \vec{W}_8^+ X \rightarrow W^+ g + X) + e^+ \nu g + X &= 1.9 \times 10^{-2} \text{ nb} \\ \sigma(p\bar{p} \rightarrow \vec{W}_8^3 X \rightarrow Zg + X + \bar{W}g + X) &= 1.0 \times 10^{-2} \text{ nb}, \end{aligned} \quad (2)$$

where the kinematical cuts applied by UA1 and UA2²⁾ in their analyses of the data have been considered. We have taken $M_8 = 160$ GeV for the mass of the \vec{W}_8 , as suggested by the data¹³⁾,

and $g^2 = g_B^2 + 4g_W^2$, where g is the standard model weak coupling constant ($g=0.65$), because the hypercolor structure of \tilde{W} and \tilde{W}_0 (in the absence of color forces) are the same. Since the integrated luminosity is $\sim 200 \text{ nb}^{-1}$ (including both experiments), the expected number of events are

$$\begin{aligned} \# \text{ events } (e^{\pm} \text{ jet } \tilde{\nu}_T) &= 3-4 \\ \# \text{ events } (\text{jet } \tilde{\nu}_T) &= 2 \end{aligned} \quad (13)$$

where $\tilde{\nu}_T$ stands for transverse missing energy. It is interesting to note that the dependence of the cross sections (12) on the applied kinematical cuts is very small. So, a release of these cuts would not modify the obtained number of events.

When comparing eq. (13) with the experimental data one sees that the order of magnitude of the expected number of events is in good agreement with the data, although the prediction of a number of e^{\pm} jet+missing energy events larger than the number of monojet events does not seem to be fulfilled by the data. The accumulated statistics, however, is very poor to stress a definite conclusion on this subject. In order to allow for a comparison with the data from the new collider run we have evaluated the cross-section for processes (10) at $\sqrt{s}=630 \text{ GeV}$,

$$\begin{aligned} \sigma(\bar{p}\bar{p} \rightarrow \tilde{W}_0^{\pm} + X \rightarrow W^{\pm} g + X \rightarrow e^{\pm} \nu g + X) &\approx 2.4 \times 10^{-2} \text{ nb} \\ \sigma(\bar{p}\bar{p} \rightarrow \tilde{W}_0^{\pm} + X \rightarrow Z g + X \rightarrow \nu \bar{\nu} g + X) &\approx 1.6 \times 10^{-2} \text{ nb} \end{aligned} \quad (14)$$

and, also, we have plotted the gluon angular distribution in the $\bar{p}\bar{p}$ center of mass system (Fig. 2) for the monojet events.

Further tests of this scenario can be provided by the study of photon-jet final states in $\bar{p}\bar{p}$ collisions. Indeed, when applying the W_3 - γ mixing mechanism to eq. (8) one obtains, in addition to the already discussed $W_3 Z g$ vertex, a $W_3 \gamma g$ interaction given by

$$L_3 = -g_B \lambda_A \lambda_{\mu} \alpha_{\nu}^2 \alpha_{\alpha}^{\mu \nu} - g_B \lambda^{\lambda \epsilon} \mu \nu \lambda_{\alpha}^{\mu} A^{\mu} W_{\alpha}^{\nu} \lambda_{\alpha}^{\alpha} \quad (15)$$

where λ_{μ} is the photon field. Thus, we can calculate the cross-section for the process $\bar{p}\bar{p} \rightarrow \tilde{W}_0^{\pm} + X \rightarrow \gamma g + X$ (Fig. 1b) without introducing any new free parameter. The results turn out to be

$$\begin{aligned} \sigma(\bar{p}\bar{p} \rightarrow \tilde{W}_0^{\pm} + X \rightarrow \gamma g + X) &= 1 \times 10^{-2} \text{ nb} \quad \text{for } \sqrt{s}=540 \text{ GeV} \\ \sigma(\bar{p}\bar{p} \rightarrow \tilde{W}_0^{\pm} + X \rightarrow \gamma g + X) &= 1.4 \times 10^{-2} \text{ nb} \quad \text{for } \sqrt{s}=630 \text{ GeV}. \end{aligned} \quad (16)$$

So, the expected number of photon-jet events is roughly the same as the expected number of monojet events. It must be noted that this is a consequence of the assumed global SU(2) symmetry and its breaking mechanism through a W_3 - γ mixing, and, thus, it will be a common fact to all the models where the weak interaction is considered as a residual interaction and the monojet events are interpreted as \tilde{W} +jet final states, with the \tilde{W} being the decay product of a Z. The W_3 production, however, can be tested using the angular distribution of the γ -jet events, which is predicted to be proportional to $(1-\cos^2\theta)$ in the $\bar{p}\bar{p}$ center of mass system.

In conclusion, the recently discovered CERN monojet and e jet+missing energy events can be considered as a preliminary indication of the existence of new particles, \tilde{W}_0 , that arise in a natural way in a large class of composite models. The low accumulated statistics, however, does not allow to stress any definite conclusion and further tests are needed. This can be achieved by studying the photon-jet events, which are predicted to appear in roughly the same number as the monojet events and would provide a clear W_3 signature.

The author is grateful to R.D. Peccei and K. Hagiwara for many useful discussions and to A. Bramon for critical reading of the manuscript. The author also acknowledges the kind hospitality extended to him by the DESY Theory Group.

FIGURE CAPTIONS

Fig. 1. - Feynman diagrams for

- a) $\bar{p}\bar{p} \rightarrow W_0^+ (W_0^+) + X + W^+ (Z) + X + e^+ \nu(\bar{\nu}) + X$
 b) $\bar{p}\bar{p} \rightarrow W_0^+ (W_0^+) + X + \gamma + X$

Fig. 2. - Gluon angular distribution in the $\bar{p}\bar{p}$ center of mass system for the monojet events.

REFERENCES

- 1.- UA1 Collab., G. Arnison et al., Phys. Lett. 122B, 189 (1983); 126B, 189 (1983).
- UA2 Collab., M. Banner et al., Phys. Lett. 122B, 476 (1983); P. Bagnaia et al., Phys. Lett. 129B, 130 (1983).
- 2.- UA1 Collab., G. Arnison et al., Phys. Lett. 139B (1984). UA2 Collab., P. Bagnaia et al., Phys. Lett. 139B (1984).
- 3.- G.J. Gounaris and A. Nicolaïdis, Phys. Lett. 148B, 239 (1984).
- 4.- H. Fritzsche and G. Mandelbaum, Phys. Lett. 102B, 319 (1981); 109B, 224 (1982).
- 5.- R.D. Peccci, Proceedings of the Workshop on Feasibility of Hadron Colliders in the LEP Tunnel, Lausanne (1984).
- 6.- U. Baur and K.H. Streng, Max Planck Institute Preprint, MPI-PAE/PTH 50/84 (1984).
- 7.- L. Lyons, Proceedings of the International Europhysics Conference on High Energy Physics, Brighton (England), (1983).
- 8.- P. Chen and J.J. Sakurai, Phys. Lett. 110B, 481 (1982).
- 9.- P.Q. Hung and J.J. Sakurai, Nucl. Phys. B143, 81 (1978).
- 10.- J.D. Bjorken, Phys. Rev. D19, 335 (1979).
- 11.- A. Bramon, E. Etim and M. Greco, Phys. Lett. 41B, 609 (1972). J.J. Sakurai, Phys. Lett. 46B, 207 (1973).
- 12.- M. Glück, E. Hoffmann and E. Reya, Z. Physik 13C, 119 (1982).
- 13.- UA1 Collab., CERN Preprint CERN-EP/84-126 (1984).




Fig.1




Fig.2