

**DEUTSCHES ELEKTRONEN-SYNCHROTRON DESY**

DESY 89-049

April 1989



**Path Integral Solution of a Class of Potentials  
Related to the Pöschl-Teller Potential**

Ch. Grosche

*II. Institut für Theoretische Physik, Universität Hamburg*

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

**DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche  
Verwertung der in diesem Bericht enthaltenen Informationen vor.**

**DESY reserves all rights for commercial use of information included in this report, especially in  
case of filing application for or grant of patents.**

To be sure that your preprints are promptly included in the  
**HIGH ENERGY PHYSICS INDEX**,  
send them to the following address ( if possible by air mail ) :

**DESY  
Bibliothek  
Notkestrasse 85  
2 Hamburg 52  
Germany**

## I. INTRODUCTION

In the last ten years there has been remarkable progress in calculating path integrals explicitly. The starting shot was given by Duru and Kleinert [6] with their treatment of the hydrogen atom. The clue for solving this path integral problem successfully was to find simultaneously a coordinate- and time-transformation ("space-time transformation") to reformulate the hydrogen problem in terms of the harmonic oscillator which was a well-known and already solved problem. This idea of attacking path integral problems by performing space-time transformations has since been marvelously fruitful and much of the following work is based on that idea.

Here we want to add some further examples. As we shall see, we must perform in three out of four cases a space-time transformation and we need in all the examples the results of the path integrals of the (modified) Pöschl-Teller potential. Let us summarize in short the results of these two path integrals and the results of the technique of space-time transformations in path integrals:

- 1) The Pöschl-Teller (PT) potential with some numbers  $\kappa$  and  $\lambda$  is defined as

$$V^{PT}(x) = \frac{1}{2m} \left[ \frac{\kappa(\kappa-1)}{\sin^2 x} + \frac{\lambda(\lambda-1)}{\cos^2 x} \right], \quad (0 < x < \frac{\pi}{2}). \quad (1)$$

This class of potentials was first discussed by Pöschl and Teller [26]. A detailed discussion can be e.g. found in Constantinescu and Magyari [4] and Nieto [22]. The path integral solution for this potential can be achieved by means of the path integral over the  $SU(2)$  manifold, and it was successfully studied by Duru [7], Inomata and Kayed [17] and Böhm and Junker [3]. Alternatively, a simple form can be studied with the help of the rigid rotator [11,28].

Some care is needed in the path integral formulation for the Pöschl-Teller potential. Looking carefully at the lattice derivation [3,7] for the path integral we see that we must use a functional measure formulation similar to the one used in the lattice formulation for the radial harmonic oscillator [11,25,29]. This has the consequence that the following interpretation scheme must be used, namely (we use units  $\hbar = 1$ ):

$$\begin{aligned} K^{PT}(x'', x'; T) \\ = \int Dx(t) \exp \left[ i \int_{x'}^{x''} \left( \frac{m}{2} \dot{x}^2 - \frac{\kappa(\kappa-1)}{2m \sin^2 x} - \frac{\lambda(\lambda-1)}{2m \cos^2 x} \right) dt \right] \\ := \int Dx(t) \mu_{\lambda, \kappa}[\sin x, \cos x] \exp \left( \frac{im}{2} \int_{x'}^{x''} \dot{x}^2 dt \right) \\ = \lim_{N \rightarrow \infty} \left( \frac{m}{2\pi i \epsilon} \right)^{\frac{N}{2}} \prod_{j=1}^{N-1} \int_0^{\frac{\pi}{2}} dx^{(j)} \prod_{j=1}^N \mu_{\lambda, \kappa}[\sin x^{(j)}, \cos x^{(j)}] \exp \left[ \frac{im}{2\epsilon} (x^{(j)} - x^{(j-1)})^2 \right], \end{aligned} \quad (2)$$

## Abstract:

In this paper some new exact path integral treatments are presented. These are the path integrals for the symmetric top, the "smooth step", the Rosen-Morse- and the Manning-Rosen-potential. By appropriate space-time transformations these path integral problems can be reduced to the path integral problem of the Pöschl-Teller potential.

\* Supported by Deutsche Forschungsgemeinschaft under contract no. DFG-Mu 757/2-3

where the functional measure  $\mu_{\lambda,\kappa}$  is given by (we use the notation  $\widehat{\theta^{(j)}} \equiv \sin \theta^{(j)} \sin \theta^{(j-1)}$  etc.):

$$\begin{aligned} \mu_{\lambda,\kappa}[\sin x, \cos x] &:= \lim_{N \rightarrow \infty} \prod_{j=1}^N \mu_{\lambda,\kappa}[\sin x^{(j)}, \cos x^{(j)}] \\ &= \lim_{N \rightarrow \infty} \left( \frac{2\pi m}{ie} \right)^N \prod_{j=1}^N \left[ \sin \widehat{x^{(j)}} \cos \widehat{x^{(j)}} \right] \\ &\quad \times \exp \left[ \frac{m}{ie} \left( \sin^2 \widehat{x^{(j)}} + \cos^2 \widehat{x^{(j)}} \right) \right] I_{\lambda-\frac{1}{2}} \left( \frac{m}{ie} \sin^2 \widehat{x^{(j)}} \right) I_{\kappa-\frac{1}{2}} \left( \frac{m}{ie} \cos^2 \widehat{x^{(j)}} \right). \quad (3) \end{aligned}$$

The first line in Eq.(2) has only the symbolical meaning that formally the potential appearing in the Schrödinger equation translates into  $\int Dz \exp(i \times \text{Action})$ . We want to emphasize that only the functional measure formulation has a well-defined lattice formulation. The usual expansion of the modified Bessel function  $I_\nu(z) \simeq (2\pi z)^{-\frac{1}{2}} \exp[z - \frac{\nu^2 - 1/4}{2z}]$  [ $z \rightarrow \infty$ ,  $\arg(z) \neq 0$ ] (or Eq.(3.15) in Ref.[18], respectively) seems very suggestive but gives in the lattice formulation the wrong boundary behaviour of the corresponding short-time kernels and wave functions because the condition  $\arg(z) \neq 0$  is violated. Instead of the correct behaviour we would get a highly singular one. But it is not the scope of this paper to discuss these features in detail; this will be done elsewhere [13].

The path integral solution for the Pöschl-Teller potential now reads:

$$K^{PT}(x'', x'; T) = \sum_{n=0}^{\infty} \exp \left[ -\frac{iT}{2m} (\kappa + \lambda + 2n)^2 \right] \Psi_n^{PT}(x') \Psi_n^{PT}(x''), \quad (4)$$

where

$$\begin{aligned} \Psi_n^{PT}(x) &= \left[ 2(\kappa + \lambda + 2n) \frac{n! \Gamma(\kappa + \lambda + n)}{\Gamma(\kappa + n + \frac{1}{2}) \Gamma(\lambda + n + \frac{1}{2})} \right]^{\frac{1}{2}} \\ &\quad \times (\sin x)^\kappa (\cos x)^\lambda P_n^{(\kappa - \frac{1}{2}, \lambda - \frac{1}{2})} (1 - 2 \sin^2 x). \quad (5) \end{aligned}$$

Here the  $P_n^{(\alpha, \beta)}(x)$  denote Jacobi-polynomials.

2) The modified Pöschl-Teller (mPT) potential with some numbers  $\eta$  and  $\nu$  is defined as

$$V^{mPT}(r) = \frac{1}{2m} \left[ \frac{\eta(\eta - 1)}{\sinh^2 r} - \frac{\nu(\nu - 1)}{\cosh^2 r} \right], \quad (r > 0). \quad (6)$$

A classical study of this problem is due to Frank and Wolf [9], whereas the path integral treatment by means of the path integral over the  $SU(1|1)$  manifold is due to Böhm and Junker [2,3]. The special case  $\nu(\nu - 1) = 0$  can be studied with the help of the path integral on the pseudosphere [12]. Again we must use a functional measure formulation similarly

to the previous one and the following interpretation scheme for the modified Pöschl-Teller potential must be used:

$$\begin{aligned} K^{mPT}(r'', r'; T) &= \int Dr(t) \exp \left[ i \int_{t'}^{t''} \left( \frac{m}{2} \dot{r}^2 - \frac{\eta(\eta - 1)}{2m \sinh^2 x} + \frac{\nu(\nu - 1)}{2m \cosh^2 x} \right) dt \right] \\ &:= \int Dr(t) \mu_{\eta,\nu}[\sinh r, \cosh r] \exp \left( \frac{im}{2} \int_{t'}^{t''} \dot{r}^2 dt \right) \\ &= \lim_{N \rightarrow \infty} \left( \frac{m}{2\pi i\epsilon} \right)^{\frac{N}{2} N-1} \prod_{j=1}^N \int_0^\infty dr^{(j)} \prod_{j=1}^N \mu_{\eta,\nu}[\sinh r^{(j)}, \cosh r^{(j)}] \exp \left[ \frac{im}{2\epsilon} (r^{(j)} - r^{(j-1)})^2 \right], \quad (7) \end{aligned}$$

where the functional measure  $\mu_{\eta,\nu}$  is given by

$$\mu_{\eta,\nu}[\sinh r, \cosh r] = \lim_{N \rightarrow \infty} \prod_{j=1}^N \mu_{\eta,\nu}[\sinh r^{(j)}, \cosh r^{(j)}]$$

$$\begin{aligned} &= \lim_{N \rightarrow \infty} \left( \frac{2\pi m}{\epsilon} \right)^N \prod_{j=1}^N \sinh \widehat{r^{(j)}} \cosh \widehat{r^{(j)}} \\ &\quad \times \exp \left[ -\frac{m}{i\epsilon} (\sinh^2 \widehat{r^{(j)}} - \cosh^2 \widehat{r^{(j)}}) \right] I_{\eta-\frac{1}{2}} \left( \frac{m}{i\epsilon} \sinh^2 \widehat{r^{(j)}} \right) I_{\nu-\frac{1}{2}} \left( \frac{m}{\epsilon} \cosh^2 \widehat{r^{(j)}} \right). \quad (8) \end{aligned}$$

Of course, the same line of reasoning as before is also valid for  $K^{mPT}$  as far as the functional measure is concerned. Adopting the notation of Frank and Wolf the path integral solution reads [define  $2s = \eta(\eta - 1)$ ,  $-2c = \nu(\nu - 1)$  and introduce the numbers  $k_1, k_2$  which are defined in terms of  $c$  and  $s$  as  $k_1 = \frac{1}{2}(1 \pm \sqrt{1/4 + 2s})$ ,  $k_2 = \frac{1}{2}(1 \pm \sqrt{1/4 + 2c})$ ];

$$K^{mPT}(r'', r'; T) = \sum_{n=1}^{N_M} e^{-iTE_n} \Psi_n^{(k_1, k_2)}(r') \Psi_n^{(k_1, k_2)*}(r'')$$

$$+ \int_0^\infty dk e^{-\frac{2\pi}{2m} k^2} \Psi_k^{(k_1, k_2)}(r') \Psi_k^{(k_1, k_2)*}(r''). \quad (9)$$

Here  $N_M$  denotes the maximal number of states with  $0, 1, \dots, n \leq N_M < k_1 - k_2 - \frac{1}{2}$ . The correct signs depend on the boundary conditions for  $r \rightarrow 0$  and  $r \rightarrow \infty$ , respectively. In particular one gets for  $s = 0$  an even and an odd wave function corresponding to  $k_1 = \frac{1}{4}, \frac{3}{4}$ . The bound states are explicitly given by:

$$\begin{aligned} \Psi_n^{(k_1, k_2)}(r) &= N_n^{(k_1, k_2)} (\sinh r)^{2k_1 - \frac{1}{2}} (\cosh r)^{-2k_1 - \frac{1}{2}} \\ &\quad \times {}_2F_1(-k_1 + k_2 + \kappa, -k_1 + k_2 - \kappa + 1; 2k_2; -\sinh^2 r), \quad (10a) \\ &= \left[ \frac{2n! (2k_1 - 1) \Gamma(2k_1 - n - 1)}{\Gamma(2k_2 + n) \Gamma(2k_1 - 2k_2 - n)} \right]^{\frac{1}{2}} (\sinh r)^{2k_2 - \frac{1}{2}} (\cosh r)^{2n - 2k_2 - \frac{1}{2}} \\ &\quad \times P_n^{(2k_2 - 1, 2(k_1 - k_2 - n) - 1)} \left( \frac{1 - \sinh^2 r}{\cosh^2 r} \right), \quad (10b) \\ N_n^{(k_1, k_2)} &= \frac{1}{\Gamma(2k_2)} \left[ \frac{(2k_1 - 1) \Gamma(k_1 + k_2 - \kappa) \Gamma(k_1 + k_2 + \kappa - 1)}{\Gamma(k_1 - k_2 + \kappa) \Gamma(k_1 - k_2 - \kappa - 1)} \right]^{\frac{1}{2}} \quad (10c) \end{aligned}$$

and  $E_n = -\frac{1}{2m} (2\kappa - 1)^2 = -\frac{1}{2m} [2(k_1 - k_2 - n) - 1]^2$ . The continuous states read:

$$\begin{aligned}\Psi_{\mathbf{k}}^{(k_1, k_2)}(r) &= N_{\mathbf{k}}^{(k_1, k_2)}(\cosh r)^{2k_1 - \frac{1}{2}}(\sinh r)^{2k_2 - \frac{1}{2}} \\ &\quad \times {}_2F_1(k_1 + k_2 - \kappa, k_1 + k_2 + \kappa - 1; 2k_2; -\sinh^2 r),\end{aligned}\quad (11a)$$

$$N_{\mathbf{k}}^{(k_1, k_2)} = \frac{1}{\pi \Gamma(2k_2)} \sqrt{\frac{k \sinh \pi k}{2}} \quad (11b)$$

$$\times \left[ \Gamma(k_1 + k_2 - \kappa) \Gamma(-k_1 + k_2 + \kappa) \Gamma(k_1 + k_2 + \kappa - 1) \Gamma(-k_1 + k_2 - \kappa + 1) \right]^{\frac{1}{2}},$$

where  $\kappa = \frac{1}{2}(1 + ik)$  ( $k > 0$ ) and  $E = \frac{k^2}{2m}$ .

3) Let us now discuss the general method for a space-time transformation [11]. It works as follows: One starts with the path integral

$$K(x'', x'; T) = \int Dx(t) \exp \left\{ i \int_{t'}^{t''} \left[ \frac{m}{2} \dot{x}^2 - V(x) - V_{qu}(x) \right] dt \right\} \quad (12)$$

$V_{qu} = \frac{1}{8m}(\Gamma'^2 + 2\Gamma')$  is a quantum potential due to a non-trivial metric, see below, where it is assumed that the effective potential  $V + V_{qu}$  is so complicated that a direct evaluation is not possible. One then defines a new “time”  $s$  together with a coordinate transformation  $x(t) \rightarrow q(s)$ ,  $s(t) = \int_{t'}^t d\sigma / f(x(\sigma))$  and  $x = F(q)$  with some well defined positive functions  $f$  and  $F$ , where we further assume that the relation  $f(F(q)) = [F'(q)]^2$  holds. Let us consider the Legendre-transformed general one-dimensional Hamiltonian

$$H_E = -\frac{1}{2m} \left( \frac{d^2}{dx^2} + \Gamma(x) \frac{d}{dx} \right) + V(x) - E \quad (13)$$

which is hermitian with respect to the inner product  $\langle f_1, f_2 \rangle = \int f_1^*(x) f_2(x) J(x) dx$ , where  $J(x) = e^{\int \Gamma(x) dx}$ . Let us define the quantities  $G(q) = \Gamma(F(q))$ ,  $\sqrt{g} = J(q) = e^{\int \Gamma(q) dq}$ ,  $\tilde{\Gamma}(q) = G(q)F'(q) - F''(q)/F'(q)$  and  $p_q = \frac{1}{i}[\frac{d}{dq} + \frac{1}{2}\tilde{\Gamma}(q)]$ , where  $p_q$  is, of course, a momentum operator which is hermitian with respect to the scalar product  $\langle f_1, f_2 \rangle = \int f_1^*(q) f_2(q) \sqrt{g(q)} dq$ . With the constraint  $f(F(q)) = F'^2(q)$  we thus obtain the space-time transformed Hamiltonian  $\tilde{H} = f H_E$ :

$$\tilde{H}(p_q, q) = \frac{1}{2m} p_q^2 + f(F(q)) [V(F(q)) - E] + \tilde{V}_{qu}(q) \quad (14)$$

with the well-defined quantum correction

$$\tilde{V}_{qu}(q) = \frac{1}{8m} \left[ 3 \left( \frac{F''(q)}{F'(q)} \right)^2 - 2 \frac{F'''(q)}{F'(q)} + (G(q)F'(q))^2 + 2G'(q)F'(q) \right]. \quad (15)$$

Now assume that the constraint  $\int_0^{s''} ds f(F(q(s))) = T [s(t'')] = s''$  has for all admissible paths a unique solution  $s'' \geq 0$ . Of course, since  $T$  is fixed, the “time”  $s''$  will be path-dependent

We must incorporate the constraint into the path integral and define the energy-dependent Feynman-kernel  $G(x'', x'; E)$  (Green's function) via the Fourier transformation

$$K(x'', x'; T) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} e^{-itE} G(x'', x'; E) dE \quad (16)$$

and obtain the transformation formula

$$G(x'', x'; E) = i[f(x'')f(x')]^{1/4} \int_0^{\infty} \tilde{K}(q'', q'; s'') ds'' \quad (17)$$

which gives the energy-dependent kernel  $G$  as a time integral over the transformed Feynman path integral  $\tilde{K}(q'', q; s'') = \langle q'' | e^{-is'' \tilde{H}} | q' \rangle$ :

$$\tilde{K}(q'', q'; s'') = \int Dq(s) \exp \left( i \int_0^{s''} \left\{ \frac{m}{2} \dot{q}^2 - f(F(q)) [V(F(q)) - E] \right\} ds \right) \quad (18)$$

$\{ \dot{q} = dq(s)/ds, q' = F^{-1}(x'), q'' = F^{-1}(x'') \}$ . Here the measure  $Dq(s)$  is defined in the same way as  $Dx(t)$  in the path integral (12). This technique of space-time transformation was originally developed by Duru and Kleinert [6]. It was further developed by Steiner [28], Pak and Sökmén [24], Inomata [16], Kleinert [19] and Grosche and Steiner [11]. The lattice derivation of the equations (16-18) is far from being trivial but can be rigorously performed [e.g. by appropriate symmetrization rules and the well-known midpoint-prescription (which must be used in path integrals on curved manifolds)].

The further content of this paper reads as follows. In section II the path integral solution for the symmetric top is presented, a problem closely related to the  $SU(2)$  path integral.

In section III the path integral solution for the “smooth step” function  $V(x) = -V_0/(1 + e^{x/R})$  ( $x \in \mathbf{R}$ ,  $V_0, R > 0$ , constants) is calculated, a potential important in solid state physics.

Section III contains finally the path integrals for the Rosen-Morse potential  $V(x) = A \tanh \frac{x}{R} - B / \cosh^2 \frac{x}{R}$  ( $x \in \mathbf{R}$ ) and for the Manning-Rosen potential  $V(r) = A \coth \frac{r}{R} - B / \sinh^2 \frac{r}{R}$  ( $r > 0$ ,  $A, B, R > 0$  denote constants).

Section V contains some summarizing remarks.

## II. THE SYMMETRIC TOP

A quantum mechanical treatment of the symmetric top (ST) is due to Demirson [5]. He used it for describing the spectra of molecules with rotational degrees of freedom. There is also a close relationship to the problem of an electron moving in the field of a magnetic monopole [30]. The simplest top is, of course, where all three principle moments of inertia are equal. This, in fact, is the  $SU(2)$  problem [7]. Here we want to discuss the symmetric top, where only two principle momenta of inertia are equal. We introduce the Eulerian angles  $\theta, \psi, \phi$  with  $\theta \in [0, \pi]$ ,  $\psi \in [-2\pi, 2\pi]$  and  $\phi \in [0, 2\pi]$ . The kinetic energy giving the rotational

energy (= classical Hamiltonian = classical Lagrangian) of a rigid body with the two principle moments of inertia  $A$  and  $B$  in these coordinates is then given by (e.g.[20]):

$$\begin{aligned} L_{\text{Cl}} &= \frac{A}{2}(\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2) + \frac{B}{2}(\dot{\phi} + \cos \theta \dot{\psi})^2 \\ &= \frac{A}{2}\dot{\theta}^2 + \frac{1}{2}(A \sin^2 \theta + B \cos^2 \theta)\dot{\phi}^2 + \frac{B}{2}\dot{\phi}^2 + B \cos \theta \dot{\phi} \dot{\psi} \\ &= \frac{A}{2}g_{ab}q^a q^b, \end{aligned} \quad (19)$$

where  $q^a$  ( $a = 1, 2, 3$ ) denote the three Eulerian angles and the metric tensor  $g_{ab}$  and its inverse  $g^{ab}$  are given by ( $\tilde{B} = B/A$ ):

$$(g_{ab}) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sin^2 \theta + \tilde{B} \cos^2 \theta & \tilde{B} \cos \theta \\ 0 & \tilde{B} \cos \theta & \tilde{B} \end{pmatrix}, \quad (g^{ab}) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sin^2 \theta} & -\frac{\cos \theta}{\sin^2 \theta} \\ 0 & -\frac{\cos \theta}{\sin^2 \theta} & \frac{1}{\tilde{B}} + \cot^2 \theta \end{pmatrix} \quad (20)$$

and the determinant  $g$  of  $g_{ab}$  reads:  $g = \det(g_{ab}) = \tilde{B} \sin^2 \theta$ . The Hamiltonian reads as:

$$H_{\text{Cl}} = \frac{1}{2A} \left[ p_\theta^2 + \frac{p_\phi^2}{\sin^2 \theta} + \left( \frac{A}{B} + \cot^2 \theta \right) p_\phi^2 - \frac{2 \cos \theta}{\sin^2 \theta} p_\psi p_\phi \right]. \quad (21)$$

To construct Hamiltonians in quantum mechanics we must respect the ordering ambiguity of position- and momentum-operators. Following our prescriptions given in [10,11] we start by considering the hermitian momenta  $p_q = \frac{1}{i}(\frac{\partial}{\partial \theta} + \frac{1}{2}\Gamma_q)$  with  $\Gamma_q = \partial_q \ln \sqrt{g}$ . Thus

$$p_\theta = \frac{1}{i} \left( \frac{\partial}{\partial \theta} + \frac{1}{2} \cot \theta \right), \quad p_\psi = \frac{1}{i} \frac{\partial}{\partial \psi}, \quad p_\phi = \frac{1}{i} \frac{\partial}{\partial \phi}. \quad (22)$$

For the Hamiltonian we can use the well-known Weyl-ordering prescription

$$H = \frac{1}{8A} [g^{ab} p_a p_b + 2p_a g^{ab} p_b + p_a p_b g^{ab}] + \Delta V_{Weyl} \quad (23)$$

with the well-defined quantum potential

$$\Delta V_{Weyl} = \frac{1}{8A} [g^{ab} \Gamma_a \Gamma_b + 2(g^{ab} \Gamma_b)_{,a} + g^{ab}_{,ab}], \quad (24)$$

or, respectively, the “product-ordering” prescription [10]

$$H = \frac{1}{2A} h^{ac} p_a p_b h^{cb} + \Delta V_{Weyl} + \frac{1}{8A} (2h^{ac} h^{bc}_{,ab} - h^{ac}_{,a} h^{bc}_{,b} - h^{ac}_{,b} h^{bc}_{,a}) \quad (25)$$

with the decomposition  $g^{ab} = h^{ac} h^{cb}$ . Due to the special nature of  $g^{ab}$  and  $g_{ab}$ , respectively, the quantum potentials turn out to be equal for both prescriptions and we find for the quantum Hamiltonian of the symmetric top:

$$\begin{aligned} H &= \frac{1}{2A} \left[ \frac{\partial^2}{\partial \theta^2} + \cot \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} + \left( \frac{A}{B} + \cot^2 \theta \right) \frac{\partial^2}{\partial \psi^2} - 2 \frac{\cos \theta}{\sin^2 \theta} \frac{\partial}{\partial \psi} \frac{\partial}{\partial \phi} \right] \\ &= \frac{1}{2A} \left[ p_\theta^2 + \frac{p_\phi^2}{\sin^2 \theta} + \left( \frac{A}{B} + \cot^2 \theta \right) p_\phi^2 - 2 \frac{\cos \theta}{\sin^2 \theta} p_\psi p_\phi \right] - \frac{1}{8A} \left( 1 + \frac{1}{\sin^2 \theta} \right). \end{aligned} \quad (26)$$

The path integral for the symmetric top can now be constructed in the usual way [10,11] and it reads in the “product-form” definition  $[\Delta q^{(j)} = q^{(j)} - q^{(j-1)}, q^{(j)} = q(t^{(j)}) = q(t' + j\epsilon)]$ :  $0, 1, \dots, N$ ,  $\epsilon = T/N = (t'' - t')/N$  ( $N \rightarrow \infty$ ):

$$\begin{aligned} K^{ST}(\theta'', \theta', \psi'', \psi', \phi'', \phi'; T) &= \int \sqrt{g} D\theta(t) D\psi(t) D\phi(t) \\ &\times \exp \left\{ i \int_t^{t''} \left[ \frac{A}{2} \dot{\theta}^2 + \frac{1}{2} (A \sin^2 \theta + B \cos^2 \theta) \dot{\psi}^2 + \frac{B}{2} \dot{\phi}^2 + B \cos \theta \dot{\psi} \dot{\phi} + \frac{1}{8A} \left( 1 + \frac{1}{\sin^2 \theta} \right) \right] dt \right\} \\ &= \lim_{N \rightarrow \infty} \left( \frac{A}{2\pi i \epsilon} \right)^{\frac{3}{2}N N^{-1}} \prod_{j=1}^N \sqrt{\frac{A}{B}} \int_0^\pi \sin \theta^{(j)} d\theta^{(j)} \int_{-2\pi}^{2\pi} d\psi^{(j)} \\ &\times \exp \left\{ i \sum_{j=1}^N \left[ \frac{A}{2\epsilon} \Delta^2 \theta^{(j)} + \frac{1}{2\epsilon} (A \sin^2 \theta^{(j)} + B \cos^2 \theta^{(j)}) \Delta^2 \psi^{(j)} + \frac{B}{2\epsilon} \Delta^2 \phi^{(j)} \right. \right. \\ &\quad \left. \left. + \frac{B}{\epsilon} \widetilde{\cos \theta^{(j)}} \Delta \psi^{(j)} \Delta \phi^{(j)} + \frac{\epsilon}{8A} \left( 1 + \frac{1}{\sin^2 \theta^{(j)}} \right) \right] \right\}. \end{aligned} \quad (27)$$

We calculate this path integral by starting with a Fourier expansion:

$$\begin{aligned} K^{ST}(\theta'', \theta', \psi'', \psi', \phi'', \phi'; T) &= \sum_{L=-\infty}^{\infty} \sum_{M=-\infty}^{\infty} K_{LM}^{ST}(\theta'', \theta', T) e^{iL(\phi'' - \phi')} e^{iM(\psi'' - \psi')} \\ K_{LM}^{ST}(\theta'', \theta'; T) &= \frac{1}{8\pi^2} \int_{-2\pi}^{2\pi} d\phi'' K^{ST}(\theta'', \theta', \psi'', \psi', \phi'', \phi'; T) e^{-iL(\phi'' - \phi') - iM(\psi'' - \psi')}. \end{aligned}$$

This yields

$$\begin{aligned} K_{LM}^{ST}(\theta'', \theta'; T) &= \frac{1}{8\pi^2} \exp \left( \frac{iT}{8A} \right) \\ &\times \lim_{N \rightarrow \infty} \left( \frac{A}{2\pi i \epsilon} \right)^{\frac{3}{2}N N^{-1}} \prod_{j=1}^N \sqrt{\frac{A}{B}} \int_0^\pi \sin \theta^{(j)} d\theta^{(j)} \exp \left[ i \sum_{j=1}^N \left( \frac{A}{2\epsilon} \Delta^2 \theta^{(j)} + \frac{\epsilon}{8A \sin^2 \theta^{(j)}} \right) \right] \\ &\times \prod_{j=1}^N \int_{-2\pi}^{2\pi} d\phi^{(j)} \exp \left\{ \frac{i}{2\epsilon} \left[ (A \sin^2 \theta^{(j)} + B \cos^2 \theta^{(j)}) \Delta^2 \psi^{(j)} + B \Delta^2 \phi^{(j)} \right. \right. \\ &\quad \left. \left. + 2B \cos \theta^{(j)} \Delta \psi^{(j)} \Delta \phi^{(j)} \right] - iL \Delta \phi^{(j)} - iM \Delta \psi^{(j)} \right\} \\ &= \frac{1}{8\pi^2} \sqrt{\frac{A}{B}} \exp \left[ -iT \left( \frac{L^2}{2B} - \frac{1}{8A} \right) \right] \\ &\times \lim_{N \rightarrow \infty} \left( \frac{A}{2\pi i \epsilon} \right)^{N N^{-1}} \prod_{j=1}^N \int_0^\pi \sin \theta^{(j)} d\theta^{(j)} \exp \left[ i \sum_{j=1}^N \left( \frac{A}{2\epsilon} \Delta^2 \theta^{(j)} + \frac{\epsilon}{8A \sin^2 \theta^{(j)}} \right) \right] \\ &\times \prod_{j=1}^N \int_{-2\pi}^{2\pi} d\psi^{(j)} \exp \left[ \frac{iA}{2\epsilon} \sin^2 \theta^{(j)} \Delta \psi^{(j)} \right] \\ &= \sqrt{\frac{A}{B}} \frac{(\sin \theta' \sin \theta'')^{-\frac{1}{2}}}{8\pi^2} \exp \left[ -iT \left( \frac{L^2}{2B} - \frac{1}{8A} \right) \right] \hat{K}_{LM}^{ST}(\theta'', \theta'; T), \end{aligned} \quad (28)$$

where the kernel  $\hat{K}_{LM}^{ST}$  is given by:

$$\begin{aligned}\hat{K}_{LM}^{ST}(\theta'', \theta'; T) &= \int D\theta(t) \exp \left[ i \int_{t'}^{t''} \left( \frac{A}{2} \dot{\theta}^2 - \frac{(M - L \cos \theta)^2 - \frac{1}{4}}{2A \sin^2 \theta} \right) dt \right] \\ &= \lim_{N \rightarrow \infty} \left( \frac{A}{2\pi i \epsilon} \right)^{\frac{N}{2}} \prod_{j=1}^{N-1} \int_0^\pi d\theta^{(j)} \exp \left[ i \sum_{j=1}^N \left( \frac{A}{2\epsilon} \Delta^2 \theta^{(j)} - \epsilon \frac{(M - L \cos \theta^{(j)})^2 - \frac{1}{4}}{2A \sin^2 \theta^{(j)}} \right) \right].\end{aligned}\quad (30)$$

This potential problem related to the symmetric top has also its own right as a path integral, especially if one analytically continues in  $L$  and  $M$  to, say, arbitrary real numbers. The path integral (30) can now be solved by a simple coordinate transformation by means of the Pöschl-Teller potential or  $SU(2)$  path integral (see also [7] for a similar calculation), respectively. We consider the Hamiltonian

$$H = -\frac{1}{2A} \frac{d^2}{d\theta^2} + \frac{(M - L \cos \theta)^2 - \frac{1}{4}}{2A \sin^2 \theta}, \quad (31)$$

perform the transformation  $(1 - \cos \theta) = 2 \sin^2 x$ , i.e.  $x = \theta/2$ , and arrive at the transformed Hamiltonian

$$\hat{H} = -\frac{1}{8A} \frac{d^2}{dx^2} + \frac{(M - L)^2 - \frac{1}{4}}{8A \sin^2 x} + \frac{(M + L)^2 - \frac{1}{4}}{8A \cos^2 x} - \frac{L^2}{2A}. \quad (32)$$

Thus we get the transformed path integral corresponding to the Hamiltonian  $\hat{H}$ :

$$\begin{aligned}\hat{K}_{LM}^{ST}(\theta'', \theta'; T) &\equiv \frac{1}{2} e^{iT L^2 / 2A} \hat{K}(x'', x'; T) \\ &= \frac{1}{2} e^{iT L^2 / 2A} \int Dx(t) \mu_{|M-L|, |M+L|} [\sin x, \cos x] \exp \left( 2iA \int_{t'}^{t''} \dot{x}^2 dt \right) \\ &\quad \times \sum_{n=0}^{\infty} \exp \left[ -iT \left( \frac{(d+s+2n+1)^2}{8A} - \frac{L^2}{2A} \right) \right] \Psi_n^{PT}(x') \Psi_n^{PT}(x'').\end{aligned}\quad (33)$$

Here we used the notation of Eq.(4), where  $m = 4A$ ,  $\kappa = d + \frac{1}{2}$ ,  $\lambda = s + \frac{1}{2}$  with  $d = |M - L|$  and  $s = |M + L|$ . Thus we get finally the path integral for the symmetric top:

$$K^{ST}(\theta'', \theta', \psi'', \psi', \phi'', \phi'; T) = \sum_{n=0}^{\infty} \sum_{M=-\infty}^{\infty} \sum_{L=-\infty}^{\infty} e^{-iT E_n^{ST}} \Psi_n^{ST}(\theta', \psi', \phi') \Psi_n^{ST*}(\theta'', \psi'', \phi''), \quad (34)$$

where the wave-functions and the energy-spectrum are given by:

$$\Psi_n^{ST}(\theta, \psi, \phi) = N_n^{ST} e^{-iL\phi} e^{-iM\psi} e^{-iM\phi} (1 - \cos \theta)^{\frac{|M-L|}{2}} (1 + \cos \theta)^{\frac{|M+L|}{2}} P_n^{(|M-L|, |M+L|)}(\cos \theta) \quad (35)$$

$$E_n^{ST} = \frac{(n + \frac{d+s+1}{2})^2 - \frac{1}{4}}{2A} + \frac{L^2}{2} \left( \frac{1}{B} - \frac{1}{A} \right). \quad (36)$$

We can introduce the quantity  $J = n + \frac{d+s}{2} = 0, 1, 2, \dots$  and can interpret the energy as

$$\begin{aligned}E^{ST} &= \frac{J(J+1)}{2A} + \frac{L^2}{2} \left( \frac{1}{B} - \frac{1}{A} \right). \quad (37) \\ \text{For } A = B \text{ we recover the } SU(2)\text{- and } SO(3)\text{-spectrum, respectively. Note that}\end{aligned}$$

$$\begin{aligned}\frac{d+s}{2} &= \frac{1}{2} |L + M| + \frac{1}{2} |L - M| = |L|, & (|L| \geq |M|) \\ &= |M|, & (|M| \geq |L|).\end{aligned}\quad (38)$$

This suggests that  $J$ ,  $L$  and  $M$  may be interpreted as representing angular momentum and it can be in fact shown that, e.g. the total angular momentum  $P$  can be described by  $P^2 = J(J+1)$ . For a detailed discussion of these features see [5].

As a final result we consider the kernel  $\hat{K}_{LM}^{ST}$ , interpret the numbers  $L$  and  $M$  as, say, arbitrary real numbers and obtain the path integral identity:

$$\begin{aligned}\int D\theta(t) \exp \left[ i \int_{t'}^{t''} \left( \frac{A}{2} \dot{\theta}^2 - \frac{(M - L \cos \theta)^2 - \frac{1}{4}}{2A \sin^2 \theta} \right) dt \right] \\ = \sum_{n=0}^{\infty} \exp \left[ -\frac{iT}{2A} \left( \frac{(2n+d+s+1)^2}{4} - L^2 \right) \right] \Psi_n^{PT}(\cos \theta) \Psi_n^{PT}(\cos \theta'') \quad (39)\end{aligned}$$

in the notation of Eqs.(4,5,34). In Eq.(39) a functional measure formulation must be used if needed.

### III. THE SMOOTH STEP

Let us consider now the path integral for the “smooth step” potential ( $x \in \mathbb{R}$ )  $V(x) = -V_0/(1 + e^{x/R})$ :

$$K(x'', x'; T) = \int Dx(t) \exp \left\{ i \int_{t'}^{t''} \left[ \frac{m}{2} \dot{x}^2 + \frac{V_0}{1 + e^{\frac{x}{R}}} \right] dt \right\}. \quad (40)$$

Potentials like this are important in solid state physics. They can smear out the potential step at metal-vacuum interfaces. Following our prescription for a general space-time transformation in path integrals we start by considering the Legendre-transformed Hamiltonian  $H_E$ :

$$H_E = -\frac{1}{2m} \frac{d^2}{dx^2} - \frac{V_0}{1 + e^{\frac{x}{R}}} - E. \quad (41)$$

We perform the transformation  $(1 + e^{\frac{x}{R}})^{-1} = \tanh^2 r$ , introduce the momentum operator  $p_r = \frac{1}{i} (\frac{d}{dr} + \frac{1}{2} \Gamma_r)$  [ $\Gamma_r = 1/(\sinh r \cosh r)$ ] and define a new “time”  $s$  by  $dt = f[q(s)]ds$ , with

$f(r) = 4R^2 \coth^2 r$ . Thus we have  $\Delta V = -\frac{1}{8m}(3/\cosh^2 r + 1/\sinh^2 r)$  and arrive at the space-time transformed Hamiltonian  $\tilde{H} = f(r)\hat{H}_E$ :

$$\tilde{H} = \frac{p_r^2}{2m} - 4R^2(E + V_0) - \frac{8mR^2E + \frac{1}{4}}{2m\sinh^2 r} - \frac{3}{8m\cosh^2 r}. \quad (42)$$

Therefore the space-time transformed path integral  $\tilde{K}$  reads  $[\nu = -\frac{1}{2}, \frac{3}{2}; \eta = \frac{1}{2} \pm \sqrt{-8mR^2E}]$ :

$$\tilde{K}(r'', r'; s'') = e^{is''4R^2(E+V_0)} \int D\tau(s) \mu_{n,\nu} [\sinh r, \cosh r] \exp\left(\frac{im}{2} \int_{r'}^{r''} r'^2 ds\right) \quad (43)$$

This path integral is a special case of the modified Pöschl-Teller potential as discussed in the introduction. Equations (9) and (11) give immediately the solution yielding (there are no bound states)

$$\tilde{K}(r'', r'; s'') = e^{is''4R^2(E+V_0)} \int_0^\infty dk e^{-\frac{i\pi}{2m}k^2} \Psi_k^{(k_1, k_2)}(r') \Psi_k^{(k_1, k_2)}(r''), \quad (44)$$

Due to the boundary conditions  $\Psi_k^{(k_1, k_2)} \rightarrow 0$  for  $r \rightarrow 0$  we get  $k_1 = 0$  and  $k_2 = \frac{1}{2}(1 + \sqrt{-8mR^2E})$ . Performing the  $s''$ -integration gives therefore

$$G(x'', x'; E) = \int_0^\infty dk \frac{\Psi_{Step}(x') \Psi_{Step}(x'')}{\frac{k^2}{8mR^2} - V_0 - E}. \quad (45)$$

The energy-spectrum reads  $E_k = \frac{k^2}{8mR^2} - V_0$  and has a constant shift  $V_0$ . The wave functions are given by  $[y \equiv (1 + e^{\frac{x}{R}})^{-1}, p \equiv \sqrt{2mER^2}, k \equiv \sqrt{2mR^2(E + V_0)}]$ :

$$\begin{aligned} \Psi_{Step}(x) &= Ny^{ip}(1-y)^{ik} {}_2F_1[i(p+k), i(p+k)+1; 1+2p; y] \\ N &= \left[ \frac{k}{8\pi R} \frac{\sinh \pi(p+k) \sinh 2\pi p}{\sinh \pi(p+k) \sinh \pi(p-k)} \right]^{\frac{1}{2}}. \end{aligned} \quad (46)$$

This is the correct solution [4].

#### IV. THE ROSEN-MORSE AND MANNING-ROSEN POTENTIAL

- 1) Let us first consider the path integral for the Rosen-Morse (RM) potential  $V^{RM}(x) = A \tanh \frac{x}{R} - B / \cosh^2 \frac{x}{R}$ , ( $x \in \mathbf{R}$ ,  $A, B, R$  positive constants):

$$K^{RM}\{x'', x'; T\} = \int D\tau(t) \exp \left[ i \int_{r'}^{r''} \left( \frac{m}{2} \dot{x}^2 - A \tanh \frac{x}{R} + \frac{B}{\cosh^2 \frac{x}{R}} \right) dt \right]. \quad (47)$$

The potential  $V^{RM}$  was introduced by Rosen and Morse [27] to discuss the spectra of polyatomic molecules. With this potential one can also study the penetration of electrons through

a potential barrier [8]. The Rosen-Morse potential has a discrete and continuous spectrum and thus a “hidden”  $SU(1,1)$  symmetry. There is already a path integral solution for  $V^{RM}$  by Junker and Ihonata [18], but their treatment is based on the  $SU(2)$  path integral instead of the  $SU(1,1)$  path integral. Therefore these authors did not get the continuous part of the spectrum.

We proceed similarly to the previous section. Let us start with the Legendre transformed Hamiltonian  $H_E$ :

$$H_E = -\frac{1}{2m} \frac{d^2}{dx^2} + A \tanh \frac{x}{R} - \frac{B}{\cosh^2 \frac{x}{R}} - E. \quad (48)$$

We perform the transformation  $\frac{1}{2}(1 + \tanh \frac{x}{R}) = \tanh^2 r$ , introduce the momentum operator  $p_r = \frac{1}{i}(\frac{d}{dr} + \frac{1}{2}\Gamma_r)$  with  $\Gamma_r = 1/(\sinh r \cosh r)$ . We define a “time”  $s$  together with the coordinate transformation  $x(t) \rightarrow r(s)$  with  $F(r) = R \operatorname{artanh}(2 \tanh^2 r - 1)$ ,  $f(r) = R^2 \coth^2 r$  and  $\Delta V = -\frac{1}{8m}(3/\cosh^2 r + 1/\sinh^2 r)$ . Thus we get the space-time transformed Hamiltonian  $\tilde{H} = f(r)H_E$ :

$$\tilde{H} = \frac{p_r^2}{2m} - \frac{2mR^2(A+E) + \frac{1}{4}}{2m \sinh^2 r} - \frac{8mBR^2 + \frac{3}{4}}{2m \cosh^2 r} + R^2(A-E) \quad (49)$$

and the space-time transformed path integral  $\tilde{K}$  reads  $[\eta = \frac{1}{2} \pm \sqrt{-2mR^2(A+E)}, \nu = \frac{1}{2} \pm \sqrt{1 + 8mBR^2}]$ :

$$\begin{aligned} \tilde{K}(r'', r'; s'') &= e^{is''R^2(A-E)} \int Dr(s) \mu_{n,\nu} [\sinh r, \cosh r] \exp \left( \frac{im}{2} \int_0^{s''} r'^2 ds \right) \\ &= \sum_{n=0}^{N_M} e^{is''[(E-A)R^2 - E_n]} \Psi_n^{(k_1, k_2)}(r') \Psi_n^{(k_1, k_2)}(r'') \\ &\quad + \int_0^\infty dk e^{is''[(E-A)R^2 - \frac{k^2}{2m}]} \Psi_k^{(k_1, k_2)}(r') \Psi_k^{(k_1, k_2)}(r''). \end{aligned} \quad (50)$$

in the notation of Eq.(9). Performing the  $s''$ -integration [c.f.Eq.(17)] and respecting the correct boundary conditions for  $x \rightarrow \pm\infty$  yields:

$$G(x'', x'; E) = \sum_{n=0}^{N_M} \frac{\Psi_n^{RM}(x') \Psi_n^{RM}(x'')}{E_n^{RM} - E} + \int_0^\infty dk \frac{\Psi_k^{RM}(x') \Psi_k^{RM}(x'')}{A + \frac{k^2}{2mR^2} - E}. \quad (51)$$

The wave-functions and the energy spectrum are given by  $[s \equiv \sqrt{1 + 8mBR^2}, 0, 1, \dots, n \leq N_M < \frac{1}{2}(s-1) - \sqrt{m|A|R^2}, k_1 = \frac{1}{2}(1+s), k_2 = \frac{1}{2}[1 + \frac{1}{2}(s-2n-1) - \frac{2nAR^2}{s-2n-1}], u = \frac{1}{2}(1 + \tanh \frac{x}{R})$ , note  $k_2 - \frac{1}{2} > 0$ :

$$\begin{aligned} \Psi_n^{RM} &= \frac{N_n^{(k_1, k_2)}}{\sqrt{R}} u^{k_2 - \frac{1}{2}} (1-u)^{\frac{1}{2}s-k_2-n} {}_2F_1(-n, s-n; 2k_2; u) \\ \Psi_n^{RM} &= \frac{N_n^{(k_1, k_2)}}{\sqrt{R}} u^{k_2 - \frac{1}{2}} (1-u)^{\frac{1}{2}s-k_2-n} {}_2F_1(-n, s-n; 2k_2; u) \end{aligned} \quad (52a)$$

$$\Psi_n^{RM} = \left[ \frac{s \cdot n! \Gamma(s-n)}{R \cdot 2^{s-2n-1} \Gamma(2k_2-n) \Gamma(s-2k_2-n+1)} \right]^{\frac{1}{2}}$$

$$x \times (1 - \tanh \frac{x}{R})^{\frac{1}{2} \cdot k_2 - n} (1 + \tanh \frac{x}{R})^{k_2 - \frac{1}{2}} P_n^{(\star - 2k_2 - 2n, 2k_2 - 1)} (\tanh \frac{x}{R}) \quad (52b)$$

$$E_n^{RM} = - \left[ \frac{(s-2n-1)^2}{8mR^2} + \frac{2mA^2R^2}{(s-2n-1)^2} \right]. \quad (53)$$

The wave-functions and the energy spectrum of the continuous states are given by [k<sub>2</sub> ≈ 1/2(1 + ik̄), k̄ ≡ √2mR<sup>2</sup>(A + E<sub>k</sub>) > 0]:

$$\Psi_k^{RM}(x) = \frac{N_k^{(k_1, k_2)}}{\sqrt{R}} (1-u)^{-\frac{1}{2}k} u^{\frac{1}{2}k} {}_2F_1\left\{\frac{1}{2}[1+s+i(\tilde{k}-k)], \frac{1}{2}[1-s+i(\tilde{k}-k)]; 1+i\tilde{k}; u\right\}$$

$$E_k = A + \frac{k^2}{2mR^2}. \quad (54)$$

2) Second we discuss the Manning-Rosen (MR) potential V<sub>MR</sub>(x) = B / sinh<sup>2</sup>  $\frac{x}{R}$  - A coth  $\frac{x}{R}$ , (x > 0, A, B, R positive constants). V<sub>MR</sub> was introduced by Manning and Rosen [21] to study vibrations of diatomic molecules. It can furthermore be used for describing the Kepler problem in a space of constant negative curvature [15]. The path integral for the Kepler problem in a space of constant positive curvature was discussed by Barut, Inomata and Junker [1]. The path integral for V<sub>MR</sub> reads:

$$K^{MR}(x'', x'; t') = \int Dx(t) \exp \left[ i \int_{t'}^{t''} \left( \frac{m}{2} \dot{x}^2 + A \coth \frac{x}{R} - \frac{B}{\sinh^2 \frac{x}{R}} \right) dt \right]. \quad (55)$$

Again we consider the Legendre-transformed Hamiltonian H<sub>E</sub>:

$$H_E = -\frac{1}{2m} \frac{d^2}{dx^2} - A \coth \frac{x}{R} + \frac{B}{\sinh^2 \frac{x}{R}} - E. \quad (56)$$

Performing the transformation  $\frac{1}{2}(1 - \coth \frac{x}{R}) = -1/\sinh^2 r$  yields:

$$\tilde{H} = \frac{p_r^2}{2m} + \frac{2mR^2(E-A)}{2m \cosh^2 r} + \frac{1}{4} + \frac{8mBR^2 + \frac{3}{4}}{2m \sinh^2 r} - (E+A)R^2.$$

Here we have introduced the momentum operator p<sub>r</sub> =  $\frac{1}{i}(\frac{d}{dr} + \frac{1}{2}\Gamma_r)$  [Γ<sub>r</sub> = -1/(sinh r cosh r)] together with the time transformation dt = f(r)ds, where f(r) = R<sup>2</sup> tanh<sup>2</sup> r. With ΔV =  $\frac{1}{8m}(1/\cosh^2 r + 3/\sinh^2 r)$  we thus arrive at the space-time transformed path integral [η =  $\frac{1}{2} \pm \sqrt{1 + 8mBR^2}$ , ν =  $\frac{1}{2} \pm \sqrt{-2mR^2(E-A)}$ ]:

$$\tilde{K}(r'', r; s'')$$

$$= e^{is''(E+A)R^2} \int D\Gamma(s) \mu_{\eta, \nu} [\sinh r, \cosh r] \exp \left( i \int_0^{s''} \frac{m}{2} \dot{r}^2 ds \right)$$

$$= \sum_{n=0}^{N_M} e^{is''[(E+A)R^2 - E_n]} \Psi_n^{(k_1, k_2)}(r') \Psi_n^{(k_1, k_2)*}(r'')$$

$$+ \int_0^{\infty} dk e^{is''[(E+A)R^2 - \frac{k^2}{2m}]} \Psi_n^{(k_1, k_2)}(r') \Psi_n^{(k_1, k_2)*}(r''), \quad (58)$$

again with the notation of Eq.(9). Performing the s''-integration [c.f.Eq.(17)] yields:

$$G(x'', x'; E) = \sum_{n=0}^{N_M} \frac{\Psi_n^{MR}(x') \Psi_n^{MR}(x'')}{E_n^{MR} - E} + \int_0^{\infty} dk \frac{\Psi_k^{MR}(x') \Psi_k^{MR*}(x'')}{\frac{k^2}{2mR^2} - A - E}. \quad (59)$$

Respecting the correct boundary conditions for x → 0 and x → ∞ gives k<sub>1</sub> =  $\frac{1}{2}[(1+\frac{1}{2}(s+2n+1)) + \frac{2mAR^2}{s+2n+1}]$  and k<sub>2</sub> =  $\frac{1}{2}(1 + \sqrt{1 + 8mBR^2}) \equiv \frac{1}{2}(1+s)$ . The wave-functions and the energy-spectrum of the bound states read [0, 1, …, n ≤ N<sub>M</sub> < √mAR<sup>2</sup> -  $\frac{1}{2}(s+1)$ , note n +  $\frac{1}{2}$  - k<sub>1</sub> < 0]:

$$\Psi_n^{MR}(x) = \frac{N_n^{(k_1, k_2)}}{\sqrt{R}} (u-1)^{\frac{1}{2}-k_1+n} u^{k_1-1-\frac{1}{2}s-n} {}_2F_1\left(-n, 2k_1 - n - 1; s + 1; \frac{1}{1-u}\right)$$

$$= \left[ \frac{(2k_1-1)n! \Gamma(2k_1-n-1)}{R \Gamma(n+s+1) \Gamma(2k_1-s-n-1)} \right]^{\frac{1}{2}}$$

$$\times (1 + e^{-2x})^{k_2} e^{-2x(k_1-\frac{1}{2}s-n-1)} P_n^{(2k_1-2n-s-2,s)}(1-2e^{-2x}) \quad (60)$$

$$E_n^{RM} = - \left[ \frac{(s+2n+1)^2}{8mR^2} + \frac{2mA^2R^2}{(s+2n+1)^2} \right]. \quad (61)$$

The wave-functions and the energy-spectrum of the continuous states are given by [k<sub>1</sub> ≡  $\frac{1}{2}(1+i\tilde{k})$ ,  $\tilde{k} \equiv \sqrt{2mR^2(E_k - A)} > 0$ ]:

$$\Psi_k^{MR}(x) = \frac{N_k^{(k_1, k_2)}}{\sqrt{R}} u^{-\frac{1}{2}k} (u-1)^{\frac{1}{2}i\tilde{k}-(1+s)} \\ \times {}_2F_1\left(\frac{1+s+i(\tilde{k}-k)}{2}, \frac{1+s-i(\tilde{k}+k)}{2}; s+1; \frac{1}{1-u}\right) \quad (62)$$

## V.SUMMARY

In this paper several further examples of exact path integral treatments have been presented. These have been the path integral for the “smooth step”, the Rosen-Morse-, the Manning-Rosen potential and the symmetric top. All examples have a close relation to the usual Pöschl-Teller potential (for a pure discrete spectrum as for the top) or to the modified Pöschl-Teller potential (if the spectrum is also continuous). As already noted these two manifolds and potential problems. But, following Duru [7], “it is well-known that all of the special functions which appear as solutions of problems in theoretical physics are the matrix elements of some Lie group. Because of this fact, if one parametrizes the problems

suitably with their symmetries, it may be possible to relate their path integrals to the ones written for the motions on the appropriate group spaces.”

We have also discussed the problem of the correct functional measure to be used in the path integral of the the Pöschl-Teller and modified Pöschl-Teller case, respectively. The usual expansion of the modified Bessel function seems very suggestive but gives in the lattice formulation the wrong boundary behaviour of the corresponding short-time kernels and wave-functions. This is very analogous to the radial path integral which has been discussed in great detail in Refs.[11,29].

The examples solved in this paper suggest that the two Pöschl-Teller path integrals play an analogous important role for applications as the path integral for the radial harmonic oscillator [25]. In a forthcoming publication we use these three path integrals (Pöschl-Teller, modified Pöschl-Teller and radial harmonic oscillator) to give a classification scheme (similar to the classical factorization method of Schrödinger on the operator level as reviewed by Infeld and Hull [14]) of the known path integral solutions [13].

**Acknowledgement:** I thank F.Steiner for reading the manuscript. I also thank F.Steiner and W.Nachmann for drawing my attention to the “smooth step” potential problem.

## REFERENCES

- [1] A.O.Barut, A.Inomata and G.Junker: Path Integral Treatment of the Hydrogen Atom in a Space of Constant Curvature; *J.Phys. A* **20** (1987), 6271.
- [2] M.Böhm and G.Junker: The SU(1,1) Propagator as a Path Integral Over Compact Groups; *Phys.Lett. A* **117** (1986), 375.
- [3] M.Böhm and G.Junker: Path Integration Over Compact and Non-Compact Rotation Groups; *J.Math. Phys. A* **28** (1987), 1978.
- [4] F.Constantinescu and E.Magyari: Problems in Quantum Mechanics; (*Pergamon Press, Oxford, 1971*).
- [5] D.M.Dennison: The Infrared Spectra of Polyatomic Molecules; *Rev.Mod.Phys.* **3** (1931), 280.
- [6] I.H.Duru and H.Kleinert: Solution of the Path Integral for the H-Atom; *Phys.Lett. B* **84** (1979), 185; Quantum Mechanics of H-Atoms From Path Integrals; *Portugr.Phys.* **30** (1982), 401.
- [7] I.H.Duru: Path Integral Over the  $SU(2)$  Manifold and Related Potentials; *Phys.Rev. D* **30** (1984), 2121.
- [8] C.Eckart: The Penetration of a Potential Barrier by Electrons; *Phys.Rev.* **35** (1930), 1303.
- [9] A.Frank and K.B.Wolf: Lie Algebras for Systems With Mixed Spectra. I. The Scattering Pöschl-Teller Potential; *J.Math.Phys.* **25** (1985), 573.
- [10] C.Grosche: The Product Form for Path Integrals on Curved Manifolds; *Phys.Lett. A* **128** (1988), 113.
- [11] C.Grosche and F.Steiner: Path Integrals on Curved Manifolds; *Z.Phys. C* **38** (1987), 699.
- [12] C.Grosche and F.Steiner: The Path Integral on the Pseudosphere; *Ann.Phys.(N.Y.)* **182** (1988), 120.
- [13] C.Grosche and F.Steiner: Exactly Solvable Path Integrals; *DESY preprint*, in preparation.

- [16] A.Inomata: Remarks on the Time Transformation Technique for Path Integration; *Bielefeld Encounters in Physics and Mathematics VII; Path Integrals from meV to MeV, 1985*, pp.435; eds. M.C.Gutzwiller et al., World Scientific 1986.
- [14] L.Infeld and T.E.Hull: The Factorization Method; *Rev.Mod.Phys.* **23** (1951), 21.
- [15] L.Infeld and A.Schild: A Note on the Kepler Problem in a Space of Constant Negative Curvature; *Phys.Rev.* **67** (1945), 121.
- [17] A.Inomata and M.A.Kayed: Path Integral Quantization of the Symmetric Pöschl-Teller Potential; *Phys.Lett. A* **108** (1985), 9.
- [18] G.Junker and A.Inomata: Path Integrals on  $S^3$  and its Applications to the Rosen-Morse Oscillator; *Bielefeld Encounters in Physics and Mathematics VII*, pp.315.
- [19] H.Kleinert: How to do the Time-Sliced Path Integral of the H-Atom; *Phys.Lett. A* **120** (1987), 361.
- [20] L.Landau and E.M.Lifschitz: Lehrbuch der theoretischen Physik, Vol.I: Mechanik; (*Akademie Verlag, Berlin, 1979*).
- [21] M.F.Manning and N.Rosen: A Potential Function for the Vibrations of Diatomic Molecules; *Phys.Rev.* **44** (1933), 953.
- [22] N.M.Nieto: Exact Wave-Function Normalization for the  $B_0 \tanh z - U_0 \cosh^2 z$  and Pöschl-Teller Potentials; *Phys.Rev. A* **17** (1978), 1273.
- [23] K.Pak and I.Sökmén: Exact Path Integral Solutions of a Class of Potentials Related to the Rigid Rotator; *Phys.Lett. A* **103** (1984), 298.
- [24] K.Pak and I.Sökmén: General New Time Formalism in the Path Integral; *Phys.Rev. A* **30** (1984), 1629.
- [25] D.Peek and A.Inomata: Summation Over Feynman Histories in Polar Coordinates; *J.Math.Phys.* **10** (1969), 1422.
- [26] G.Pöschl and E.Teller: Bemerkungen zur Quantenmechanik des anharmonischen Oszillators; *Z.Phys.* **83** (1933), 143.
- [27] N.Rosen and P.M.Morse: On the Vibrations of Polyatomic Molecules; *Phys.Rev.* **42** (1932), 210.
- [28] F.Steiner: Space-Time Transformations in Radial Path Integrals; *Phys.Lett. A* **108** (1984), 356.
- [29] F.Steiner: Path Integrals in Polar Coordinates from eV to GeV; *Bielefeld Encounters in Physics and Mathematics VII*, pp.335.
- [30] Ig.Tamm: Die verallgemeinerte Kugelfunktionen und die Wellenfunktionen eines Elektrons im Felde eines Magnetpols; *Z.Phys.* **71** (1931), 141.