


DEUTSCHES ELEKTRONEN-SYNCHROTRON



DESY 95-099
May 1995



Conditionally Solvable Path Integral Problems

C. Grosche

II. Institut für Theoretische Physik, Universität Hamburg

ISSN 0418-9833

NOTKESTRASSE 85 - 22607 HAMBURG

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX,
send them to (if possible by air mail):

DESY
Bibliothek
Notkestraße 85
22607 Hamburg
Germany

DESY-IfH
Bibliothek
Platanenallee 6
15738 Zeuthen
Germany

CONDITIONALLY SOLVABLE PATH INTEGRAL PROBLEMS

Christian Grosche

*II. Institut für Theoretische Physik
Universität Hamburg, Luruper Chaussee 149
22761 Hamburg, Germany*

1 Introduction.

In recent years there has been enormous success in solving path integrals exactly. Milestones in the development have been the path integral solutions of Feynman of the harmonic oscillator [1, 2], the path integral solution of the radial harmonic oscillator [3]–[5], and the path integral solution of the Pöschl-Teller and modified Pöschl-Teller potential [6, 7], respectively. All these kind of problems have in common that they correspond to either a Gaussian-, Besselian- or Legendrian path integration. The couplings and parameters of the potentials are always assumed to take on arbitrary real values. An extensive list of all potential problems along with other path integral solutions, say in homogeneous spaces [8], will appear soon in a table of Feynman path integrals [9]; a classification scheme has been already announced in [10, 11]. It is remarkable that almost all of these solutions can be understood in terms of a group path integration, may it be a group path integration on the entire group space, or where an extension with the introduction of additional dummy variables is necessary [12, 13].

However, there are a couple of problems which generalize the well-known problems which are not entirely soluble by their own accord in the sense that all possible parameters are free to choose. These considerations can be made, of course, within the Schrödinger equation approach or within the path integral formalism. In the following I will be only concerned with the path integral approach. One set of such problems is called “quasi-exactly solvable” [14]. This means that a certain constraint on the parameters must be imposed, and then only some few low lying energy-levels together with the wavefunctions can be stated. Another set of problems is called “conditionally exactly solvable” [15]–[18]. They modify the usual potentials in quantum mechanics in a specific way such that they are quantum mechanically exactly solvable, however, the parameters and the coupling of the potentials are not completely free to choose.

In this article I discuss four different potentials of the later kind. The first potential generalizes and modifies the Coulomb potential. A $1/\sqrt{r}$ -term is incorporated and adds a significant long-range behaviour to the usual Coulomb interaction. As it turns out a specific form of the centrifugal barrier must be chosen (actually attractive) in order that the corresponding path integral can be solved. As has been pointed out by Stillinger [19] this particular set-up can produce a wide barrier around an attractive origin. By manipulations of the parameters the ground state energy can be moved up and down, and can also be moved to zero-energy, hence a potential with a bound-state character and a resonance character can be studied at the same time. Artificially as this looks alike, there exists an electrostatic charge density which actually produces such a potential field. A point charge located at the origin is accompanied by a strong $1/r^4$ and $1/r^{5/2}$ behaviour [19].

The second potential is a confinement potential with a dominant $r^{2/3}$ -behaviour for $r \rightarrow \infty$. Again a specific form of the radial potential (attractive) is required in order that the path integral can be solved. The potential is therefore an anharmonic radial oscillator. It has played a rôle in the modeling of quark-antiquark forces for mesons in nuclear physics [20].

As it turns out, both potentials are rather complicated concerning the proper formulation of the quantization condition. In each case it is required to solve a transcendental equation involving a parabolic cylinder function. This very point has been ignored in [15, 18], where a naive solution was claimed. The authors did not take into account

Some specific conditionally exactly solvable potentials are discussed within the path integral formalism. They generalize the usually known potentials by the incorporation of a fractional power behaviour and strongly anharmonic terms. We find four different kinds of such potentials, the first is related to the Coulomb potential, the second is an anharmonic confinement potential, and the third and the fourth are related to the Manning-Rosen potential.

ABSTRACT

that the radial problem remains a radial problem even after the transformations, and the deceiving regularity of the transformed problem (a shifted harmonic oscillator) does not allow a coordinate continuation to the entire \mathbb{R} (I sketch the naïve solution, though). As we will see, it is not possible to state the propagator exactly. However, the corresponding energy dependent Green function can be stated in closed form. The proper quantization conditions follows from the poles of the Green functions.

The third and the fourth potential are modifications of a Eckart potential [21] or a Rosen-Morse oscillator [22] and model potential troughs. These kinds of potentials play a rôle in the theory of molecules [21, 22], solitons and reflectionless potentials [23]. In the first of the two potentials the proper transformed potential is of the Manning-Rosen type [24], which was also used as a screened Coulomb potential [25, 26] with exponential decay (*s*-wave Yukawa potential). The second leads to hyperbolic Scarf-like potential [27, 28] with an even stronger screening of the potential energy. The connection to supersymmetric quantum mechanics with all these potentials has been pointed out by Dutt et al. [17], Nag et al. [18], and Papp [29].

Although exactly solvable, these potentials are complicated enough to be of serious consideration in modeling of actual physical forces. By choosing a path integral approach we succeed in gaining comprehensive information of the bound-state solutions of these potentials (if they exist), and what is often more important, in the scattering states which eventually allow for the calculation of cross-sections and phase-shifts which will be taken into consideration elsewhere.

This article is organized as follows. In the following part I sketch some necessary information concerning transformation techniques in the path integral. I the third section I present the four "conditionally exactly solvable" potentials labeled V_1 , V_2 , V_3 and V_4 . The well-established space-time transformation technique reduces each path integral problem to an already known one. The final result in each case includes the statement of the corresponding Green function. For the first two potentials, where the bound-state solutions are only implicitly known through a transcendental equation, this is sufficient. In the other two the bound-state wave-functions, the energy-spectrum, and the continuous states are displayed explicitly. The last section contains a summary and a short discussion.

2 Space-Time Transformation Technique.

In order to make the article self-contained, let us cite shortly the space-time transformation technique, e.g., Refs. [9]–[11, 30]–[39], and references therein. We consider a path integral

$$K(x'', x'; T) = \int_{x(t')=x'}^{x(t'')=x''} \mathcal{D}x(t) \exp \left\{ \frac{i}{\hbar} \int_{t'}^{t''} \left[\frac{m}{2} \dot{x}^2 - V(x) \right] dt \right\}, \quad (1)$$

where it is assumed that the potential $V(x)$ is so complicated that a direct path integral evaluation is not possible. Now a transformation $x = F(q, t)$ and the so-called "time transformation" is implemented by introducing a new "pseudo-time" s'' . In order to do this, one first makes use of the operator identity

$$\frac{1}{H-E} = f_r(x, t) \frac{1}{f_l(x, t)(H-E)} f_l(x, t), \quad (2)$$

where H is the Hamiltonian corresponding to the path integral $K(T)$, and $f_r(x, t)$ are functions in x and t , multiplying from the left or from the right, respectively, onto the operator $(H-E)$. Secondly, the introduced pseudo-time s'' is assumed to obey the constraint

$$\int_0^{s''} ds f_l(F(q(s), s)) f_r(F(q(s), s)) = T = t'' - t', \quad (3)$$

and has for all admissible paths a unique solution $s'' > 0$ given by

$$s'' = \int_{t'}^{t''} \frac{dt}{f_l(x, t) f_r(x, t)} = \int_{t'}^{t''} \frac{ds}{F'^2(q(s), s)}. \quad (4)$$

Here one has made the choice $f_l(F(q(s), s)) = f_r(F(q(s), s)) = F'(q(s), s)$ in order that in the final result the metric coefficient in the kinetic energy term is equal to one. A convenient way to derive the corresponding transformation formulæ uses the energy dependent Green function $G(E)$ of the kernel $K(T)$ defined by $(F(q, t) \equiv F(q)$ time-independent)

$$G(q'', q'; E) = \left\langle q'' \left| \frac{1}{H-E-ic} \right| q' \right\rangle = \frac{i}{\hbar} \int_0^{\infty} dT e^{i(E+ic)T/\hbar} K(q'', q'; T), \quad (5)$$

where a small positive imaginary part ($\epsilon > 0$) has been added to the energy E . (Usually we not explicitly write the ic , but will tacitly assume that the various expressions are regularized according to this rule). For the path integral $K(T)$ one obtains the following transformation formulæ

$$K(x'', x'; T) = \int_{-\infty}^{\infty} \frac{dE}{2\pi i} e^{-iET/\hbar} G(q'', q'; E), \quad (6)$$

$$G(q'', q'; E) = \frac{i}{\hbar} \left[F''(q'') F'(q') \right]^{1/2} \int_0^{\infty} ds'' \hat{K}(q'', q'; s''), \quad (7)$$

with the transformed path integral \hat{K} given by

$$\hat{K}(q'', q'; s'') = \int_{q(0)=q'}^{q(s'')=q''} \mathcal{D}q(s) \exp \left\{ \frac{i}{\hbar} \int_0^{s''} \left[\frac{m}{2} \dot{q}^2 - F'^2(q) (V(F(q)) - E) - \Delta V(q) \right] ds \right\} \quad (8)$$

and the quantum potential ΔV given by

$$\Delta V(q) = 8m \left(\frac{F''^2}{F'^2} - 2 \frac{F'''}{F'} \right). \quad (9)$$

These formulæ are sufficient for our purposes.

3 The Potentials.

3.1 The modified Coulomb potential.

The first potential we are going to study has the following form ($r > 0$)

$$V_1(r) = \frac{\hbar^2}{2m} \frac{\gamma}{r^2} - \frac{Zq}{r} + \frac{b}{\sqrt{r}}. \quad (10)$$

Zq^2 is a Coulomb-coupling, $b \in \mathbb{R}$, and γ is a constant which will be determined. Obviously, this potential is a generalization of a pure Coulomb-potential. The centrifugal term usually makes no difficulty, but the $r^{-1/2}$ -long-range term significantly alters the behaviour of the potential for $r \rightarrow \infty$. I proceed in the canonical manner and perform a coordinate transformation $r = u^2$, $t = 2uu'$ together with a time transformation $\Delta t_j = 4u_j u_{j-1} \Delta s_j$ in each short-time interval Δt_j to the new time $s(t) = \int_t^t d\sigma u^{-2}(\sigma)$. I obtain

$$\begin{aligned} K^{(V_1)}(r'', r'; T) &= \int_{r'(t')=r'}^{r(t'')=r''} \mathcal{D}r(t) \exp \left\{ \frac{i}{\hbar} \int_{t'}^{t''} \left[\frac{m}{2} \dot{r}^2 - \left(\frac{\hbar^2}{2m} \frac{\gamma}{r^2} - \frac{Zq^2}{r} + \frac{b}{\sqrt{r}} \right) \right] dt \right\} \\ &= 2(r' r'')^{1/4} \int_{\mathbb{R}} \frac{dE}{2\pi \hbar} e^{-iET/\hbar} \int_0^\infty ds s'' e^{iZq^2 s''/\hbar} \\ &\quad \times \int_{u(0)=u'}^{u(s'')=u''} \mathcal{D}u(s) \exp \left\{ \frac{i}{\hbar} \int_0^{s''} \left[\frac{m}{2} \dot{u}^2 - \left(\frac{\hbar^2}{2m} \frac{16\gamma + 3}{4u^2} - 4Eu^2 + 4bu \right) \right] ds \right\}. \end{aligned} \quad (11)$$

This path integral is at it stands not solvable. However, if we set $\gamma = -3/16$, the centrifugal term vanishes due to $\Delta V = 3\hbar^2/8mu^2$, and we seem to have a shifted harmonic oscillator path integral. Let us for the moment continue with the naive analysis according to [15, 18], where $u \in \mathbb{R}$ is assumed. Performing the additional variable shift $v = u - b/2E$ we get

$$\begin{aligned} K^{(V_1)}(r'', r'; T) &= 2(r' r'')^{1/4} \int_{\mathbb{R}} \frac{dE}{2\pi \hbar} e^{-iET/\hbar} \int_0^\infty ds s'' e^{i s'' (4Zq^2 - b^2/E)/\hbar} \\ &\quad \times \int_{v(0)=v'}^{v(s'')=v''} \mathcal{D}v(s) \exp \left[\frac{im}{2\hbar} \int_0^{s''} (v^2 - \omega^2 v^2) ds \right], \end{aligned} \quad (12)$$

where $\omega^2 = -8E/m$. Provided $v \in \mathbb{R}$, the last expression can now be analysed in various ways: In order to obtain the discrete spectrum one can insert the explicit form of the propagator of the harmonic oscillator and can expand it by means of the Mehler formula to obtain the wave-functions and the energy-spectrum of the bound state of the path integral (12); secondly, one can insert the explicit form of the propagator and use a dispersion relation to obtain the wave-functions of the continuous spectrum. The third possibility is to insert the Green function of the harmonic oscillator: By analyzing its analytic pole structure it follows that the Green function of the harmonic oscillator must be evaluated at the energy $E \rightarrow 4Zq^2 - b^2/E$, i.e., we obtain by performing the s'' -integration the Green function of the path integral (12)

$$G(v'', v'; E) = \sqrt{\frac{m}{\pi \hbar^3 \omega}} \Gamma(-\nu) D_\nu \left(\sqrt{\frac{2m\omega}{\hbar}} v_> \right) D_\nu \left(-\sqrt{\frac{2m\omega}{\hbar}} v_< \right). \quad (13)$$

Here $\nu = -\frac{1}{2} + (4Zq^2 - b^2/E)/\hbar\omega$, and $D_\nu(z)$ is a parabolic cylinder function. The poles of $G(E)$ yield the bound-state energy-spectrum and wave-functions and the cut the continuous spectrum. The poles are determined by the poles of the Γ -function and we obtain the quantization condition

$$4Zq^2 - b^2/E_n = 2\hbar(n + \frac{1}{2})\sqrt{-2E_n}/m, \quad n \in \mathbb{N}_0. \quad (14)$$

With some algebraic manipulations this can be cast into a cubic equation in terms of E_n yielding

$$(n + \frac{1}{2})^2 \hbar^2 E_n^3 + 2mZ^2 q^4 E_n^2 - Zq^2 b^2 m E_n + mb^4/8 = 0. \quad (15)$$

From the generally three solutions of this equation, the physically relevant one is selected by requiring that for $b = 0$ a Coulombic spectrum should emerge. Hence we get

$$E_n = \sqrt[3]{\sqrt{D} - \frac{Q}{2}} - \sqrt[3]{\sqrt{D} + \frac{Q}{2}} - \frac{R}{3} \quad (16)$$

$$D = \left(\frac{P}{3} \right)^3 + \left(\frac{Q}{2} \right)^2, \quad P = \frac{3S - R^2}{3}, \quad Q = \frac{2R^3}{27} - \frac{RS}{3} + T, \quad (17)$$

$$R = \frac{2mZ^2 q^4}{\hbar^2 (n + \frac{1}{2})^2}, \quad S = -\frac{mZq^2 b^2}{\hbar^2 (n + \frac{1}{2})^2}, \quad T = \frac{mb^4}{8\hbar^2 (n + \frac{1}{2})^2}. \quad (18)$$

The bound state wave-functions then have the form

$$\begin{aligned} \Psi_n^{(V_1)}(r) &= N_{E_n} \left(\sqrt{\frac{m\omega_n}{\pi \hbar}} r \frac{1}{2^n n!} \right)^{1/2} \\ &\quad \times H_n \left[\sqrt{\frac{m\omega_n}{\hbar}} \left(\sqrt{r} - \frac{b}{2E_n} \right) \right] \exp \left[-\frac{m\omega_n}{2\hbar} \left(\sqrt{r} - \frac{b}{2E_n} \right)^2 \right], \end{aligned} \quad (19)$$

$$N_{E_n} = \left(\frac{2|E_n|^{3/2}}{(n + \frac{1}{2}) [3(E_n + R/3)^2 - R^2/3 + S]} \right)^{1/2}. \quad (20)$$

Up to the additional proper normalization factor N_{E_n} this is the result of [15, 18]. Let us note that for the case $Z = 0$ we can repeat the analysis and obtain a discrete energy-spectrum according to

$$E_n = -\sqrt[3]{\frac{mb^4}{8\hbar^2 (n + \frac{1}{2})^2}}. \quad (21)$$

This energy-spectrum is of genuine non-Coulombic behaviour.

However, this easy-to-obtain solution can not be considered as correct! As pointed out in [19] a wave-function with a $r^{1/4}$ -behaviour at the origin of a singular potential is physically unacceptable [40]. Therefore we must discard the above solution of [15, 18] entirely. In [11] I have made the implicit assumption that it is possible to extend the variable u to the entire \mathbb{R} . This is in contrast to the one-dimensional Kustaaheimo-Stiefel transformation which maps $\mathbb{R}^+ \rightarrow \mathbb{R}^+$ [41, 42]. In the usual radial Coulomb problem this is an obvious mapping because the radial Coulomb problem in the variable r is mapped onto a radial harmonic oscillator in the variable u . Therefore the path integral (11) is a radial path integral with $u > 0$, and the path integral (12) is a radial path integral with $v > -b/2E$ for fixed energy E . The additional linear term spoils the symmetry with respect to reflections in the variable u . In [43, 44] I have developed a procedure how to deal with such problems within the path integral. We assume that we have evaluated a path integral problem with a potential $V(x)$ in, say, the entire \mathbb{R} . This path integral is called $K^{(V)}(T)$. The corresponding Green function is denoted by $G^{(V)}(E)$. Now we consider the path integral problem with the same potential V , but with Dirichlet boundary conditions at the location $x = a$ and we consider the half-space $x > a$. Then the Green function in

the half-space $x > a$ is given by [43, 44]

$$\begin{aligned} & \frac{1}{\hbar} \int_0^\infty dT e^{iET/\hbar} \int_{x^{(v')}=x'}^{x^{(v'')}=x''} \mathcal{D}_{(x>a)}^{(D)} x(t) \exp \left\{ \frac{i}{\hbar} \int_r^{t''} \left[\frac{m}{2} \dot{x}^2 - V(x) \right] dt \right\} \\ & = G^{(V)}(x'', x'; E) - \frac{G^{(V)}(x'', a; E) G^{(V)}(a, x'; E)}{G^{(V)}(a, a; E)}. \end{aligned} \quad (22)$$

We identify the Green function $G^{(V)}(E)$ with (13) and obtain consequently

$$\begin{aligned} G^{(V_1)}(r'', r'; E) & = 2(r'r'')^{1/4} \sqrt{\frac{m}{\pi \hbar^3 \omega}} \Gamma(-\nu) \\ & \times \left\{ D_\nu \left[\sqrt{\frac{2m\omega}{\hbar}} \left(\sqrt{r>} - \frac{b}{2E} \right) \right] D_\nu \left[-\sqrt{\frac{2m\omega}{\hbar}} \left(\sqrt{r<} - \frac{b}{2E} \right) \right] \right. \\ & - D_\nu \left[\sqrt{\frac{2m\omega}{\hbar}} \left(\sqrt{r'} - \frac{b}{2E} \right) \right] D_\nu \left[-\sqrt{\frac{2m\omega}{\hbar}} \left(\sqrt{r'} - \frac{b}{2E} \right) \right] \\ & \left. \times D_\nu \left(\sqrt{\frac{2m\omega}{\hbar}} \frac{b}{2E} \right) / D_\nu \left(-\sqrt{\frac{2m\omega}{\hbar}} \frac{b}{2E} \right) \right\}. \end{aligned} \quad (23)$$

This determines the energy-spectrum by the zeros of the parabolic cylinder function, i.e.,

$$D_{\nu_n} \left(-\sqrt{\frac{2m\omega_n}{\hbar}} \frac{b}{2E_n} \right) = 0. \quad (24)$$

This result is in accordance with [19]. I have indicated by $\nu_n = \nu(E_n)$ and $\omega_n = \omega(E_n)$ the explicit dependence on E_n . The analysis in [44] showed that the poles coming from the prefactor in (23) play no rôle in the corresponding boundary condition problem. The case $b = 0$ is contained in (23) by noting that for $b = 0$ the path integral (11) is a radial path integral in u with angular momentum $l = 0$ [32].

3.2 A radial confinement potential.

The second potential we want to consider has the form ($r > 0$)

$$V_2(r) = \frac{m\omega^2 r^{2/3}}{2} + \frac{B}{r^{2/3}} + \frac{\hbar^2 \gamma}{2m r^2}. \quad (25)$$

This potentials models a confinement potential for quark-antiquark interactions [20], however not quite oscillator-like, and has a singularity at the origin. We perform a space-time transformation as before with $r = \tilde{x}^{3/2}$. In doing that we set $\gamma = -5/36$ in order to cancel the corresponding term in the emerging quantum potential $\Delta V = 5\hbar^2/32m\tilde{x}^2$ due to the transformation. Hence we obtain

$$\begin{aligned} K^{(V_2)}(r'', r'; E) & = \int_{r^{(v')}=r'}^{r^{(v'')}=r''} D\tilde{x}(t) \\ & \times \exp \left\{ \frac{i}{\hbar} \int_r^{t''} \left[\frac{m}{2} \dot{\tilde{x}}^2 - \left(\frac{m}{2} \omega^2 r^{2/3} + \frac{B}{r^{2/3}} - \frac{\hbar^2 \gamma}{2m} \frac{5}{36r^2} \right) \right] dt \right\} \end{aligned} \quad (26)$$

$$\begin{aligned} & = \frac{3}{2} (r'r'')^{1/6} \int_{\mathbb{R}} \frac{dE}{2\pi\hbar} e^{-iET/\hbar} \int_0^\infty ds'' e^{-9Bs''/4\hbar} \\ & \times \int_{u^{(v')}=u'}^{u^{(v'')}=u''} D u(s) \exp \left[\frac{i}{\hbar} \int_0^{s''} \left(\frac{m}{2} \dot{u}^2 + \frac{9}{4} Eu - \frac{9m}{4} \omega^2 u^2 \right) ds \right] \\ & = \frac{3}{2} (r'r'')^{1/6} \int_{\mathbb{R}} \frac{dE}{2\pi\hbar} e^{-iET/\hbar} \int_0^\infty ds'' \exp \left[\frac{i}{\hbar} s''^9 \left(\frac{E^2}{4} (2m\omega^2 - B) \right) \right] \\ & \times \int_{v^{(v')}=v'}^{v^{(v'')}=v''} D v(s) \exp \left[\frac{im}{2\hbar} \int_0^{s''} (v^2 - \tilde{\omega}^2 v^2) ds \right]. \end{aligned} \quad (27)$$

In the last step we have set $\tilde{\omega} = 3\omega/2$ and $v = u - E/m\omega^2$. Inserting the Green function for the harmonic oscillator gives the Green function for path integral in the variable v ($v \in \mathbb{R}$):

$$G(v'', v'; E) = \sqrt{\frac{3m}{2\pi\hbar^3 \omega}} \Gamma(-\nu) D_\nu \left(\sqrt{\frac{3m\omega}{\hbar}} v_{>} \right) D_\nu \left(-\sqrt{\frac{3m\omega}{\hbar}} v_{<} \right), \quad (29)$$

with $\nu = -\frac{1}{2} - \frac{3}{2}(E^2/2m\omega^2 - B)/\hbar\omega$. Provided $v \in \mathbb{R}$, the poles of the Green function would determine the energy-spectrum given by $E_n = \sqrt{2m\omega^2[2(n + \frac{1}{2})\hbar\omega/3 + B]}$ for $n \in \mathbb{N}_0$ ($\omega^2 > 0, B > -\hbar\omega/3$) [18]. But for the same reasons as in sect.3.1 this "solution" must be discarded.

In order to obtain the proper solution in the half-space $v > -E/m\omega^2$ for fixed energy E , i.e., for the potential V_2 we proceed similarly as for the previous case and obtain

$$\begin{aligned} G^{(V_2)}(r'', r'; E) & = \frac{3}{2} (r'r'')^{1/6} \sqrt{\frac{3m}{2\pi\hbar^3 \omega}} \Gamma(-\nu) \\ & \times \left\{ D_\nu \left[\sqrt{\frac{3m\omega}{\hbar}} \left(r_{>}^{2/3} - \frac{E}{m\omega^2} \right) \right] D_\nu \left[-\sqrt{\frac{3m\omega}{\hbar}} \left(r_{<}^{2/3} - \frac{E}{m\omega^2} \right) \right] \right. \\ & + D_\nu \left[\sqrt{\frac{3m\omega}{\hbar}} \left(r^{2/3} - \frac{E}{m\omega^2} \right) \right] D_\nu \left[\sqrt{\frac{3m\omega}{\hbar}} \left(r^{\mu 2/3} - \frac{E}{m\omega^2} \right) \right] \\ & \left. \times D_\nu \left(\sqrt{\frac{3m\omega}{\hbar}} \frac{E}{m\omega^2} \right) / D_\nu \left(-\sqrt{\frac{3m\omega}{\hbar}} \frac{E}{m\omega^2} \right) \right\}. \end{aligned} \quad (30)$$

This determines the energy-spectrum by the zeros of the parabolic cylinder function, i.e.

$$D_{\nu_n} \left(-\sqrt{\frac{3m\omega}{\hbar}} \frac{E_n}{m\omega^2} \right) = 0. \quad (31)$$

In the case that $\omega^2 < 0$ we have to replace $\omega \rightarrow i\omega$ in (30). In the special case $\omega = 0$ the corresponding Green function can be obtained by using the Green function for the linear potential [44] and we get

$$\begin{aligned} G_{\omega=0}^{(V_2)}(r'', r'; E) & = 2(r'r'')^{1/6} \frac{m}{\hbar^2} \left[\left(r^{2/3} - \frac{B}{E} \right) \left(r^{\mu 2/3} - \frac{B}{E} \right) \right]^{1/2} \\ & \times \left\{ K_{1/3} \left[\frac{\sqrt{-2mE}}{\hbar} \left(r_{>}^{2/3} - \frac{B}{E} \right)^{3/2} \right] I_{1/3} \left[\frac{\sqrt{-2mE}}{\hbar} \left(r_{<}^{2/3} - \frac{B}{E} \right)^{3/2} \right] \right\} \end{aligned}$$

$$-K_{1/3} \left[\frac{\sqrt{-2mE}}{\hbar} \left(r^{2/3} - \frac{B}{E} \right)^{3/2} \right] K_{1/3} \left[\frac{\sqrt{-2mE}}{\hbar} \left(r^{2/3} - \frac{B}{E} \right)^{3/2} \right] \\ \times I_{1/3} \left(-\frac{B}{\hbar E} \sqrt{2mB} \right) / K_{1/3} \left(-\frac{B}{\hbar E} \sqrt{2mB} \right) \} \quad (32)$$

Possible bound states are determined by

$$K_{1/3} \left(-\frac{B}{\hbar E_n} \sqrt{2mB} \right) = 0. \quad (33)$$

Due to the property

$$\text{Ai}(z) = \frac{1}{\pi} \left(\frac{z}{3} \right)^{1/2} K_{1/3} \left(\frac{2}{3} z^{3/2} \right) \quad (34)$$

the zeros are determined by the zeros of the Airy function $\text{Ai}(z)$, $z = -\alpha_n$ ($\alpha_n > 0$, $n \in \mathbb{N}_0$). From the relation

$$\text{Ai} \left(-\frac{B}{E} \sqrt{\frac{-9mE}{2\hbar^2}} \right) = 0, \quad (35)$$

we see that negative bound states are allowed only if $B < 0$. Hence it follows

$$E_n = -\sqrt{\frac{9m}{2\hbar^2}} \left(\frac{|B|}{\alpha_n} \right)^{3/2}, \quad n \in \mathbb{N}_0. \quad (36)$$

Therefore the singular term $B/r^{2/3}$ must be attractive in order that bound states can exist. Note that no resonance states exist because all zeros of $\text{Ai}(z)$ are located on the negative real axis [45, p.166]. The ground state energy is $E_0 = -\sqrt{9m/2\hbar^2} (|B|/2.341\dots)^{3/2}$, and the accumulation point is $E_\infty = 0$.

3.3 A modified Rosen-Morse potential I.

The third potential I want to consider has the form ($x \in \mathbb{R}$)

$$V_3(x) = -\frac{A}{\sqrt{1+e^{-2x}}} + \frac{B}{1+e^{-2x}} + \frac{C}{(1+e^{-2x})^2}. \quad (37)$$

I perform the transformation $x = \ln(\sinh u)$, $u > 0$, together with the appropriate time transformation. The emerging quantum potential is

$$\Delta V(u) = -\frac{\hbar^2}{8m} \left(\frac{3}{\cosh^2 u} + \frac{1}{\sinh^2 u} \right). \quad (38)$$

In order that the terms $\alpha^{-1}/\cosh^2 u$ cancel we must set $C = -3\hbar^2/8m$. This gives for the path integral

$$K^{(V_3)}(r'', r'; T) \\ = \int_{x(t')=x'}^{x(t'')=x''} \mathcal{D}x(t) \exp \left\{ \frac{i}{\hbar} \int_{t'}^{t''} \left[\frac{m}{2} \dot{x}^2 - \left(\frac{B}{1+e^{-2x}} - \frac{A}{\sqrt{1+e^{-2x}}} - \frac{3\hbar^2}{8m(1+e^{-2x})^2} \right) \right] dt \right\} \\ = \left(\coth u' \coth u'' \right)^{1/2} \int_{\mathbb{R}} \frac{dE}{2\pi\hbar} e^{-iET/\hbar} \int_0^\infty ds \int_0^\infty ds' \exp \left[\frac{i}{\hbar} \left(\frac{3\hbar^2}{8m} - B + E \right) s'' \right] \\ \times \int_{u(0)=u'}^{u(s'')=u''} \mathcal{D}u(s) \exp \left[\frac{i}{\hbar} \int_0^{s''} \left(\frac{m}{2} \dot{u}^2 + A \coth u + \hbar^2 \frac{2mE/\hbar^2 + 1/4}{2m \sinh^2 u} \right) ds \right]. \quad (39)$$

The path integral in the variable u is the path integral for the Manning-Rosen potential $V_{MR}(u) = -A \coth u + B'/\sinh^2 u$ [46]. From the spectral representation of the Manning-Rosen potential we therefore derive the following quantization condition for the bound states ($n = 0, 1, \dots, N_{Max}$)

$$B - \frac{3\hbar^2}{8m} - E_n = \frac{\hbar^2}{2m} \left(n + \frac{1}{2} + \sqrt{-2mE_n/\hbar} \right)^2 + \frac{mA^2/2}{\left(n + \frac{1}{2} + \sqrt{-2mE_n/\hbar} \right)^2}. \quad (40)$$

This can be rewritten into a cubic equation for $\sqrt{-E_n}$

$$\frac{2(n + \frac{1}{2})}{\hbar} \sqrt{2m(-E_n)^{3/2}} + \left[5(n + \frac{1}{2})^2 - \frac{2m}{\hbar^2} \left(B - \frac{3\hbar^2}{8m} \right) \right] (-E_n) \\ + 2 \left[\frac{\hbar^2(n + \frac{1}{2})^2}{m} - \left(B - \frac{3\hbar^2}{8m} \right) \right] \left(n + \frac{1}{2} \right) \frac{\sqrt{2m(-E_n)^{1/2}}}{\hbar} \\ + \left[\frac{\hbar^2}{2m} \left(n + \frac{1}{2} \right)^4 + \frac{mA^2}{2} - \left(B - \frac{3\hbar^2}{8m} \right) \right] = 0. \quad (41)$$

On the other hand we obtain from the Green function representation of the Manning-Rosen potential [9, 27, 47]

$$G^{(V_3)}(x'', x'; E) = \left(\coth u' \coth u'' \right)^{1/2} \frac{m}{\hbar^2} \frac{\Gamma(m_1 - L_{E'}) \Gamma(L_{E'} + m_1 + 1)}{\Gamma(m_1 + m_2 + 1) \Gamma(m_1 - m_2 + 1)} \\ \times \left(\frac{1}{1+u'} \cdot \frac{1}{1+u''} \right)^{\frac{1}{2}(m_1+m_2+1)} \left(\frac{u'}{1+u'} \cdot \frac{u''}{1+u''} \right)^{\frac{1}{2}(m_1-m_2)} \\ \times {}_2F_1 \left(-L_{E'} + m_1, L_{E'} + m_1 + 1; m_1 - m_2 + 1; \frac{u_-}{1+u_-} \right) \\ \times {}_2F_1 \left(-L_{E'} + m_1, L_{E'} + m_1 + 1; m_1 + m_2 + 1; \frac{1}{1+u_+} \right). \quad (42)$$

Here are $\sinh u = e^x$, $L_{E'} = -\frac{1}{2} + \frac{1}{2} \sqrt{2m(A-E')/\hbar}$, $m_{1/2} = \frac{1}{2}(s \pm \sqrt{-2m(A+E')/\hbar})$, $s = 2\sqrt{-2mE}/\hbar$, and $E' = 3\hbar^2/8m - B + E$. The poles of the Green function determine the energy spectrum which coincides with the one determined in (40), and the corresponding residues give the wave-functions expansion ($s_n = s(E_n)$, $k_1 = \frac{1}{2}[1 + \frac{1}{2}(s_n + 2n + 1) + 2mA/\hbar^2(s_n + 2n + 1)]$)

$$\Psi_n^{(V_3)}(x) = N_{E_n}^{(V_3)} \left[\left(1 + \frac{4mA}{\hbar^2(s_n + 2n + 1)^2} \right)^{2k_1 - n - s_n - 2} m! \Gamma(2k_1 - n - 1)^{1/2} \right. \\ \left. \times \sqrt{\coth u} (1 - e^{-2u})^{s_n+1/2} e^{-2u(k_1 - s_n/2 - n - 1)} P_n^{(2k_1 - 2n - s_n - 2, s_n)}(1 - 2e^{-2u}) \right], \quad (43)$$

$$E_n = -\frac{\hbar^2}{2m} \left(\sqrt[3]{\sqrt{D} + \frac{Q}{2}} - \sqrt[3]{\sqrt{D} - \frac{Q}{2}} + \frac{R}{3} \right)^2, \quad (44)$$

$$R = \frac{5(n + \frac{1}{2})^2 - 2m(B - \frac{3\hbar^2}{8m})/\hbar^2}{2(n + \frac{1}{2})^4 + mA^2/2 - (B - \frac{3\hbar^2}{8m})}, \quad S = 2 \left[\left(n + \frac{1}{2} \right)^2 - \frac{m}{\hbar^2} \left(B - \frac{3\hbar^2}{8m} \right) \right], \quad (45)$$

$$T = m \frac{2(n + \frac{1}{2})\hbar^2}{(n + \frac{1}{2})\hbar^2}, \\ N_{E_n}^{(V_3)} = \left(\frac{\sqrt{-2mE_n} (n + \frac{1}{2} + \sqrt{-2mE_n/\hbar})^2}{\hbar(n + \frac{1}{2}) [3(\sqrt{-2mE_n/\hbar} + R/3)^2 - E^2/3 + S]} \right)^{1/2}. \quad (46)$$

Bound states exist if $A > 0$ and the maximal number N_{Max} of bound states is found by requiring $E_n < 0$. The cut of the Green function determine the continuous spectrum, and the corresponding wave-functions are determined by the method of [47]. Thus the wave functions and the energy-spectrum of the continuous states are given by

$$\Psi_p^{(V_3)}(x) = N_p^{(V_3)} \sqrt{\coth u} u^{-\nu/2} (u-1)^{[\nu-(1+s_p)]/2} \times {}_2F_1\left(\frac{1+s_p+i(\tilde{p}-p)}{2}, \frac{1+s_p-i(\tilde{p}+p)}{2}; s_p+1; \frac{1}{1-u}\right) \quad (47)$$

$$N_p^{(V_3)} = \frac{1}{\Gamma(2k_2)} \sqrt{\frac{p \sinh \pi p}{2\pi^2}} \left[\Gamma(k_1+k_2-\kappa) \Gamma(-k_1+k_2+\kappa) \right]^{1/2} \times \Gamma(k_1+k_2+\kappa-1) \Gamma(-k_1+k_2-\kappa+1) \quad (48)$$

$[\kappa = \frac{1}{2}(1+ip), k_2 = \frac{1}{2}(1+s_p), s_p = s(E_p), \tilde{p} = \sqrt{2m(E_p-A)/\hbar}, p \in \mathbb{R}]$ with energy spectrum $E_p = \hbar^2 p^2/2m - A + B - 3\hbar^2/8m$. The results concerning the bound states coincide with [17]. Note that the transformation $x = \frac{1}{2} \ln(-\cosh^2 v)$ leads to a Rosen-Morse potential. However, this transformation is not longer a real transformation which causes interpretation difficulties, and therefore not used.

3.4 A modified Rosen-Morse potential II.

The fourth potential I treat has the form ($x \in \mathbb{R}$)

$$V_4(x) = \frac{A}{1+e^{-2x}} - \frac{Be^{-x}}{\sqrt{1+e^{-2x}}} - \frac{3\hbar^2}{8m(1+e^{-2x})^2}. \quad (49)$$

I perform the same transformation $x = \ln(\sinh u)$ together with the appropriate time transformation, and I have set the coupling in the third term equal to $C = -3\hbar^2/8m$ in order that it cancels with the corresponding term in the emerging quantum potential. This gives for the path integral

$$\begin{aligned} K^{(V_4)}(r'', r'; T) &= \int_{x(t')=x'}^{x(t'')=x''} \mathcal{D}x(t) \exp \left\{ \frac{i}{\hbar} \int_{t'}^{t''} \left[\frac{m}{2} \dot{x}^2 - \left(\frac{A}{1+e^{-2x}} - \frac{Be^{-x}}{\sqrt{1+e^{-2x}}} - \frac{3\hbar^2}{8m(1+e^{-2x})^2} \right) \right] dt \right\} \\ &= \left(\coth u' \coth u'' \right)^{1/2} \int_{\mathbb{R}} \frac{dE}{2\pi\hbar} e^{-iET/\hbar} \int_0^\infty ds'' \exp \left[\frac{i}{\hbar} \left(\frac{\hbar^2}{8m} - A \right) s'' \right] \\ &\quad \times \int_{u(0)=u'}^{u(s'')=u''} \mathcal{D}u(s) \exp \left[\frac{i}{\hbar} \int_0^{s''} \left(\frac{m}{2} \dot{u}^2 + B \frac{\coth u}{\sinh u} + \hbar^2 \frac{2mE/\hbar^2 + 1/4}{2m \tanh^2 u} \right) ds \right]. \end{aligned} \quad (50)$$

The path integral in the variable u is a path integral of a hyperbolic Scarf-like potential $V_{HS}(u) = \frac{\hbar^2}{2m}(V_0+V_1 \coth^2 u + V_2 \frac{\cosh u}{\sinh u})$ as discussed in [27]. We identify $V_0 = 2m\lambda/\hbar^2 - \frac{1}{2}$, $V_1 = -(2mE/\hbar^2 + \frac{1}{4})$, $V_2 = -2mB/\hbar^2$. It has energy eigenvalues $E_n = \frac{\hbar^2}{2m}(V_0 + V_1 - \frac{\hbar^2}{8m}[2(k_1 - k_2 - n) - 1])^2$. From its spectral expansion we derive the quantization condition for the potential V_4 ($n = 0, 1, \dots, N_{Max}$)

$$\sqrt{A - E_n - \frac{3\hbar^2}{8m}} = \frac{1}{2} \left(\sqrt{B - E_n} - \sqrt{-B - E_n} \right) - \frac{\hbar}{\sqrt{2m}} (n + \frac{1}{2}), \quad (51)$$

This give after some algebra a cubic equation in $(-E_n)$ ($\lambda = A+C+\tilde{n}^2$, $C = -3\hbar^2/8m$, $\tilde{n} = \hbar(n + \frac{1}{2})/\sqrt{2m}$)

$$\begin{aligned} &4\tilde{n}^2(-E_n)^3 + [12\tilde{n}^2(\tilde{n}^2 + \lambda) - \lambda^2](-E_n)^2 \\ &+ [16\tilde{n}^2\lambda(A+C+\lambda) - 2(\lambda + 4\tilde{n}^2)(\lambda^2 + \frac{B^2}{4} + 4\tilde{n}^2(A+C))](-E_n) \\ &+ [16\tilde{n}^2\lambda^2(A+C) - (\lambda^2 + \frac{B^2}{4} + 4\tilde{n}^2(A+C))^2] = 0. \end{aligned} \quad (52)$$

From the Green function of the hyperbolic Scarf-like potential we derive the Green function for the potential V_4

$$\begin{aligned} G^{(V_4)}(x'', x'; E) &= (\coth u' \coth u'')^{1/2} \frac{2m}{\hbar^2} \frac{\Gamma(m_1 - L_\nu) \Gamma(L_\nu + m_1 + 1)}{\Gamma(m_1 + m_2 + 1) \Gamma(m_1 - m_2 + 1)} \\ &\quad \times (\cosh u' \cosh u'')^{-(m_1 - m_2)} (\tanh u' \tanh u'')^{m_1 + m_2 + \frac{1}{2}} \\ &\quad \times {}_2F_1\left(-L_\nu + m_1, L_\nu + m_1 + 1; m_1 - m_2 + 1; \frac{1}{\cosh^2 u_C}\right) \\ &\quad \times {}_2F_1\left(-L_\nu + m_1, L_\nu + m_1 + 1; m_1 + m_2 + 1; \tanh^2 u_C\right) \end{aligned} \quad (53)$$

with $\sinh u = e^\pm$, $m_{1,2} = \eta/2 \pm \sqrt{V_0 + V_1 - 8mE/\hbar^2}$, where $\eta = \sqrt{V_1 + V_2 + 1/4}$, $L_\nu = \frac{1}{2}(\nu - 1)$ and $\nu = \sqrt{V_1 - V_2 + 1/4}$. Again, the poles of the Green function determine the energy-spectrum which coincidences with the one determined in (51), and the corresponding residua give the wave-functions expansions. We obtain ($k_1 = \frac{1}{2}(1+\nu)$, $k_2 = \frac{1}{2}(1-\eta)$, $\eta = \sqrt{-2m(E_n + B)/\hbar}$, $\nu = \sqrt{2m(B - E_n)/\hbar}$)

$$E_n = \sqrt[3]{\sqrt{D} + \frac{Q}{2}} - \sqrt[3]{\sqrt{D} - \frac{Q}{2}} + \frac{R}{3}, \quad (54)$$

$$\begin{aligned} R &= \frac{12\tilde{n}^2(\tilde{n}^2 + \lambda) - \lambda^2}{4\tilde{n}^2}, \\ T &= \frac{16\tilde{n}^2\lambda(A+C) - [\lambda^2 + B^2/4 + 4\tilde{n}^2(A+C)]^2}{8\tilde{n}^2\lambda(A+C+\lambda) - (\lambda^2 + 4\tilde{n}^2)(\lambda^2 + B^2/4 + 4\tilde{n}^2(A+C))}, \\ S &= \frac{4\tilde{n}^2}{2\tilde{n}^2}, \end{aligned} \quad (55)$$

$$\begin{aligned} \Psi_n^{(V_4)}(x) &= N_{E_n}^{(V_4)} \left[\frac{(2k_1 - 2k_2 - 2n - 1)n! \Gamma(2k_1 - n - 1)^{1/2}}{2! \Gamma(2k_2 + n) \Gamma(2k_1 - 2k_2 - n)} \right]^{1/2} \sqrt{\coth u} \\ &\quad \times \left(\sinh \frac{u}{2} \right)^{2k_2 - 1/2} \left(\cosh \frac{u}{2} \right)^{2n - 2k_1 + 3/2} P_n^{[2k_2 - 1, 2(k_1 - k_2 - n) - 1]} \left(\frac{2}{\cosh^2 \frac{u}{2}} - 1 \right) \end{aligned} \quad (56)$$

$$N_{E_n}^{(V_4)} = \left(\frac{(2\lambda - E_n) \sqrt{E_n^2 - B^2} (\sqrt{A - E_n} + \tilde{n})^{1/2}}{\tilde{n} [3(R/3 - E_n)^2 - R^2/3 + S]} \right)^{1/2}. \quad (57)$$

Bound states exist if $A < 0$, $0 < B < |A|$, and the number N_{Max} of bound states is found by requiring $|E_n| > B$. The scattering states have the form [$\kappa = \frac{1}{2}(1+ip)$, $p \in \mathbb{R}$]

$$\Psi_p^{(V_4)}(u) = N_p^{(V_4)} \sqrt{\coth u} \left(\cosh \frac{u}{2} \right)^{2n - 2k_1 + 3/2} \left(\sinh \frac{u}{2} \right)^{2k_2 - 1/2}$$

$${}_2F_1\left(k_1 + k_2 - \kappa, k_2 - k_1 - \kappa + 1; 2k_2; \tanh^2 \frac{u}{2}\right), \quad (58)$$

$$N_p^{(\kappa)} = \frac{1}{\Gamma(2k_2)} \sqrt{\frac{p \sinh 2\pi p}{2\pi^2}} \left[\Gamma(k_1 + k_2 - \kappa) \Gamma(-k_1 + k_2 + \kappa) \right. \\ \left. \times \Gamma(k_1 + k_2 + \kappa - 1) \Gamma(-k_1 + k_2 - \kappa + 1) \right]^{1/2}. \quad (59)$$

The results concerning the bound states coincide with [17].

4 Summary.

In this article the path integral treatments of four so-called "conditionally exactly solvable" potentials have been presented. Our approach showed that the path integral in the present cases is far more superior in comparison to other methods. In spite of the fact that the bound state energy-levels could not be stated in closed form in the first two cases, closed form solution in terms of the Green function were still possible. The poles of the Green functions (transcendental equations in terms of parabolic cylinder functions) gave the bound state energy-levels, the cuts provided the scattering states.

In the second set of potentials the bound-state solutions are determined by a cubic equation which considerably complicated the expressions analytically. In each two cases the bound-state energy-levels with the wave-functions and the scattering solutions could be obtained.

The results are rather satisfactory. In the Schrödinger approach, may it be the usual study in non-relativistic quantum mechanics or a super-symmetric investigation, fails to see the potential problem as a whole. In comparison, the path integral provides comprehensive information, about the propagator, in case it can be explicitly computed, the Green function which its poles and cuts, the bound-state wave-functions, the continuous spectrum, and the necessary boundary conditions. We also see that the interplay of various techniques was needed to obtain the proper solutions. In all four cases a space-time transformation was essential. In the first two cases, it was not only necessary to know about the path integral solution of a (shifted) harmonic oscillator or the linear potential, but even more essential was to know, how to incorporate explicit boundary conditions into the path integral. In the two modified Rosen-Morse oscillators two special path integral solutions had to be known which are in turn based on the path integral solution of the modified Pöschl-Teller potential.

It is to be expected that in the future some other specific path integral solutions can be found by relating known problems to more complicated (and therefore more realistic) potentials which can incorporate more parameters. These considerations can, of course, also be extended to two and three-dimensions and will be among other things presented elsewhere.

References

- [1] Feynman R P (1948) "Space-Time Approach to Non-Relativistic Quantum Mechanics", *Rev.Mod.Phys.* **20** 367-387
- [2] Feynman R P and Hibbs A (1965) "Quantum Mechanics and Path Integrals" (*McGraw Hill*, New York)
- [3] Peak D and Inomata A (1969) "Summation Over Feynman Histories in Polar Coordinates", *J.Math.Phys.* **10** 1422-1428
- [4] Govaerts M J (1975) "Path-Integral Evaluation of a Nonstationary Calogero Model", *J.Math.Phys.* **16** 720-723
- [5] Inomata A (1994) "Reckoning of the Besselian Path Integral", in "On Klauder's Path: A Field Trip", 99-106, eds.: Emch G G, Hergfeldt G C and Streit L (*World Scientific*, Singapore)
- [6] Böhm M and Junker G (1986) "The SU(1,1) Propagator as a Path Integral Over Noncompact Groups", *Phys.Lett. A* **117** 375-380
- [7] Böhm M and Junker G (1987) "Path Integration Over Compact and Noncompact Rotation Groups", *J.Math.Phys.* **28** 1978-1994
- [8] Grosche C (1995) "Path Integrals, Hyperbolic Spaces, and Selberg Trace Formulae", *DESY Report*, DESY 95-021, Habilitationsschrift, Hamburg University, 228pp., to be published by *World Scientific*
- [9] Grosche C and Steiner F (1995) "Table of Feynman Path Integrals", to appear in: *Springer Tracts in Modern Physics*
- [10] Grosche C and Steiner F (1993) "Classification of Solvable Feynman Path Integrals", in the Proceedings of the IV. International Conference on "Path Integrals from meV to MeV", Tutzing, Germany 1992, 276-288, eds.: Grabert H et al. (*World Scientific*, Singapore)
- [11] Grosche C and Steiner F (1995) "How to Solve Path Integrals in Quantum Mechanics" *J.Math.Phys.* **36** 2354-2385
- [12] Inomata A and Junker G (1993) "Path Integrals and Lie Groups", In "Noncompact Lie Groups and Some of Their Applications", San Antonio, Texas, USA, 1993, 199-224; eds.: Tanner E A and Wilson R (*Kluwer Academic Publishers*, Dordrecht)
- [13] Inomata A, Kuratsuji H and Gerry C (1992) "Path Integrals and Coherent States of SU(2) and SU(1,1)" (*World Scientific*, Singapore)
- [14] Lucht M W and Jarvis P D (1993) "Quasireactly Solvable Problems in the Path-Integral Formalism", *Phys.Rev. A* **47** 817-822
- [15] De Souza Dutra A (1993) "Conditionally Exactly Soluble Quantum Potentials", *Phys.Rev. A* **47** R2435-R2437
- [16] de Souza Dutra A and Boschi-Filho H (1994) "so(2,1) Lie algebra and the Green Functions for the Conditionally Exactly Solvable Potentials", *Phys.Rev. A* **50** 2915-2920

- [17] Dutt R, Khare A and Varshni Y P (1995) "New Classes of Conditionally Exactly Solvable Potentials in Quantum Mechanics", *J. Phys. A: Math. Gen.* **28** L107-L113
- [18] Nag N, Roychoudhury R and Varshni Y P (1994) "Conditionally Exactly Soluble Potentials and Supersymmetry", *Phys. Rev. A* **49** 5098-5099
- [19] Stillinger F H (1979) "Solution of a Quantum Mechanical Eigenvalue Problem with Long Range Potentials", *J. Math. Phys.* **20** 1891-1895
- [20] Song X (1979) "An Effective Quark-Antiquark Potential for Both Heavy and Light Mesons", *J. Phys. G: Nucl. Phys.* **17** 49-55
- [21] Eckart C (1930) "The Penetration of a Potential Barrier by Electrons", *Phys. Rev.* **35** 1303-1309
- [22] Rosen N and Morse P M (1932) "On the Vibrations of Polyatomic Molecules", *Phys. Rev.* **42** 210-217
- [23] Crandall R E (1983) "Exact Propagator for Reflectionless Potentials", *J. Phys. A: Math. Gen.* **16** 3005-3011
- [24] Manning M F and Rosen N (1933) "A Potential Function for the Vibrations of Diatomic Molecules", *Phys. Rev.* **44** 953
- [25] Bondjeddaa T, Chétouani L, Guéchi L and Hammann T F (1991) "Path Integral Treatment for a Screened Potential", *J. Math. Phys.* **32** 441-446
- [26] Hulthén L (1942) "Über die Eigenlösungen der Schrödingergleichung des Deuterons", *Arkiv Matematik, Astron. Fysik* **28** no.5
- [27] Grosche C (1993) "Path Integral Solution of Scarf-Like Potentials", *Nuovo Cimento B* **108** 1365-1376
- [28] Pertsch D (1990) "Exact Solution of the Schrödinger Equation for a Potential Well with Barrier and Other Potentials", *J. Phys. A: Math. Gen.* **23** 4145-4145
- [29] Papp E (1993) "Derivation of Quantum-Mechanical β Functions From Ground-State Energies", *Phys. Lett. A* **178** 231-235
- [30] Duru I H and Kleinert H (1979) "Solution of the Path Integral for the H-Atom", *Phys. Lett. B* **84** 185-188. (1982) "Quantum Mechanics of H-Atoms From Path Integrals", *Fortschr. Phys.* **30** 401-435
- [31] Fischer W, Leschke H and Müller P (1992) "Changing Dimension and Time: Two Well-Founded and Practical Techniques for Path Integration in Quantum Physics", *J. Phys. A: Math. Gen.* **25** 3835-3853. (1993) "Path Integration in Quantum Physics by Changing the Drift of the Underlying Diffusion Process: Application of Legendre Processes", *Ann. Phys. (N.Y.)* **227** 206-221
- [32] Grosche C and Steiner F (1987) "Path Integrals on Curved Manifolds", *Zeitschr. Phys. C* **36** 699-714
- [33] Kleinert H (1990) "Path Integrals in Quantum Mechanics, Statistics and Polymer Physics" (*World Scientific*, Singapore)
- [34] Pak N K and Sökmen I (1984) "General New-Time Formalism in the Path Integral", *Phys. Rev. A* **30** 1629-1635
- [35] Pelster A and Wunderlin A (1992) "On the Generalization of the Duru-Kleinert-Propagator Transformations", *Zeitschr. Phys. B* **89** 373-386
- [36] Steiner F (1984) "Space-Time Transformations in Radial Path Integrals", *Phys. Lett. A* **106** 356-362
- [37] Storčák S N (1988) "Path Reparametrization in a Path Integral on a Finite-Dimensional Manifold", *Theor. Math. Phys.* **75** 610-618
- [38] Young A and DeWitt-Morette C (1986) "Time Substitution in Stochastic Processes as a Tool in Path Integration", *Ann. Phys.* **169** 140-166
- [39] Inomata A (1986) "Remarks on the Time Transformation Technique for Path Integration", in "Path Integrals From meV to MeV", Bielefeld 1985, 433-448, eds.: Gutzwiller M C et al. (*World Scientific*, Singapore). (1988) "Recent Developments of Techniques for Solving Nontrivial Path Integrals", in "Path Summation: Achievements and Goals", Trieste 1987, 114-146, eds.: Lundquist S et al. (*World Scientific*, Singapore). (1989) "Time Transformation Techniques in Path Integration", in "Path Integrals from meV to MeV", 112-146, eds.: Sa-yakanit V et al. (*World Scientific*, Singapore)
- [40] Landau L D and Lifshitz (1958) "Quantum Mechanics", pp.188-121 (*Addison-Wesley*, Reading, Massachusetts)
- [41] Inomata A (1984) "Alternative Exact-Path-Integral-Treatment of the Hydrogen Atom", *Phys. Lett. A* **101** 253-257
- [42] Steiner F (1984) "Exact Path Integral Treatment of the Hydrogen Atom", *Phys. Lett. A* **106** 363-367
- [43] Grosche C (1993) "Path Integration via Summation of Perturbation Expansions and Application to Totally Reflecting Boundaries and Potential Steps", *Phys. Rev. Lett.* **71** 1-4
- [44] Grosche C (1993) " δ -Function Perturbations and Boundary Problems by Path Integration", *Ann. Physik* **2** 557-589
- [45] Abramowitz M and Stegun I A (1984) "Pocketbook of Mathematical Functions" (*Verlag Harry Deutsch*, Frankfurt/Main)
- [46] Grosche C (1989) "Path Integral Solution of a Class of Potentials Related to the Pöschl-Teller Potential", *J. Phys. A: Math. Gen.* **22** 5073-5087
- [47] Kleinert H and Mustapic I (1992) "Summing the Spectral Representations of Pöschl-Teller and Rosen-Morse Fixed-Energy Amplitudes", *J. Math. Phys.* **33** 643-662