

## Derivations on Murray–von Neumann algebras

A. F. Ber, K. K. Kudaybergenov, and F. A. Sykochev

For a given algebra  $\mathcal{A}$ , a linear operator  $D: \mathcal{A} \rightarrow \mathcal{A}$  is called a *derivation* if  $D$  satisfies the Leibniz rule, that is,  $D(xy) = D(x)y + xD(y)$  for all  $x, y \in \mathcal{A}$ . Each element  $a \in \mathcal{A}$  implements a derivation  $\text{ad}(a)$  on  $\mathcal{A}$  defined by  $\text{ad}(a)(x) = [a, x] = ax - xa$ ,  $x \in \mathcal{A}$ . Such derivations are said to be *inner*.

Let  $H$  be a Hilbert space, let  $B(H)$  be the  $*$ -algebra of all bounded linear operators on  $H$ , and let  $\mathcal{M}$  be a von Neumann algebra, that is, a weakly closed unital  $*$ -subalgebra of  $B(H)$  (for details see [11]).

A densely defined closed linear operator  $x: \text{dom}(x) \rightarrow H$  (here the domain  $\text{dom}(x)$  of  $x$  is a linear subspace of  $H$ ) is said to be *affiliated* with  $\mathcal{M}$  if  $yx \subset xy$  for all  $y$  in the commutant  $\mathcal{M}'$  of the algebra  $\mathcal{M}$ .

Denote the set of all projections in  $\mathcal{M}$  by  $P(\mathcal{M})$ . Recall that two projections  $e, f \in P(\mathcal{M})$  are said to be *equivalent* if there exists an element  $u \in \mathcal{M}$  such that  $u^*u = e$  and  $uu^* = f$ . A projection  $p \in \mathcal{M}$  is said to be *finite* if the conditions  $q \leq p$  and  $q \sim p$  imply that  $q = p$ . A linear operator  $x$  affiliated with  $\mathcal{M}$  is said to be *measurable* with respect to  $\mathcal{M}$  if  $\chi_{(\lambda, +\infty)}(|x|)$  is a finite projection for some  $\lambda > 0$ . (Here  $\chi_{(\lambda, +\infty)}(|x|)$  is the spectral projection of  $|x|$  corresponding to the interval  $(\lambda, +\infty)$ ). We denote the set of all measurable operators by  $S(\mathcal{M})$ .

Let  $x, y \in S(\mathcal{M})$ . It is well known that  $x + y$  and  $xy$  are densely-defined and preclosed operators. Moreover, the closures of  $x + y$ ,  $xy$ , and  $x^*$  are also in  $S(\mathcal{M})$ . The closures of  $x + y$  and  $xy$  are called the *strong sum* and *strong product*, respectively. When equipped with these operations,  $S(\mathcal{M})$  becomes a unital  $*$ -algebra over  $\mathbb{C}$  (see [12] and [15]). It is clear that  $\mathcal{M}$  is a  $*$ -subalgebra of  $S(\mathcal{M})$ . In the case when  $\mathcal{M}$  is a finite von Neumann algebra,  $S(\mathcal{M})$  is referred to as the *Murray–von Neumann algebra* associated with  $\mathcal{M}$  [9].

The hypothesis that all derivations of the algebra  $S(\mathcal{M})$  associated with a von Neumann algebra  $\mathcal{M}$  of type II are inner was first conjectured by Ayupov (see [2] and [3]). As Kadison and Liu noted in [10], pp. 210–211 (see also [9], p. 2090), for type II<sub>1</sub> algebras “the complete cohomological result would say that each derivation of  $S(\mathcal{M})$  is inner. . . . The authors *strongly* feel that this is true; but it is still open”. In this paper we announce the complete solution of this cohomological problem for type II<sub>1</sub> von Neumann algebras  $\mathcal{M}$ .

**Theorem 1.** *Let  $\mathcal{M}$  be a type II<sub>1</sub> von Neumann algebra, and let  $S(\mathcal{M})$  be the Murray–von Neumann algebra of all operators affiliated with  $\mathcal{M}$ . Then any derivation of  $S(\mathcal{M})$  is inner.*

In fact, we prove that any derivation of  $S(\mathcal{M})$  is continuous in the topology of convergence in measure on  $S(\mathcal{M})$ , and then we use known results from [4], [5], and [7] giving us that any derivation of  $S(\mathcal{M})$  which is continuous in this topology is necessarily inner.

When  $\mathcal{M}$  is an arbitrary von Neumann algebra, Sankaran [14] and Yeadon [16] introduced the algebra  $LS(\mathcal{M})$  of locally measurable operators affiliated with  $\mathcal{M}$ , with the operations of strong sum and strong multiplication. An operator  $x$  affiliated with  $\mathcal{M}$  is said to be locally measurable (with respect to  $\mathcal{M}$ ) if there is a sequence  $\{z_n\}_{n=0}^\infty \subset Z(\mathcal{M})$  of projections in the centre  $Z(\mathcal{M})$  of  $\mathcal{M}$  such that  $z_n \uparrow \mathbf{1}$ ,  $z_n(H) \subset \text{dom}(x)$ , and  $xz_n \in S(\mathcal{M})$  for all  $n \geq 0$ .

Using Theorem 1 and results from [1], [6], and [7], we obtain a necessary and sufficient condition for the existence of a non-inner derivation of the algebras  $S(\mathcal{M})$  and  $LS(\mathcal{M})$ . This result provides a complete answer to the problem posed by Ayupov in [2] and an adaptation of the celebrated Kadison–Sakai theorem [8], [13] to algebras of unbounded operators.

**Corollary 2.** *Let  $\mathcal{M}$  be an arbitrary von Neumann algebra. Then the following assertions are equivalent:*

- (a) *any derivation of  $LS(\mathcal{M})$  (of  $S(\mathcal{M})$ ) is inner;*
- (b) *a type  $I_{\text{fin}}$  direct summand of  $\mathcal{M}$  is atomic.*

In other words, the algebra  $S(\mathcal{M})$  (or  $LS(\mathcal{M})$ ) admits non-inner derivations if and only if the type  $I_{\text{fin}}$  direct summand of  $\mathcal{M}$  is non-trivial and non-atomic.

The authors thank J. Huang, G. Levitina, and D. Zanin for a detailed check of the results presented above and for useful comments.

## Bibliography

- [1] S. Albeverio, Sh. A. Ayupov, and K. K. Kudaybergenov, *J. Funct. Anal.* **256**:9 (2009), 2917–2943.
- [2] Ш. А. Аюпов, *Докл. АН РУз*, 2000, № 3, 14–17. [Sh. A. Ayupov, *Dokl. Uzbek Akad. Nauk*, 2000, no. 3, 14–17.]
- [3] Sh. A. Ayupov and K. K. Kudaybergenov, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **13**:2 (2010), 305–337.
- [4] Sh. Ayupov and K. Kudaybergenov, *J. Math. Anal. Appl.* **408**:1 (2013), 256–267.
- [5] А. Ф. Бер, *Матем. заметки* **93**:5 (2013), 658–664; English transl., A. F. Ber, *Math. Notes* **93**:5 (2013), 654–659.
- [6] A. F. Ber, V. I. Chilin, and F. A. Sukochev, *Extracta Math.* **21**:2 (2006), 107–147.
- [7] A. F. Ber, V. I. Chilin, and F. A. Sukochev, *Proc. Lond. Math. Soc.* (3) **109**:1 (2014), 65–89.
- [8] R. V. Kadison, *Ann. of Math.* (2) **83**:2 (1966), 280–293.
- [9] R. V. Kadison and Z. Liu, *Proc. Natl. Acad. Sci. USA* **111**:6 (2014), 2087–2093.
- [10] R. V. Kadison and Z. Liu, *Math. Scand.* **115**:2 (2014), 206–228.
- [11] R. V. Kadison and J. R. Ringrose, *Fundamentals of the theory of operator algebras*, vol. II: *Advanced theory*, Pure Appl. Math., vol. 100, Part 2, Academic Press, Inc., Orlando, FL 1986, i–xiv, 399–1074 pp.
- [12] E. Nelson, *J. Funct. Anal.* **15**:2 (1974), 103–116.
- [13] S. Sakai, *Ann. of Math.* (2) **83**:2 (1966), 273–279.
- [14] S. Sankaran, *J. Lond. Math. Soc.* **34**:3 (1959), 337–344.
- [15] I. E. Segal, *Ann. of Math.* (2) **57**:3 (1953), 401–457.

- [16] F. J. Yeadon, *Proc. Cambridge Philos. Soc.* **74**:2 (1973), 257–268.

**Aleksei F. Ber**

National University of Uzbekistan,  
Tashkent, Uzbekistan

*E-mail:* [aber1960@mail.ru](mailto:aber1960@mail.ru)

Presented by A. G. Sergeev

Accepted 20/JUN/19

Translated by F. SUKOCHEV

**Karimbergen K. Kudaybergenov**

Department of Mathematics,  
Karakalpak State University,  
Nukus, Uzbekistan

*E-mail:* [karim2006@mail.ru](mailto:karim2006@mail.ru)

**Fedor A. Sukochev**

School of Mathematics and Statistics,  
University of New South Wales,  
Kensington, Australia

*E-mail:* [f.sukochev@unsw.edu.au](mailto:f.sukochev@unsw.edu.au)