

Optical solitons with M-truncated derivative and conservation laws for NLSE equation which describe pseudospherical surfaces

Abdullahi Yusuf¹ and Mustafa Inc^{2,3} 

¹Federal University Dutse, Science Faculty, Department of Mathematics, 7156 Jigawa, Nigeria

²Firat University, Science Faculty, Department of Mathematics, 23119 Elazig, Turkey

E-mail: yusufabdullahi@fud.edu.ng and minc@firat.edu.tr

Received 26 August 2019, revised 15 October 2019

Accepted for publication 29 October 2019

Published 6 February 2020



Abstract

The present work analyzes the optical solitons with M-truncated and conservation laws (CLs) of the nonlinear Schrodinger equation that explain pseudospherical surfaces. The well-known integration scheme which is generalized Bernoulli sub-ODE method is utilized to construct such optical soliton solutions. For the successful existence of the solutions, the constraints conditions have been presented. The discussion for the physical features of the obtained solutions is reported. Moreover, we establish CLs for the equation under consideration by means of multiplier approach.

Keywords: generalized Bernoulli sub-ODE method, M-truncated derivative, optical solitons

(Some figures may appear in colour only in the online journal)

1. Introduction

In both theoretical and applied sciences, the research of physical events through mathematical models is an important component. Such models often lead to nonlinear systems and some prototypical equations in amazingly big numbers of instances. Solitons are an intrinsically nonlinear phenomenon and their history is closely linked to the growth of nonlinear wave equations theory. In latest decades there has been a steady increase in interest in optical solitons. The field has significant potential for technological applications and poses many interesting study issues from both a basic and an applied perspective. Many fields of science and engineering employ a nonlinear partial differential equations (PDEs) to determine the characteristic of these phenomena. In the scrutiny of PDEs, a lot of authors give a priority in constructing soliton and exact solutions.

NLSEs are a PDEs with exciting performances for many years. This is because of its expansive spectrum of applicability. There are a types of NLSEs that can be employed to interpret real phenomena in various domains among which is

Bose–Einstein, deliquescence [1, 2], nonlinear optics [3–5], fluid dynamics [6] to mention a few. In the present time, many studies can be found in this regard [7–13]. Optical solitons are localized electromagnetic waves that spread in nonlinear dispersive media with the unchanged intensity as a result of the effects of the stability for the dispersion and nonlinearity [14–23].

Furthermore, CLs generate numerous ideas on the systems of equations that are modelled by PDEs. The CLs have been very important aspects in the investigation of system of PDEs, because the integrability, internal properties, existence and uniqueness of equations can be reached through their CLs. Several schemes have been suggested in the literature for the constructions of CLs of a system of a particular equations [24–29].

Pseudospherical surfaces is described by a differential system for a 2-vector-valued function $U(x, t)$ if it possesses some certain conditions with regard to the existence of smooth real functions F_{ij} , $i \in [1, 3]$, $j \in [1, 2]$, depending on U only and a finite number of derivatives, thus we have

$$\sigma_i = F_{i1}dx + F_{i2}dt, \quad i \in [1, 3], \quad (1)$$

hold for the following

$$d\sigma_1 = \sigma_3 \wedge \sigma_2, \quad d\sigma_2 = \sigma_1 \wedge \sigma_3, \quad d\sigma_3 = \sigma_1 \wedge \sigma_2. \quad (2)$$

³ Author to whom any correspondence should be addressed.

A differential equation for a real valued function $U(x, t)$ is said to be integrable kinematically if it has the integrability property of a family of one-parameter for linear situations [30–36]

$$V_x = P(\gamma)V, V_t = Q(\gamma)V, \quad (3)$$

with $P(\gamma)$ and $Q(\gamma)$ depicting the $SL(2, R)$ -valued functions for x, t . Therefore, an equation is said to be integrable kinematically if it has the same property with the zero curvature property

$$\frac{\partial P(\gamma)}{\partial t} - \frac{\partial Q(\gamma)}{\partial x} + [P(\gamma), Q(\gamma)] = 0, \quad (4)$$

in which $\text{tr}P(\gamma) = \text{tr}Q(\gamma) = 0$, for every γ . Additionally, a differential equation is said to be strictly integrable kinematically if its integrable kinematically and diagonal entries of the matrix $P(\gamma)$ presented above are γ and $-\gamma$.

Many renowned physicists and applied mathematicians have been contributing to some or all of the aforementioned fields over the past decades. The involved-engineers, physicists, and mathematicians-are still operating within their own limits to a big extent. One of the aims of this article is to establish some optical soliton solutions and CIs, hoping to stimulate future communication with distinct backgrounds among scientists. The optical solitons are computed via a well-known analytical technique with m-truncated derivative, while the CIs are constructed by means of multiplier technique

The rest of the paper is organized in the following direction: In section two we state the description of m-truncated derivative and the form of the governing equation, in section three we establish some optical solitons solutions for the governing equation. Section four provides the CIs for the governing equation by means of multiplier approach and finally the paper is concluded in section five.

2. Truncated M-fractional derivative

We define the truncated Mittag–Leffler function of one parameter by

$${}_iE_\beta(z) = \sum_{k=0}^i \frac{z^k}{\Gamma(\beta k + 1)}. \quad (5)$$

Truncated M-fractional derivative (TMD) is a fractional derivative that has been introduced in [37]. This derivative has expunged the obstacles with the existing derivatives. It is defined in the following definition.

Definition 2.1. Assume that $f: (0, \infty) \rightarrow \mathbb{R}$, the TMD of f with order γ exhibited ${}_iT_M^{\gamma, \beta}$ is given by

$${}_iT_M^{\gamma, \beta} f(\tau) = \lim_{\epsilon \rightarrow 0} \frac{f(\tau + {}_iE_\beta(\epsilon\tau^{-\gamma})) - f(\tau)}{\epsilon}, \quad (6)$$

for $\tau > 0$, and ${}_iE_\beta \gamma \in (0, 1)$, $\beta > 0$ is a truncated Mittag–Leffler function of one parameter, as defined in (5). Note that, if f is γ -differentiable in some open interval $(0, a)$, $a > 0$, and

$\lim_{\tau \rightarrow 0^+} ({}_iT_M^{\gamma, \beta} f(\tau))$. Then we attain

$${}_iT_M^{\gamma, \beta} f(0) = \lim_{\tau \rightarrow 0^+} ({}_iT_M^{\gamma, \beta} f(\tau)). \quad (7)$$

Theorem 2.2. Surmise that $f: (0, \infty) \rightarrow \mathbb{R}$ is γ -differentiable for $\tau_0 > 0$, with $\gamma \in (0, 1]$, $\beta > 0$, then f is continuous at τ_0 .

Theorem 2.3. Let $0 < \gamma \leq 1$, $\beta > 0$, $a, b \in \mathbb{R}$, f, g , γ -differentiable, at a point $\tau > 0$. Then

- ${}_iT_M^{\gamma, \beta}(af + bg) = a{}_iT_M^{\gamma, \beta}(f) + b{}_iT_M^{\gamma, \beta}(f)$, $a, b \in \mathbb{R}$
- ${}_iT_M^{\gamma, \beta}(t^\mu) = \mu\tau^{\mu-\gamma}$, $\mu \in \mathbb{R}$
- ${}_iT_M^{\gamma, \beta}(fg) = f_iT_M^{\gamma, \beta}(g) + g_iT_M^{\gamma, \beta}(f)$,
- ${}_iT_M^{\gamma, \beta}\left(\frac{f}{g}\right) = \frac{g_iT_M^{\gamma, \beta}(f) - f_iT_M^{\gamma, \beta}(g)}{g^2}$,
- If f is differentiable, then ${}_iT_M^{\gamma, \beta}(f)(\tau) = \frac{\tau^{1-\gamma}}{\Gamma(\beta+1)} \frac{df}{d\tau}$,
- ${}_iT_M^{\gamma, \beta}(fog)(\tau) = f'(g(\tau)){}_iT_M^{\gamma, \beta}g(\tau)$, for f differentiable at g .

2.1. Governing equation

Assume that M^2 is a differentiable surface [38], parameterized by x, t . Let also, the following

$$\begin{aligned} \sigma_1 &= 2\sigma dx + (-4\gamma\sigma + 2V_x)dt, \\ \sigma_2 &= 2\gamma dx + (2(V^2 + \sigma^2) - 4\gamma^2)dt, \\ \sigma_3 &= -2Vdx + (4\gamma V + 2\sigma_x)dt. \end{aligned} \quad (8)$$

Then M^2 is a PSS if and only if U satisfies the NLSE

$${}_i^E\mathcal{D}_{M,t}^{\gamma, \beta}U + {}_0^E\mathcal{D}_{M,x}^{2\gamma, \beta} + 2|U|^2U = 0, \quad (9)$$

where $\mathcal{D}_{M,x}^{2\gamma, \beta}$ denotes the m-truncated derivative. To obtain a travelling wave solutions for equation (9), we substitute $U(x, t) = V + i\sigma$ in equation (9) to obtain

$$\begin{aligned} {}_0^E\mathcal{D}_{M,t}^{\gamma, \beta}V + {}_0^E\mathcal{D}_{M,x}^{2\gamma, \beta}\sigma + 2(V^2 + \sigma^2)\sigma &= 0, \\ -{}_0^E\mathcal{D}_{M,t}^{\gamma, \beta}\sigma + {}_0^E\mathcal{D}_{M,x}^{2\gamma, \beta}V + 2(V^2 + \sigma^2)V &= 0, \end{aligned} \quad (10)$$

where $\mathcal{D}_{M,x}^{2\gamma, \beta}$ denotes the m-truncated derivative. Plugging the transformation $V(x, t) = u(\xi)$, $\sigma(x, t) = v(\xi)$, $\xi = \frac{\lambda\Gamma(\beta+1)}{\gamma}(x^\gamma - kt^\gamma + c)$, where λ, k, c are real constants and will be found later, into equation (10), we have

$$\begin{aligned} -k\lambda u'(\xi) + 2v(\xi)(u(\xi)^2 + v(\xi)^2) + \lambda^2 v''(\xi) &= 0, \\ k\lambda v'(\xi) + \lambda^2 u''(\xi) + 2u(\xi)(u(\xi)^2 + v(\xi)^2) &= 0. \end{aligned} \quad (11)$$

3. Application

According to Bernoulli sub-ODE method [39], equation (11) has possessed the solution as comes next

$$u = a_0 + a_1\Phi(\xi), v = b_0 + b_1\Phi(\xi), \quad (12)$$

where $\Phi(\xi)$ satisfies

$$\Phi' = \mu\Phi^2 - \lambda\Phi. \quad (13)$$

where μ is a nonzero constant. Plugging equation (13) along with equation (12) into equation (11), we get

$$\begin{aligned} & 2(b_1\Phi(\xi) + b_0)((a_1\Phi(\xi) + a_0)^2 + (b_1\Phi(\xi) + b_0)^2) \\ & + a_1k\lambda\Phi(\xi)(\lambda - \mu\Phi(\xi)) + b_1\lambda^2\Phi(\xi)(\lambda - \mu\Phi(\xi)) \\ & \times (\lambda - 2\mu\Phi(\xi)) = 0, \\ & 2(a_1\Phi(\xi) + a_0)((a_1\Phi(\xi) + a_0)^2 + (b_1\Phi(\xi) + b_0)^2) \\ & + a_1\lambda^2\Phi(\xi)(\lambda - \mu\Phi(\xi))(\lambda - 2\mu\Phi(\xi)) \\ & + b_1k\lambda\Phi(\xi)(\mu\Phi(\xi) - \lambda) = 0. \end{aligned} \quad (14)$$

Collecting the terms in $\Phi^i (i = 0, 1, 2, 3, 4, 5, 6, 7)$, one reaches

$$\begin{aligned} & 2b_0(a_0^2 + b_0^2) = 0, \\ & b_1(2a_0^2 + 6b_0^2 + \lambda^4) + a_1(4a_0b_0 + k\lambda^2) = 0, \\ & 2b_1(a_1^2 + b_1^2 + \lambda^2\mu^2) = 0, \\ & a_1(4a_0b_1 - k\lambda\mu) \\ & + 2a_1^2b_0 + 3b_1(2b_0b_1 - \lambda^3\mu) = 0, \\ & 2a_0(a_0^2 + b_0^2) = 0, \\ & a_1(6a_0^2 + 2b_0^2 + \lambda^4) + b_1(4a_0b_0 - k\lambda^2) = 0, \\ & 2a_1(a_1^2 + b_1^2 + \lambda^2\mu^2) = 0, \\ & a_1(4b_0b_1 - 3\lambda^3\mu) \\ & + b_1(2a_0b_1 + k\lambda\mu) + 6a_0a_1^2 = 0. \end{aligned} \quad (15)$$

Solving (15), we obtain

$$\begin{aligned} & a_0 = a_0, \quad k = k, \quad \lambda = \lambda, \quad \mu = \mu, \quad a_1 = \frac{-4a_0^2\mu - k^2\mu}{4a_0\lambda}, \\ & b_0 = \frac{a_0k}{\lambda^2}, \quad b_1 = \frac{\lambda^2(a_1k + 2b_0\lambda\mu)}{k^2}. \end{aligned}$$

Using the obtained results, we reach the following dark and singular optical solitons

$$\begin{aligned} & V(x, t) = a_0 - \frac{4a_0^2 + k^2}{8a_0} \\ & \times \left(\tanh \left(\frac{\lambda}{2} \left(\frac{\lambda\Gamma(\beta + 1)}{\gamma} (x^\gamma - kt^\gamma + c) \right) \right) - 1 \right), \\ & \sigma(x, t) = \frac{a_0k}{\lambda^2} + \frac{\lambda^3(a_1k + 2b_0\lambda\mu)}{2\mu k^2} \\ & \times \left(\tanh \left(\frac{\lambda}{2} \left(\frac{\lambda\Gamma(\beta + 1)}{\gamma} (x^\gamma - kt^\gamma + c) \right) \right) - 1 \right), \end{aligned} \quad (16)$$

or

$$\begin{aligned} & V(x, t) = a_0 - \frac{4a_0^2 + k^2}{8a_0} \\ & \times \left(\coth \left(\frac{\lambda}{2} \left(\frac{\lambda\Gamma(\beta + 1)}{\gamma} (x^\gamma - kt^\gamma + c) \right) \right) - 1 \right), \\ & \sigma(x, t) = \frac{a_0k}{\lambda^2} + \frac{\lambda^3(a_1k + 2b_0\lambda\mu)}{2\mu k^2} \\ & \times \left(\coth \left(\frac{\lambda}{2} \left(\frac{\lambda\Gamma(\beta + 1)}{\gamma} (x^\gamma - kt^\gamma + c) \right) \right) - 1 \right). \end{aligned} \quad (17)$$

4. Results and discussion

The novel GBM and multiplier scheme have been employed to establish soliton solutions and CIs for the equation under consideration, respectively. We computed CIs because of its importance in determining the integrability of differential equation and in providing partial information on the existence of a particular movement, even if the equations are too complicated to require a complete solution. The governing equation has possessed conserved vectors and internal properties which are presented in the section 5. These results has affirmed its integrability. We established soliton solutions because such solutions tend to play a significant role in explaining various phenomena and processes throughout the natural sciences. This is due to their capability to check the accuracy or estimate errors in statistical, asymptotic and estimated analytical solutions. To this aim, we get motivated and computed these results which will be of great benefit to the literature.

The physical and outlook views of the solutions obtained by GBM are portrayed by means of graphical representations. The types of solutions attained are the dark optical soliton reported in (16), the singular optical soliton reported in (17).


Dark optical soliton describes the solitary waves with less strength than the context, the different soliton solutions represent a solitary wave with discontinuous derivatives; an example of such solitary waves are compactons with finite (compact) support, and peaks with discontinuous first derivative peaks. Due to their efficiency and, of course, versatility in long distance optical communication, these types of solitary waves are extremely important.

It should be remembered that optical fibers are thin long strings of ultra-pure glass or plastic so that light can be transmitted from one end to the next without any attenuation or loss. In order to have a clear vision on the effect of parameters to the transmission of solitons, suitable values of parameters have been considered in the following graphs: In figure 1, we choose $a_0 = 0.2$, $\lambda = 1.2$, $k = 0.7$, $c = 0.1$, $a_1 = 0.5$, $b_0 = 0.9$, $\mu = 0.55$. And in figure 2, we choose $a_0 = 0.2$, $\lambda = 1.2$, $k = 0.9$, $c = 0.1$, $a_1 = 0.55$, $b_0 = 0.19$, $\mu = 0.5$.


5. Conservation laws via multiplier approach

Following the multipliers approach as depicted in [40], we construct the first-order multipliers $\Lambda^1(x, t, \sigma, V, \sigma_x, V_x, \sigma_t, V_t, \dots)$, $\Lambda^2(x, t, \sigma, V, \sigma_x, V_x, \sigma_t, V_t, \dots)$ for the equation under consideration and they are given by


$$\begin{aligned} \Lambda^1 &= \frac{1}{2}(c_1x + 2c_2)V - \frac{1}{2}(2c_1t + 2c_3)\sigma_x - c_4\sigma_t, \\ \Lambda^2 &= \frac{1}{2}(c_1x + 2c_2)\sigma + \frac{1}{2}(2c_1t + 2c_3)V_x - c_4V_t, \end{aligned} \quad (18)$$




(a) $V(x,t)$ with $\gamma = 0.5$, $\beta = 0.9$ for (16).



(b) $\sigma(x,t)$ with $\gamma = 0.5$ for (16).




(c) $V(x,t)$ with $\gamma = 0.5$, $\beta = 0.9$ for (17).




(d) $\sigma(x,t)$ with $\gamma = 0.5$, $\beta = 0.9$ for (17).


Figure 1. Physical features with suitable values of the parameters.




(a) $V(x,t)$ with $\gamma = 0.55$, $\beta = 0.99$ for (16).



(b) $\sigma(x,t)$ with $\gamma = 0.5$ for (16).



(c) $V(x,t)$ with $\gamma = 0.55$, $\beta = 0.99$ for (17).



(d) $\sigma(x,t)$ with $\gamma = 0.55$, $\beta = 0.99$ for (17).

Figure 2. Physical features with suitable values of the parameters.

where c_1, c_2 are some arbitrary constants. Subsequently, one attains the following fluxes:

$$\begin{aligned} T^t &= -\frac{1}{2}V^4c_4 - V^2\sigma^2c_4 + \frac{1}{4}V^2c_1x + \frac{1}{2}V^2c_2\sigma_x\sigma^3c_1t^2 \\ &\quad + \frac{1}{2}c_4\sigma_x^2 - 2\sigma^3\sigma_xc_3t - V\sigma_xc_1t - V\sigma_xc_3 + \frac{1}{2}c_4V_x^2, \\ T^x &= \frac{1}{2}Vc_1\sigma + Vc_3\sigma_t + Vc_3\sigma_t - \frac{1}{2}c_3V^4 \\ &\quad + 2\sigma^3\sigma_t c_3t - \frac{1}{2}V_xc_1\sigma_x + Vc_1\sigma_{tt} \\ &\quad - \sigma_xc_4\sigma_t + \sigma_xVc_2 - \frac{1}{2}\sigma_x^2c_1t + \frac{1}{2}\sigma_xVc_1x \\ &\quad - \frac{1}{2}\sigma_x^2c_3 - t\sigma^2c_1V^2 + \sigma_t\sigma^3c_1t^2 \\ &\quad - \frac{1}{2}c_1tV^4 - \sigma^2c_3V^2 - \frac{1}{2}V_x^2c_3 - V_xc_4V_t \\ &\quad - V_xc_2\sigma - \frac{1}{2}V_x^2c_1t. \end{aligned} \quad (19)$$

By situations disconnected without a constants, one reaches the following fluxes:

Case 1.

$$\Lambda^1 = \frac{1}{2}V_x - \sigma_{xt}, \quad \Lambda^2 = V_{xt} + \frac{1}{2}\sigma_x. \quad (20)$$

Subsequently, we obtain the following fluxes

$$\begin{aligned} T^t &= \frac{1}{4}V^2x - \sigma_x\sigma^3t^2 - V\sigma_{xt}, \\ T^x &= \frac{1}{2}V\sigma - \frac{1}{2}V_x\sigma_x + V\sigma_{tt} - \frac{1}{2}\sigma_x^2t + \frac{1}{2}V\sigma_{xx} - tV^2\sigma^2 \\ &\quad + \sigma_t\sigma^3t^2 - \frac{1}{2}tV^4 - \frac{1}{2}V_x^2t. \end{aligned} \quad (21)$$

Case 2.

$$\Lambda^1 = \sigma, \quad \Lambda^2 = V. \quad (22)$$

Subsequently, we obtain the following fluxes

$$\begin{aligned} T^t &= \frac{1}{2}V^2, \\ T^x &= V\sigma_x - \sigma V_x. \end{aligned} \quad (23)$$

Case 3.

$$\Lambda^1 = -\sigma_x, \quad \Lambda^2 = V_x. \quad (24)$$

Subsequently, we obtain the following fluxes

$$\begin{aligned} T^t &= -2\sigma^3\sigma_{xt} - V\sigma_x, \\ T^x &= V\sigma_t - \frac{1}{2}V^4 + 2\sigma^3\sigma_{tt} - \frac{1}{2}\sigma_x^2 - \sigma^2V^2 - \frac{1}{2}V_x^2. \end{aligned} \quad (25)$$

Case 4.

$$\Lambda^1 = -\sigma_t, \quad \Lambda^2 = V_t. \quad (26)$$

Subsequently, we obtain the following fluxes

$$\begin{aligned} T^t &= -\frac{1}{2}V^4 - \sigma^2V^2 + \frac{1}{2}\sigma_x^2 + \frac{1}{2}V_x^2, \\ T^x &= -V_tV_x - \sigma_t\sigma_x. \end{aligned} \quad (27)$$

6. Conclusion

In this work we have studied the optical solitons with M-truncated derivative and CIs for the NLSE which describe Pseudospherical Surfaces. The novel GBM have been utilized to extract such novel solutions. For the successful existence of the solutions, the constraints conditions have been reported. The discussion for the physical features of the obtained solutions have been presented. Moreover, the construction for the conservation laws have been done by means of multiplier approach.

ORCID iDs

Mustafa Inc  <https://orcid.org/0000-0003-4996-8373>

References

- [1] Bronski J C, Carr L D, Deconinck B and Kutz J N 2001 Bose-Einstein condensates in standing waves: the cubic nonlinear Schrodinger equation with a periodic potential *Phys. Rev. Lett.* **86** 1402–5
- [2] Trombettoni A and Smerzi A 2001 Discrete solitons and breathers with dilute Bose-Einstein condensates *Phys. Rev. Lett.* **86** 2353–6
- [3] Triki H and Biswas A 2011 Dark solitons for a generalized nonlinear Schrodinger equation with parabolic law and dual power nonlinearity *Math. Methods Appl. Sci.* **34** 958–62
- [4] Zhang L H and Si J G 2010 New soliton and periodic solutions of (1+2)-dimensional nonlinear Schrodinger equation with dual power nonlinearity *Commun. Nonlinear Sci. Numer. Simul.* **15** 2747–54
- [5] Biswas A 2001 A perturbation of solitons due to power law nonlinearity *Chaos Solitons Fractals* **12** 579–88
- [6] Nore C, Brachet M E and Fauve S 1993 Numerical study of hydrodynamics using the nonlinear Schrodinger equation *Physica D* **65** 154–62
- [7] Xi L X and Wang M L 2009 The (G'/G) -expansion method and traveling wave solutions for higher-order nonlinear Schrodinger equation *Appl. Math. Comput.* **208** 440–5
- [8] Tsang S C, Chiang K S and Chow K W 2004 Solitons interaction in two-core optical fiber *Opt. Commun.* **229** 431–9
- [9] Dai C, Wang Y and Zhang J 2010 Analytical spatiotemporal localization for the generalized (3 + 1)-dimensional nonlinear Schrodinger equation *Opt. Lett.* **35** 1437–9
- [10] Tian H, Tian J, Li Z and Zi J 2005 Optical solitary wave solutions in quadratic nonlinear media *Opt. Commun.* **247** 225–32

- [11] Huang D J, Li D S and Zhang H Q 2007 Explicit and exact traveling solutions for the generalized derivative Schrödinger equation *Chaos Solitons Fractals* **31** 586–93
- [12] Beitia B, Garcia J P, Vekslerchik V M V and Torres P J 2008 Lie symmetries and qualitative analysis and exact solutions of the nonlinear Schrödinger equation with inhomogeneous nonlinearities *Discrete Continuous Dyn. Syst. B* **9** 221–33
- [13] Biswas A and Konar S 2007 *Introduction to Non-kerr Law Optical Solitons* (Boca Raton, FL: CRC Press)
- [14] Eslami M, Zhou Q, Moshokoa S P, Biswas A, Belic M, Ekici M and Mirzazadeh M 2016 Optical soliton perturbation with fractional temporal evolution by first integral method with conformable fractional derivatives *Optik* **127** 10659–69
- [15] Zhou Q, Moshokoa S P, Biswas A, Belic M, Ekici M and Mirzazadeh M 2016 Solitons in optical metamaterials with fractional temporal evolution *Optik* **127** 10879–97
- [16] Mirzazadeh M, Eslami M, Vajargah B F and Biswas A 2014 Application of the first integral method to fractional partial differential equations *Indian J. Phys.* **88** 177–84
- [17] Sonomezoglu A, Eslami M, Zhou Q, Zerrad E, Biswas A, Belic M, Mirzazadeh M and Ekici M 2016 Optical solitons in nano-fibers with fractional temporal evolution *J. Comput. Theor. Nanosci.* **13** 5361–74
- [18] Sonomezoglu A, Ortakaya S, Eslami M, Biswas A, Mirzazadeh M and Ekici M 2016 Solitons solutions to a few fractional nonlinear evolution equations in shallow water wave dynamics *Eur. Phys. J., Plus* **131** 166–77
- [19] Vajargah B F, Mirzazadeh M, Eslami M and Biswas A 2014 Solitons and periodic solutions to a couple of fractional nonlinear evolution equations *Pramana* **82** 465–76
- [20] Zerrad E, Biswas A, Song M and Ahmed B 2013 Domain wall and bifurcation analysis of the klein-gordon zakharov-kuznetsov equation in (1+2)-dimensions with power law nonlinearity *Chaos* **23** 033115
- [21] Biswas A and Suarez P 2011 Exact 1-soliton solution of the Zakharov–Kuznetsov equation in plasmas with power law nonlinearity *Appl. Math. Comput.* **217** 7372–5
- [22] Zerrad E, Biswas A, Kohl R and Milovic D 2009 Optical solitons by Heas variational principle in a non-kerr law media *J. Infrared, Millim., Terahertz Waves* **30** 526–37
- [23] Biswas A, Morris R and Kara A H 2013 Soliton solution and conservation laws of the Zakharov-Kuznetsov equation in plasmas with power law nonlinearity *Nonlinear Anal.: Modelling Control* **18** 153–9
- [24] Noether E 1918 Invariante variationsprobleme, königliche gesellschaft der wissenschaften zu göttingen, nachrichten *Math.-Phys. Kl. Heft* **2** 235–57
Noether E 1971 *Transp. Theory Stat. Phys.* **1** 186–207 (English transl.)
- [25] Ibragimov N H 2007 A new conservation theorem *J. Math. Anal. Appl.* **333** 311–28
- [26] Anco S C and Bluman G 1997 Direct construction of conservation laws from field equations *Phys. Rev. Lett.* **78** 2869–73
- [27] Inc M, Aliyu A I and Yusuf A 2017 Dark optical, singular solitons and conservation laws to the nonlinear Schrödinger's equation with spatio-temporal dispersion *Mod. Phys. Lett. B* **31** 1750163
- [28] Aliyu A I, Tchier F, Inc M, Yusuf A and Baleanu D 2018 Dynamics of optical solitons, multipliers and conservation laws to the nonlinear Schrödinger equation in (2+1)-dimensions with non-Kerr law nonlinearity *J. Mod. Opt.* **66** 136–42
- [29] Aliyu A I, Inc M, Yusuf A and Baleanu D 2018 Optical solitary waves and conservation laws to the (2 + 1)-dimensional hyperbolic nonlinear Schrödinger equation *Mod. Phys. Lett. B* **32** 1850373
- [30] Beals R, Rabelo M and Tenenblat K 1989 Backlund transformations and inverse scattering solutions for some pseudospherical surface equations *Stud. Appl. Math.* **81** 125–51
- [31] Cavalcante J A and Tenenblat K 1988 Conservation laws for nonlinear evolution equations *J. Math. Phys.* **29** 1044–9
- [32] Chadan K and Sabatier P C 1977 *Inverse Problems in Quantum Scattering Theory, Texts and Monographs in Physic* (New York, NY, USA: Springer)
- [33] Chern S S and Tenenblat K 1986 Pseudospherical surfaces and evolution equations *Stud. Appl. Math.* **74** 55–83
- [34] Khater A H, Callebaut D K and Sayed S M 2004 Conservation laws for some nonlinear evolution equations which describe pseudo-spherical surfaces *J. Geom. Phys.* **51** 332–52
- [35] Reyes E G 2000 Conservation laws and Calapso-Guichard deformations of equations describing pseudo-spherical surfaces *J. Math. Phys.* **41** 2968–89
- [36] Reyes E G 2001 *On Geometrically Integrable equations and Hierarchies of Pseudo-spherical Type in The Geometrical Study of Differential equations (Washington, DC, 2000)*, Vol. 285 of *Contemporary Mathematics* (Providence, RI, USA: American Mathematical Society) pp 145–55
- [37] Sousa J V D C and de Oliveira E C 2018 A new truncated M-fractional derivative type unifying some fractional derivative types with classical properties *Int. J. Anal. Appl.* **16** 83–96
- [38] Sayed S M, Elhamahmy O O and Gharib G M 2008 Travelling wave solutions for the kdv-burgers-kuramoto and nonlinear Schrödinger equations which describe pseudospherical surfaces *J. Appl. Math.* **2008** 576783
- [39] Taghizadeh N and Najand M 2012 The generalized Bernoulli sub-ODE method *Math. Aeterna* **2** 487
- [40] Buhe E and Bluman G W 2015 Symmetry reductions, exact solutions and conservation laws of the generalized Zakharov equations *J. Math. Phys.* **56** 101501