

# Analytical Approximate Solutions of Non Linear Partial Differential Equations using VIM, VIADM and New Modified KVIADM

Shruti S Sheth<sup>1</sup> & Dr Twinkle R Singh<sup>2</sup>

<sup>1</sup>Assistant Professor, Sarvajanik College of Engineering & Technology, Dr. R.K.Desai Marg, Opp. Mission Hospital, Athwalines, Suart-395001, Gujarat, India.

shruti.sheth@scet.ac.in

<sup>2</sup>Assistant Professor, Applied Mathematics & Humanities Department, Sardar Vallabhbhai National Institute of Technology, Surat-395007, Gujarat, India.

twinklesingh.svnit@gmail.com

**Abstract.** This paper examines the Analytical Approximate Solutions of the Non Linear Partial Differential Equations such as Non Linear Wave Equations and In viscid Burgers' Equation using Variational Iteration Method (VIM), Variational Iteration Adomian Decomposition Method (VIADM) and New Modified Kamal Variational Iteration Adomian Decomposition Method (New MKVIADM). VIM is a powerful tool to solve the differential equations which gives fast consecutive approximations without any conditional assumptions or any further transformations which may change the physical behaviour of the problem. Adomian Decomposition method is also an efficient method which handles the linear and non linear differential and integral equations with Initial and Boundary Conditions. It provides an efficient numerical solution in the form of an infinite series which is obtained iteratively. It usually converges to the exact solution using Adomian polynomials. Kamal Transform is a very recent new arrival of an integral transform which is commanding to solve the linear initial value problems. To check the efficiency of these methods, we have illustrated two non linear wave equations and one In Viscid Burgers' equation. Objective of this paper is that, how rapidly these methods converge to the exact solution in the closed form, in the given domain for the given initial conditions, and still how it sustains the high accuracy and precision. Our aim in this paper is try to employ the combination of three different kinds of methods. The strategy of the methods is outlined and in view of the convergence of the methods and to show how it fulfils the objectives.



## 1. Introduction

To solve these equations we have used the recent analytical approximate methods such as VIM , Combination of VIM and ADM that is VIADM and over VIADM we have applied new arrival of integral transform-Kamal Transform, that is New Modified Kamal Variational Iteration Adomian Decomposition Method.

### 1.1 Variational Iteration Method

In 1998, Variational Iteration Method has been developed and used by J. Huen He, to study and to solve the Non linear Partial Differential Equations. VIM is used to handle the both differential and Integral equations. VIM gives very rapidly convergent consecutive approximations of the exact solution, if such a solution exists, in the given domain, otherwise only some approximations can be used for numerical purposes only [4-7]. VIM effectively and accurately has been used by many authors herein and elsewhere.

### 1.2 Adomian Decomposition Method

In the 1980's, George Adomian introduced a new powerful method for solving nonlinear functional equations. Since then, this method has been known as the Adomian decomposition method (ADM) [2,3]. ADM is based on a decomposition of a solution of a nonlinear operator equation in a series of functions. Each term of the series is obtained from a polynomial generated from an expansion of an analytic function into a power series. The non linear part is expressed in terms of the Adomian polynomials. The initial or boundary condition and the terms that contain the independent variables will be considered as the initial approximation.

### 1.3 Kamal Transform [11,13,14]

Kamal transform was introduced by Abdelilah Kamal in 2016, for soft growth of the process of solving linear ordinary and partial differential equations in the time domain. It is derived from the traditional Fourier integral. Kamal transform is based on its elementary properties for mathematical straightforwardness same as like Fourier, Laplace , Sumudu , Elzaki , Aboodh and Mahgoub transforms are the expedient mathematical tools for solving differential equations, Kamal transform defined for function of exponential order we consider functions in the set  $A$  defined by:

$$A = \left\{ f(t) : \exists M, m_1, m_2 > 0, |f(t)| < Me^{\frac{|t|}{m_j}}, \text{ if } t \in (-1)^j \times [0, \infty) \right\}$$

For a given function in the set  $A$ , the constant  $M$  must be finite number,  $m_1, m_2$  may be finite or infinite.

Kamal Transform is denoted and defined as,

$$K[f(t)] = T(u) = \int_0^{\infty} e^{-\frac{t}{u}} f(t) dt ; t \geq 0, m_1 < u < m_2$$

The reason of this study is to show the applicability of this attractive new transform and its competence in solving the partial differential equations combined with the method VIM.

#### 1.3.1 Kamal Transform and Inverse Kamal Transform of some Basic Functions

$f(t)$	$K[f(t)] = T(u)$	$K^{-1}[T(u)] = f(t)$
1	$K[1] = u$	$K^{-1}[u] = 1$
$t$	$K[t] = u^2$	$K^{-1}[u^2] = t$
$t^2$	$K[t^2] = 2u^3$	$K^{-1}[u^3] = \frac{t^2}{2!}$
$t^n$	$K[t^n] = n!u^{n+1}$	$K^{-1}[u^n] = \frac{t^{n-1}}{(n-1)!}$
$e^{at}$	$K[e^{at}] = \frac{u}{1-au}$	$K^{-1}\left[\frac{u}{1-au}\right] = e^{at}$
$e^{-at}$	$K[e^{-at}] = \frac{u}{1+au}$	$K^{-1}\left[\frac{u}{1+au}\right] = e^{-at}$
$\sin at$	$K[\sin at] = \frac{au^2}{1+a^2u^2}$	$K^{-1}\left[\frac{u^2}{1+a^2u^2}\right] = \frac{\sin at}{a}$
$\cos at$	$K[\cos at] = \frac{u}{1+a^2u^2}$	$K^{-1}\left[\frac{u}{1+a^2u^2}\right] = \cos at$
$\sinh at$	$K[\sinh at] = \frac{au^2}{1-a^2u^2}$	$K^{-1}\left[\frac{u^2}{1-a^2u^2}\right] = \frac{\sinh at}{a}$
$\cosh at$	$K[\cosh at] = \frac{u}{1-a^2u^2}$	$K^{-1}\left[\frac{u}{1-a^2u^2}\right] = \cosh at$

### 1.3.2 Kamal Transform of Derivatives of the functions

- (1)  $K[f'(t)] = \frac{T(u)}{u} - f(0)$
- (2)  $K[f''(t)] = \frac{T(u)}{u^2} - \frac{f(0)}{u} - f'(0)$
- (3)  $K[f'''(t)] = \frac{T(u)}{u^3} - \frac{f(0)}{u^2} - \frac{f'(0)}{u} - f''(0)$
- (4)  $K[f^{(n)}(t)] = \frac{T(u)}{u^n} - \frac{f(0)}{u^{n-1}} - \frac{f'(0)}{u^{n-2}} - \dots - f^{(n-1)}(0)$

## 2. Methodology

### 2.1 Variational Iteration Method (VIM) [4-7]

In 1978, Inokuti et.al.[1] has proposed a general Lagrange Multiplier method to solve problems arise in quantum mechanics. In 1998, Chinese Mathematician, J.Huan He [4-7], has modified the Lagrange Multiplier Method into an Iteration Method, known as the Variational Iteration Method. The VIM gives successive approximations of the solution that may converge rapidly to the exact solution if such a solution exists. For concrete problems, obtained approximations can be used for numerical purpose.

To illustrate the equation consider,  $L[\chi(x,t)] + N[\chi(x,t)] = f(x,t)$  (1)

Where  $L$  is Linear Operator,  $N$  is a Non Linear Operator and  $f(x,t)$  is given continuous source function. VIM gives the possibility to write the solution of eq. (1) with the support of the correction functional in  $t$ - direction,

$$\chi_{n+1}(x,t) = \chi_n(x,t) + \int_0^t \lambda(x,\tau) \left[ L[\chi_n(x,\tau)] + N[\bar{\chi}_n(x,\tau)] - f(x,\tau) \right] d\tau \quad (2)$$

Here ' $\lambda$ ' is a general Lagrange Multiplier can be identified optimally via variational theory.  $\bar{\chi}_n$  is restricted variation. i.e.  $\delta \bar{\chi}_n = 0$ .

Making the correction functional stationary, we obtained,

$$\delta \chi_{n+1}(x,t) = \delta \chi_n(x,t) + \delta \int_0^t \lambda(x,\tau) \left[ L[\chi_n(x,\tau)] + N[\bar{\chi}_n(x,\tau)] - f(x,\tau) \right] d\tau \quad (3)$$

The solution of the differential equation is considered as the fixed point of the following functional under the suitable choice of the initial term  $\chi_0(x,t)$ . So, the given equation (2) will be reduces to,

$$\chi_{n+1}(x,t) = \chi_n(x,t) + \int_0^t \lambda(x,\tau) \left[ L[\chi_n(x,\tau)] + N[\chi_n(x,\tau)] - f(x,\tau) \right] d\tau \quad (4)$$

Using the selective initial value  $\chi_0(x,t)$  and the obtained Lagrange multiplier  $\lambda$ , the solution  $\chi(x,t)$  will be readily obtained. The solution will be obtained by taking the limit of the successive approximations as,  $\lim_{n \rightarrow \infty} \chi_n(x,t) = \chi(x,t)$ ; where,  $\chi_n(x,t); n \geq 0$  are the  $n$ th consecutive approximations [2-5].

## 2.2 Variational Iteration Adomian Decomposition Method (VIADM) [9]

For the Non Linear term in eq.(4), we can use Adomian Polynomials  $N(u) = \sum_{n=0}^{\infty} A_n$ , where;

$$A_0 = f(u_0)$$

$$A_1 = f'(u_0)u_1$$

$$A_2 = f''(u_0)u_2 + \frac{f''(u_0)}{2!}u_1^2$$

And so on..

So, Iteration formula eq.(4) reduces to;

$$\chi_{n+1}(x,t) = \chi_n(x,t) + \int_0^t \lambda(x,\tau) \left[ L[\chi_n(x,\tau)] + \sum_{n=0}^{\infty} A_n - f(x,\tau) \right] d\tau \quad (5)$$

This is known as Iteration formula for the Variational Iteration Adomian Decomposition Method.

## 2.3 New Modified Kamal Variational Iteration Adomian Decomposition Method (New MKVIADM)

Taking Kamal Transform on both the sides of eq.(5) as the integration is basically the single convolution with respect to 't' and hence Kamal Transform is appropriate to use.

$$\begin{aligned}
 K[\chi_{n+1}(x,t)] &= K[\chi_n(x,t)] + K\left[\int_0^t \lambda(x,\tau) \left[ L[\chi_n(x,\tau)] + \sum_{n=0}^{\infty} A_n - f(x,\tau) \right] d\tau \right] \\
 \therefore K[\chi_{n+1}(x,t)] &= K[\chi_n(x,t)] + K\left\{ [\lambda(x,\tau)] * \left[ L[\chi_n(x,\tau)] + \sum_{n=0}^{\infty} A_n - f(x,\tau) \right] \right\} \\
 \therefore K[\chi_{n+1}(x,t)] &= K[\chi_n(x,t)] + K[\lambda(x,\tau)] \cdot K\left[ L[\chi_n(x,\tau)] + \sum_{n=0}^{\infty} A_n - f(x,\tau) \right] \quad (6)
 \end{aligned}$$

Where \* is the single convolution with respect to 't'. To find the optimal value of  $\lambda = \lambda(x, \tau - t)$ , we first take the variation w.r.t.  $\chi_n(x, t)$ . Thus,

$$\begin{aligned}
 \therefore \frac{\delta}{\delta \chi_n} K[\chi_{n+1}(x,t)] &= \frac{\delta}{\delta \chi_n} K[\chi_n(x,t)] + \frac{\delta}{\delta \chi_n} K[\lambda(x,\tau)] \cdot K\left[ L[\chi_n(x,\tau)] + \sum_{n=0}^{\infty} A_n - f(x,\tau) \right] \\
 \therefore K[\delta \chi_{n+1}(x,t)] &= K[\delta \chi_n(x,t)] + \delta \left\{ K[\lambda(x,t)] \cdot K[L[\chi_n(x,t)]] \right\}
 \end{aligned}$$

Simplifying this equation and finding the  $\lambda$  value, substituting into eq.(6) and solving, we obtain the iteration equations.

### 3. Trial Problems

**Problem 1** Consider the Non Linear Wave Equation:

$$\chi_{tt} - \chi_{xx} + \chi^2 = (x^2 + t^2)^2 ; \chi(x,0) = x^2, \chi_t(x,0) = 0 \quad [15] \quad (7)$$

**Solution: Method 1 VIM**

Using VIM, the correction functional in t- direction will be,

$$\chi_{n+1}(x,t) = \chi_n(x,t) + \int_0^t \lambda(x,\tau) \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \bar{\chi}_n}{\partial x^2} + \bar{\chi}_n^2 - (x^2 + \tau^2)^2 \right] d\tau \quad (8)$$

Making correction functional stationary we get,

$$\delta \chi_{n+1}(x,t) = \delta \chi_n(x,t) + \delta \int_0^t \lambda(x,\tau) \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \bar{\chi}_n}{\partial x^2} + \bar{\chi}_n^2 - (x^2 + \tau^2)^2 \right] d\tau$$

Here ' $\lambda$ ' is a general Lagrange Multiplier that can be identified optimally via variational theory.  $\bar{\chi}_n$  is restricted variation. That is there is no variation in that direction. i.e.  $\delta \bar{\chi}_n = 0$ . Taking integration by parts on both the sides and simplifying the equations, we obtain,  $\lambda = \tau - t$ .

So, from (8), Iteration equation will be,

$$\chi_{n+1}(x,t) = \chi_n(x,t) + \int_0^t (\tau-t) \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \chi_n^2 - (x^2 + \tau^2)^2 \right] d\tau \quad (9)$$

Now, Initial condition is,  $\chi_0(x,t) = \chi(x,0) + t \chi_t(x,0) = x^2$ .

$$\therefore (9) \Rightarrow \chi_1(x,t) = \chi_0(x,t) + \int_0^t (\tau-t) \left[ \frac{\partial^2 \chi_0}{\partial \tau^2} - \frac{\partial^2 \chi_0}{\partial x^2} + \chi_0^2 - (x^2 + \tau^2)^2 \right] d\tau$$

$$\therefore \chi_1(x,t) = x^2 + t^2 + \frac{1}{6} x^2 t^4 + \frac{1}{30} t^6$$

$$\therefore \chi_2(x,t) = x^2 + t^2 + \frac{1}{90} t^6 - \frac{23}{420} x^2 t^8 - \frac{1}{3240} x^4 t^{10} - \frac{1}{11880} x^2 t^{12} - \frac{1}{163800} t^{14}$$

⋮

And so on. Neglecting the noise terms and taking  $\lim_{n \rightarrow \infty} \chi_n(x,t) = \chi(x,t)$ , we get the solution in

closed form as ;  $\chi(x,t) = x^2 + t^2$ .

### Method 02 VIADM

On the iteration eq. (9), using Adomian polynomials for non linear terms, eq.(9) will become;

$$\chi_{n+1}(x,t) = \chi_n(x,t) + \int_0^t (\tau-t) \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \sum_{n=0}^{\infty} A_n - (x^2 + \tau^2)^2 \right] d\tau \quad (10)$$

Adomian polynomials are,

$$A_0 = f(\chi_0) = \chi_0^2 = x^4$$

$$A_1 = f'(\chi_0) \chi_1 = 2\chi_0 \chi_1 = 2x^2 \left( x^2 + t^2 + \frac{1}{6} x^2 t^4 + \frac{1}{30} t^6 \right)$$

$$A_2 = f''(\chi_0) \chi_2 + \frac{f''(\chi_0)}{2!} \chi_1^2 = 2\chi_0 \chi_2 + \chi_1^2$$

$$A_3 = f'(\chi_0) \chi_3 + \frac{f''(\chi_0)}{2!} 2\chi_1 \chi_2 + \frac{f'''(\chi_0)}{3!} \chi_1^3 = 2\chi_0 \chi_3 + 2\chi_1 \chi_2$$

And so on.

$$\therefore (10) \Rightarrow \chi_1(x,t) = \chi_0(x,t) + \int_0^t (\tau-t) \left[ \frac{\partial^2 \chi_0}{\partial \tau^2} - \frac{\partial^2 \chi_0}{\partial x^2} + A_0 - (x^2 + \tau^2)^2 \right] d\tau$$

$$\therefore \chi_1(x,t) = x^2 + t^2 + \frac{1}{6} x^2 t^4 + \frac{1}{30} t^6$$

$$\text{llly, } \chi_2(x,t) = \chi_1(x,t) + \int_0^t (\tau-t) \left[ \frac{\partial^2 \chi_1}{\partial \tau^2} - \frac{\partial^2 \chi_1}{\partial x^2} + A_1 - (x^2 + \tau^2)^2 \right] d\tau$$

$$= x^2 + t^2 + \frac{2}{45} t^6 - \frac{1}{2} x^4 t^2 - \frac{1}{90} x^4 t^6 - \frac{1}{840} x^2 t^8$$

And so on. Neglecting the noise terms and taking  $\lim_{n \rightarrow \infty} \chi_n(x, t) = \chi(x, t)$ , we get the solution in closed form as ;  $\chi(x, t) = x^2 + t^2$ .

### Method 03 New MKVIADM

Applying Kamal Transform on both the sides of (9),

$$K[\chi_{n+1}(x, t)] = K[\chi_n(x, t)] + K \left[ \int_0^t \lambda(x, \tau) \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \chi_n^2 - (x^2 + \tau^2)^2 \right] d\tau \right]$$

Using Convolution theorem,

$$\begin{aligned} \therefore K[\chi_{n+1}(x, t)] &= K[\chi_n(x, t)] + K \left\{ \lambda(x, \tau) * \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \chi_n^2 - (x^2 + \tau^2)^2 \right] \right\} \\ \therefore K[\chi_{n+1}(x, t)] &= K[\chi_n(x, t)] + K[\lambda(x, \tau)] \cdot K \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \chi_n^2 - (x^2 + \tau^2)^2 \right] \end{aligned} \quad (11)$$

Making correction functional eq. (11) stationary,

$$\begin{aligned} \therefore (11) \Rightarrow K\delta\chi_{n+1}(x, t) &= K[\delta\chi_n(x, t)] + K[\lambda(x, \tau)] \\ &\cdot \delta \left[ \left( \frac{1}{u^2} \right) K[\chi_n(x, t)] - \frac{1}{u} \chi_n(x, 0) - (\chi_n)_t(x, 0) - K(\chi_n)_{xx} + K\chi_n^2 - K(x^2 + \tau^2)^2 \right] \\ &= K[\delta\chi_n(x, t)] + K[\lambda(x, \tau)] \cdot \delta \left[ \left( \frac{1}{u^2} \right) K[\chi_n(x, t)] - \frac{1}{u} x^2 - K(\chi_n)_{xx} + K\chi_n^2 - K(x^2 + \tau^2)^2 \right] \\ &= K[\delta\chi_n(x, t)] + K[\lambda(x, \tau)] \cdot \left[ \left( \frac{1}{u^2} \right) K[\delta\chi_n(x, t)] \right] \\ \therefore K[\delta\chi_{n+1}(x, t)] &= K[\delta\chi_n(x, t)] \left\{ 1 + \frac{1}{u^2} K[\lambda(x, t)] \right\} \\ \therefore u^2 + K[\lambda(x, t)] &= 0 \Rightarrow K[\lambda(x, t)] = -u^2 \therefore \lambda(x, t) = -t \end{aligned}$$

$$\therefore (11) \Rightarrow K[\chi_{n+1}(x, t)] = K[\chi_n(x, t)] + K[-t] \cdot K \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \chi_n^2 - (x^2 + \tau^2)^2 \right] \quad (12)$$

Now for non linear term, applying the Adomian polynomials,

$$\therefore (12) \Rightarrow K[\chi_{n+1}(x, t)] = K[\chi_n(x, t)] + K[-t] \cdot K \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \sum_{n=0}^{\infty} A_n - (x^2 + \tau^2)^2 \right] \quad (13)$$

This is the New Modified Kamal Variational Iteration Adomian Decomposition Method.

Now,  $\chi_0(x, t) = \chi(x, 0) + t \chi_t(x, 0) = x^2$

$$K[\chi_1(x,t)] = K[[\chi_0(x,t)]] + K[-t] \cdot K\left[(\chi_0)_{tt} - (\chi_0)_{xx} + A_0 - (x^2 + t^2)^2\right]$$

$$= u x^2 - u^2 \{-2u - 2x^2 2u^3 - 4!u^5\}$$

$$\therefore \chi_1(x,t) = x^2 + t^2 + \frac{1}{6}x^2t^4 + \frac{1}{30}t^6$$

Similarly,  $K[\chi_2(x,t)] = K[[\chi_1(x,t)]] + K[-t] \cdot K\left[(\chi_1)_{tt} - (\chi_1)_{xx} + A_1 - (x^2 + t^2)^2\right]$


$$\therefore \chi_2(x,t) = x^2 + t^2 + \frac{2}{45}t^6 - \frac{1}{2}x^4t^2 - \frac{1}{90}x^4t^6 - \frac{1}{840}x^2t^8$$

And so on. Neglecting the noise terms and taking  $\lim_{n \rightarrow \infty} \chi_n(x,t) = \chi(x,t)$ , we get the solution in closed form as ;  $\chi(x,t) = x^2 + t^2$ .

**Comparison Table** Comparison between Exact Solution, VIM, VIADM & New MKVIADM at t=1.

X	$\chi$ - Exact	$\chi_2$ - VIM	$\chi_2$ - VIADM	$\chi_2$ - MKVIADM	$ \chi_{Exact} - \chi_2_{VIM} $	$ \chi_{Exact} - \chi_2_{VIAADM} $ or $ \chi_{Exact} - \chi_2_{NewMKVIADM} $
0.	1	1.01111	1.04444	1.04444	0.01111	0.04444
0.1	1.01	1.02056	1.05438	1.05438	0.01056	0.04438
0.2	1.04	1.04891	1.08358	1.08358	0.00891	0.04358
0.3	1.09	1.09617	1.1302	1.1302	0.00617	0.0402
0.4	1.16	1.16232	1.19117	1.19117	0.00232	0.03117
0.5	1.25	1.24737	1.2622	1.2622	0.00263	0.0122
0.6	1.36	1.35132	1.33778	1.33778	0.00868	0.02222
0.7	1.49	1.47416	1.41114	1.41114	0.01584	0.07886
0.8	1.64	1.61588	1.47433	1.47433	0.02412	0.16567
0.9	1.81	1.77648	1.51814	1.51814	0.03352	0.29186
1.0	2.0	1.95595	1.53214	1.53214	0.04405	0.46786

**Graphical Representation**



**Figure 1.** Blue Line-Exact, Purple Line-VIM, Green Line-VIADM

**Problem 2** Consider the Non Linear Wave Equation:

$$\chi_{tt} - \chi_{xx} + \chi^2 = 1 + 2xt + x^2t^2 ; \chi(x, 0) = 1, \chi_t(x, 0) = x. \quad [15]$$

**Solution: Method 1 VIM**

Using VIM, the correction functional in  $t$ - direction will be,

$$\chi_{n+1}(x, t) = \chi_n(x, t) + \int_0^t \lambda(x, \tau) \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \bar{\chi}_n}{\partial x^2} + \bar{\chi}_n^2 - 1 - 2x\tau - x^2\tau^2 \right] d\tau \quad (14)$$

Making correction functional stationary we get,

$$\delta \chi_{n+1}(x, t) = \delta \chi_n(x, t) + \delta \int_0^t \lambda(x, \tau) \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \bar{\chi}_n}{\partial x^2} + \bar{\chi}_n^2 - 1 - 2x\tau - x^2\tau^2 \right] d\tau$$

Here ' $\lambda$ ' is a general Lagrange Multiplier that can be identified optimally via variational theory.  $\bar{\chi}_n$  is restricted variation. That is there is no variation in that direction. i.e.  $\delta \bar{\chi}_n = 0$ . Taking integration by parts on both the sides and simplifying the equations, we obtain,  $\lambda = \tau - t$ .

So, from (14), Iteration equation will be,

$$\chi_{n+1}(x, t) = \chi_n(x, t) + \int_0^t (\tau - t) \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \bar{\chi}_n}{\partial x^2} + \bar{\chi}_n^2 - 1 - 2x\tau - x^2\tau^2 \right] d\tau \quad (15)$$

Now, Initial condition is,  $\chi_0(x, t) = \chi(x, 0) + t \chi_t(x, 0) = 1 + xt$ .

$$\therefore (15) \Rightarrow \chi_1(x, t) = \chi_0(x, t) + \int_0^t (\tau - t) \left[ \frac{\partial^2 \chi_0}{\partial \tau^2} - \frac{\partial^2 \bar{\chi}_0}{\partial x^2} + \bar{\chi}_0^2 - 1 - 2x\tau - x^2\tau^2 \right] d\tau$$

$$\therefore \chi_1(x, t) = 1 + xt$$

$$\therefore \chi_2(x, t) = 1 + xt$$

⋮

And so on. Neglecting the noise terms and taking  $\lim_{n \rightarrow \infty} \chi_n(x, t) = \chi(x, t)$ , we get the solution in closed form as ;  $\chi(x, t) = 1 + xt$ .

**Method 02 VIADM**

On the iteration eq. (15), using Adomian polynomials for non linear terms, eq.(15) will become;

$$\chi_{n+1}(x, t) = \chi_n(x, t) + \int_0^t (\tau - t) \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \bar{\chi}_n}{\partial x^2} + \sum_{n=0}^{\infty} A_n - 1 - 2x\tau - x^2\tau^2 \right] d\tau \quad (16)$$

Adomian polynomials are,

$$A_0 = f(\chi_0) = \chi_0^2 = (1 + xt)^2$$

$$A_1 = f'(\chi_0) \chi_1 = 2\chi_0 \chi_1 = 2(1 + xt)^2$$

$$A_2 = 2\chi_0 \chi_2 + \chi_1^2 = 3 + 6xt + 3x^2t^2 - t^2 - \frac{5}{3}xt^3 - \frac{5}{6}x^2t^4 - \frac{1}{6}x^3t^5$$

And so on.

$$\therefore (16) \Rightarrow \chi_1(x, t) = \chi_0(x, t) + \int_0^t (\tau - t) \left[ \frac{\partial^2 \chi_0}{\partial \tau^2} - \frac{\partial^2 \chi_0}{\partial x^2} + A_0 - 1 - 2x\tau - x^2\tau^2 \right] d\tau$$

$$\therefore \chi_1(x, t) = 1 + xt$$

$$\text{llly, } \chi_2(x, t) = \chi_1(x, t) + \int_0^t (\tau - t) \left[ \frac{\partial^2 \chi_1}{\partial \tau^2} - \frac{\partial^2 \chi_1}{\partial x^2} + A_1 - 1 - 2x\tau - x^2\tau^2 \right] d\tau$$

$$= 1 + xt - \frac{t^2}{2} - \frac{1}{3}xt^3 - \frac{1}{12}x^2t^4$$

$$\chi_3(x, t) = \chi_2(x, t) + \int_0^t (\tau - t) \left[ \frac{\partial^2 \chi_2}{\partial \tau^2} - \frac{\partial^2 \chi_2}{\partial x^2} + A_2 - 1 - 2x\tau - x^2\tau^2 \right] d\tau$$

$$= 1 + xt - t^2 + \frac{1}{12}t^4 - \frac{2}{3}xt^3 - \frac{1}{6}x^2t^4 - \frac{1}{8}xt^5 + \frac{1}{36}x^2t^6 + \frac{1}{252}x^3t^7$$

And so on. Neglecting the noise terms and taking  $\lim_{n \rightarrow \infty} \chi_n(x, t) = \chi(x, t)$ , we get the solution in closed form as ;  $\chi(x, t) = 1 + xt$ .

### Method 03 New MKVIADM

Applying Kamal Transform on both the sides of (15),

$$K[\chi_{n+1}(x, t)] = K[\chi_n(x, t)] + K \left[ \int_0^t \lambda(x, \tau) \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \chi_n^2 - 1 - 2x\tau - x^2\tau^2 \right] d\tau \right]$$

Using Convolution theorem,

$$\therefore K[\chi_{n+1}(x, t)] = K[\chi_n(x, t)] + K \left\{ \lambda(x, \tau) * \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \chi_n^2 - 1 - 2x\tau - x^2\tau^2 \right] \right\}$$

$$\therefore K[\chi_{n+1}(x, t)] = K[\chi_n(x, t)] + K[\lambda(x, \tau)] \cdot K \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \chi_n^2 - 1 - 2x\tau - x^2\tau^2 \right] \quad (17)$$

Making correction functional eq. (17) stationary,

$$\therefore (17) \Rightarrow K\delta\chi_{n+1}(x, t) = K[\delta\chi_n(x, t)] + K[\lambda(x, \tau)]$$

$$\cdot \delta \left[ \left( \frac{1}{u^2} \right) K[\chi_n(x, t)] - \frac{1}{u} \chi_n(x, 0) - (\chi_n)_t(x, 0) - K(\chi_n)_{xx} + K\chi_n^2 - K(1 - 2x\tau - x^2\tau^2) \right]$$

$$= K[\delta\chi_n(x,t)] + K[\lambda(x,\tau)] \cdot \delta \left[ \left( \frac{1}{u^2} \right) K[\chi_n(x,t)] - \frac{1}{u} x^2 - K(\chi_n)_{xx} + K\chi_n^2 - K(1 - 2x\tau - x^2\tau^2) \right]$$

$$= K[\delta\chi_n(x,t)] + K[\lambda(x,\tau)] \cdot \left[ \left( \frac{1}{u^2} \right) K[\delta\chi_n(x,t)] \right]$$

$$\therefore K[\delta\chi_{n+1}(x,t)] = K[\delta\chi_n(x,t)] \left\{ 1 + \frac{1}{u^2} K[\lambda(x,t)] \right\}$$

$$\therefore u^2 + K[\lambda(x,t)] = 0 \Rightarrow K[\lambda(x,t)] = -u^2 \therefore \lambda(x,t) = -t$$

$$\therefore (17) \Rightarrow K[\chi_{n+1}(x,t)] = K[\chi_n(x,t)] + K[-t] \cdot K \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \chi_n^2 - 1 - 2x\tau - x^2\tau^2 \right] \quad (18)$$

Now for non linear term, applying the Adomian polynomials,

$$\therefore (18) \Rightarrow K[\chi_{n+1}(x,t)] = K[\chi_n(x,t)] + K[-t] \cdot K \left[ \frac{\partial^2 \chi_n}{\partial \tau^2} - \frac{\partial^2 \chi_n}{\partial x^2} + \sum_{n=0}^{\infty} A_n - 1 - 2x\tau - x^2\tau^2 \right] \quad (19)$$

This is the New Modified Kamal Variational Iteration Adomian Decomposition Method.

$$\text{Now, } \chi_0(x,t) = \chi(x,0) + t \chi_t(x,0) = 1 + xt$$

$$\begin{aligned} K[\chi_1(x,t)] &= K[\chi_0(x,t)] + K[-t] \cdot K \left[ (\chi_0)_{tt} - (\chi_0)_{xx} + A_0 - 1 - 2x\tau - x^2\tau^2 \right] \\ &= K[1 + xt] \end{aligned}$$

$$\therefore \chi_1(x,t) = 1 + xt$$

$$\text{Similarly, } K[\chi_2(x,t)] = K[\chi_1(x,t)] + K[-t] \cdot K \left[ (\chi_1)_{tt} - (\chi_1)_{xx} + A_1 - 1 - 2x\tau - x^2\tau^2 \right]$$

$$\therefore \chi_2(x,t) = 1 + xt - \frac{t^2}{2} - \frac{1}{3}xt^3 - \frac{1}{12}x^2t^4$$

$$K[\chi_3(x,t)] = K[\chi_2(x,t)] + K[-t] \cdot K \left[ (\chi_2)_{tt} - (\chi_2)_{xx} + A_2 - 1 - 2x\tau - x^2\tau^2 \right]$$

$$\therefore \chi_3(x,t) = 1 + xt - t^2 + \frac{1}{12}t^4 - \frac{2}{3}xt^3 - \frac{1}{6}x^2t^4 - \frac{1}{8}xt^5 + \frac{1}{36}x^2t^6 + \frac{1}{252}x^3t^7$$

And so on. Neglecting the noise terms and taking  $\lim_{n \rightarrow \infty} \chi_n(x,t) = \chi(x,t)$ , we get the solution in closed


form as ;  $\chi(x,t) = x^2 + t^2$ .

### Comparison Table

Comparison between Exact Solution, VIM, VIADM & New MKVIADM at t=1.

X	$\chi$ - Exact	$\chi_2$ - VIM	$\chi_2$ - VIADM	$\chi_2$ - MKVIADM	$ \chi_{Exact} - \chi_2_{VIM} $	$ \chi_{Exact} - \chi_2_{VIAADM} $ or $ \chi_{Exact} - \chi_2_{NewMKVIADM} $
0.	1.0	1.0	0.083333	0.083333	0.00	0.916667
0.1	1.1	1.1	0.102782	0.102782	0.00	0.997218
0.2	1.2	1.2	0.119476	0.119476	0.00	1.080524
0.3	1.3	1.3	0.13344	0.13344	0.00	1.16656
0.4	1.4	1.4	0.144698	0.144698	0.00	1.255302
0.5	1.5	1.5	0.153274	0.153274	0.00	1.346726
0.6	1.6	1.6	0.15919	0.15919	0.00	1.44081
0.7	1.7	1.7	0.162472	0.162472	0.00	1.537528
0.8	1.8	1.8	0.163143	0.163143	0.00	1.636857
0.9	1.9	1.9	0.161226	0.161226	0.00	1.738774
1.0	2.0	2.0	0.156746	0.156746	0.00	1.843254

### Graphical Representation



**Figure 2.** Blue Line & Purple Line –same line for Exact & VIM, Green Line-VIADM

**Problem 3** Consider the In Viscid Burgers' Equation:

$$\chi_t + \chi\chi_x = 0; \chi(x,0) = 2x. \quad [15] \quad (20)$$

**Solution: Method 1 VIM**

Using VIM, the correction functional in  $t$ - direction will be,

$$\chi_{n+1}(x,t) = \chi_n(x,t) + \int_0^t \lambda(x,\tau) \left[ \frac{\partial \chi_n}{\partial \tau} + \bar{\chi}_n \frac{\partial \bar{\chi}_n}{\partial x} \right] d\tau \quad (21)$$

Making correction functional stationary we get,

$$\delta \chi_{n+1}(x,t) = \delta \chi_n(x,t) + \delta \int_0^t \lambda(x,\tau) \left[ \frac{\partial \chi_n}{\partial \tau} + \bar{\chi}_n \frac{\partial \bar{\chi}_n}{\partial x} \right] d\tau$$

Here ' $\lambda$ ' is a general Lagrange Multiplier that can be identified optimally via variational theory.  $\bar{\chi}_n$  is restricted variation. That is there is no variation in that direction. i.e.  $\delta\bar{\chi}_n = 0$ . Taking integration by parts on both the sides and simplifying the equations, we obtain,  $\lambda = -1$ .

So, from (21), Iteration equation will be,

$$\chi_{n+1}(x,t) = \chi_n(x,t) - \int_0^t \left[ \frac{\partial \chi_n}{\partial \tau} + \bar{\chi}_n \frac{\partial \bar{\chi}_n}{\partial x} \right] d\tau \quad (22)$$

Now, Initial condition is,  $\chi_0(x,t) = 2x$ .

$$\therefore (22) \Rightarrow \chi_1(x,t) = \chi_0(x,t) - \int_0^t \left[ \frac{\partial \chi_0}{\partial \tau} + \chi_0 \frac{\partial \chi_0}{\partial x} \right] d\tau$$

$$\therefore \chi_1(x,t) = 2x(1-2t)$$

$$\therefore \chi_2(x,t) = 2x \left[ 1 - 2t + (2t)^2 - \frac{1}{3}(2t)^3 \right]$$

And so on. Continuing in this manner, we can say that,  $\chi_n(x,t) \leq 2x \left( \frac{1}{1+2t} \right)$  and taking

$$\lim_{n \rightarrow \infty} \chi_n(x,t) = \chi(x,t), \text{ we get the solution as; } \chi(x,t) = \frac{2x}{1+2t}.$$

### Method 02 VIADM

On the iteration eq. (21), using Adomian polynomials for non linear terms, eq.(21) will become;

$$\chi_{n+1}(x,t) = \chi_n(x,t) - \int_0^t \left[ \frac{\partial \chi_n}{\partial \tau} + \sum_{n=0}^{\infty} A_n \right] d\tau \quad (23)$$

Adomian polynomials are,

$$A_0 = f(\chi_0) = \chi_0(\chi_0)_x = 4x$$

$$A_1 = f'(\chi_0)\chi_1 = 8x - 16xt$$

And so on.

$$\therefore (23) \Rightarrow \chi_1(x,t) = \chi_0(x,t) - \int_0^t \left[ \frac{\partial \chi_0}{\partial \tau} + A_0 \right] d\tau$$

$$\therefore \chi_1(x,t) = 2x - 4xt$$

$$\begin{aligned} \text{llly, } \chi_2(x,t) &= \chi_1(x,t) - \int_0^t \left[ \frac{\partial \chi_1}{\partial \tau} + A_1 \right] d\tau \\ &= 2x - 8xt + 8xt^2 \end{aligned}$$

And so on. Continuing in this manner, we can say that,  $\chi_n(x,t) \leq 2x \left( \frac{1}{1+2t} \right)$  and taking

$$\lim_{n \rightarrow \infty} \chi_n(x,t) = \chi(x,t), \text{ we get the solution as; } \chi(x,t) = \frac{2x}{1+2t}.$$

### **Method 03 New MVIADKTM**

Applying Kamal Transform on both the sides of (18),

$$\begin{aligned} K[\chi_t] + K[\chi\chi_x] &= K[0] \\ \therefore \frac{1}{u} K[\chi(x,t)] - \chi(x,0) + K[\chi\chi_x] &= 0 \\ \therefore \frac{1}{u} K[\chi(x,t)] &= 2x - K[\chi\chi_x] \\ \therefore K[\chi(x,t)] &= 2xu - uK[\chi\chi_x] \\ \therefore \chi(x,t) &= K^{-1}[2xu - uK[\chi\chi_x]] \end{aligned}$$

So, the new correction functional is,

$$\chi_{n+1}(x,t) = 2x - K^{-1} \left[ uK \left[ \chi_n(\chi_n)_x \right] \right] \quad (24)$$

Now, for Non Linear Terms, Adomian polynomials are,

$$\begin{aligned} A_0 &= f(\chi_0) = \chi_0(\chi_0)_x = 4x \\ A_1 &= f'(\chi_0)\chi_1 = 8x - 16xt \end{aligned}$$

And so on.

So, the correction functional for New Modified Variational iteration Adomian Decomposition Kamal Transform will be,

$$\begin{aligned} \chi_{n+1}(x,t) &= 2x - K^{-1} \left[ uK \left[ \sum_{n=0}^{\infty} A_n \right] \right] \\ \chi_1(x,t) &= 2x - K^{-1} \left[ uK [A_0] \right] \\ &= 2x - 4xt \\ \therefore \chi_2(x,t) &= 2x - K^{-1} \left[ uK [A_1] \right] \\ &= 2x - 8xt + 8xt^2 \end{aligned}$$

And so on. Continuing in this manner, we can say that,  $\chi_n(x,t) \leq 2x \left( \frac{1}{1+2t} \right)$  and taking


$$\lim_{n \rightarrow \infty} \chi_n(x,t) = \chi(x,t), \text{ we get the solution as; } \chi(x,t) = \frac{2x}{1+2t}.$$

### **Comparison Table**

Comparison between Exact Solution, VIM, VIADM & New MKVIADM at t=1.

X	$\chi$ -Exact	$\chi_2$ -VIM	$\chi_2$ - VIADM	$\chi_2$ - MKVIADM	$ \chi_{Exact} - \chi_2 VIM $	$ \chi_{Exact} - \chi_2 VIAADM $ or $ \chi_{Exact} - \chi_2 NewMKVIADM $
0.	0.	0.	0.	0.	0.00	0.00
0.1	0.0666667	0.0666667	0.2	0.2	0.00	0.1333333
0.2	0.1333333	0.1333333	0.4	0.4	0.00	0.2666667
0.3	0.2	0.2	0.6	0.6	0.00	0.4
0.4	0.2666667	0.2666667	0.8	0.8	0.00	0.5333333
0.5	0.3333333	0.3333333	1.0	1.0	0.00	0.6666667
0.6	0.4	0.4	1.2	1.2	0.00	0.8
0.7	0.4666667	0.4666667	1.4	1.4	0.00	1.0666667
0.8	0.5333333	0.5333333	1.6	1.6	0.00	1.0666667
0.9	0.6	0.6	1.8	1.8	0.00	1.2
1.0	0.6666667	0.6666667	2.0	2.0	0.00	1.3333333

### Graphical Representation



**Figure 3.** Blue Line & Green Line –same line for Exact & VIM, Purple Line-VIADM

## 4. Conclusion

Our aim in this paper is to check the accuracy of VIM and the combination of two methods VIM and ADM that is VIADM which is greatly fulfilled. We also introduce the new method as New Modified Kamal Variational Iteration Adomian Decomposition Method, which is in addition successfully employed. Introducing New Modified KVIADM is just utilize for getting joy and satisfied for doing somewhat novel. From the Comparison table and Graphical Representation, we can conclude that for the undertaken trial problems, VIM is decidedly accurate than VIADM. Objective of this paper that how rapidly these methods are converges to the exact solution is also fulfilled.

## References

- [1] M. Inokuti H Sekine T Mura 1978 General use of the Lagrange multiplier in nonlinear mathematical physics, in: S Nemat-Nasser (Ed.) Variational Method in the Mechanics of solids Pergamon Press New York pp. 156–162
- [2] G. Adomian 1989 Nonlinear stochastic systems: Theory and applications to Physics Kluwer Academic press
- [3] G. Adomian, 1994 Solving Frontier problem of Physics: The Decomposition Method Kluwer

Academic press

- [4] J.-H. He, 1997 A new approach to nonlinear partial differential equations Communications in Nonlinear Science and Numerical Simulation vol. 2 no 4 pp. 230–235
- [5] J.-H. He 1999 Variational iteration method-a kind of non-linear analytical technique: some examples, International Journal of Non-Linear Mechanics vol. 34 no 4 pp. 699–708
- [6] Ji Huan He Xu-Hong Wu 2007 Variational iteration method: New development and applications An International Journal Computers and Mathematics with Applications Science Direct Elsevier 54 881–894
- [7] Ji Huan He 2007 Variational iteration method-Some recent results and new interpretations An International Journal of Computational and Applied Mathematics Science Direct Elsevier 207 pp.3 – 17.
- [8] Tamer A Abassy Magdy A El-Tawil H El Zoheiry 2007 Toward a modified variational iteration method Journal of Computational and Applied Mathematics 207 pp. 137 – 147
- [9] Syed Tauseef Mohyud-Din Muhammad Aslam Noor and Khalida Inayat Noor 2009 Modified Variational Iteration Method for Solving Sine-Gordon Equations World Applied Sciences Journal 6 (7) 999-1004 ISSN 1818-4952
- [10] S.A. Khuri A Sayfy 2012 A Laplace variational iteration strategy for the solution of differential equations Applied Mathematics Letters 25 pp.2298–2305
- [11] Abdelilah Kamal H Sedeeg 2016 The New Integral Transform Kamal Transform Advances in Theoretical and Applied Mathematics ISSN 0973-4554 Volume 11 Number 4 pp. 451-458
- [12] Alla M Elsheikh and Tarig M Elzaki 2016 Modified Variational Iteration Method for Solving Fourth Order Parabolic PDEs With Variable Coefficients Global Journal of Pure and Applied Mathematics ISSN 0973-1768 Volume 12 Number 2 pp. 1587-1592
- [13] Abdelilah K Hassan Sedeeg and Zahra I Adam Mahamoud 2017 The Use of Kamal Transform for Solving Partial Differential Equations Advances in Theoretical and Applied Mathematics ISSN 0973-4554 Volume 12 Number 1 pp. 7-13
- [14] A S Khan I I Khan M M Janolkar 2018 Solution of Ordinary Differential Equations with Variable Coefficients Using Kamal Transform International Journal of Scientific Research and Review Volume 7 Issue 3 pp 173-178
- [15] Tarig M Elzaki Solution of Nonlinear Partial Differential Equations by New Laplace Variational Iteration Method Chapter-9 Intech Open 73291