

# General improved Kudryashov method for exact solutions of nonlinear evolution equations in mathematical physics

Abd-Allah Hyder<sup>1,2,5</sup>  and M A Barakat<sup>3,4</sup>

<sup>1</sup> King Khalid University, College of Science, Department of Mathematics, P. O. Box 9004, 61413, Abha, Saudi Arabia

<sup>2</sup> Department of Engineering Mathematics and Physics, Faculty of Engineering, Al-Azhar University, Cairo, Egypt

<sup>3</sup> Department of Mathematics, Faculty of Sciences, Al-Azhar University, Assiut, Egypt

<sup>4</sup> Department of Mathematics, College of Al Wajh, University of Tabuk, Saudi Arabia

E-mail: [abahahmed@kku.edu.sa](mailto:abahahmed@kku.edu.sa) and [Barakat14285@yahoo.com](mailto:Barakat14285@yahoo.com)

Received 10 September 2019, revised 11 December 2019

Accepted for publication 23 December 2019

Published 14 February 2020



## Abstract

This paper is devoted to improving the general Kudryashov method by a new and general auxiliary equation. So, a new method is introduced, which we call ‘the general improved Kudryashov method’, to produce exact solutions for nonlinear evolution equations arising in mathematical physics. As application examples, exact traveling wave solutions for the combined Korteweg–de Vries and modified Korteweg–de Vries (KdV–mKdV) equation and the (2+1)-dimensional Zakharov–Kuznetsov equation are obtained. These solutions can be classified as solitary and periodic wave solutions. Some of the obtained solutions are graphically sketched.

Keywords: Kudryashov method, nonlinear evolution equations, exact solutions

(Some figures may appear in colour only in the online journal)

## 1. Introduction

Nonlinear evolution equations play an important role in physics and applied sciences. Searching for exact traveling wave solutions of nonlinear evolution equations in mathematical physics plays a crucial role in miscellaneous fields of science and engineering, such as soliton theory, plasma physics, nonlinear optics, fluid dynamics, biophysics and many others. So, many powerful methods have been introduced, such as the homogeneous balance method [1, 2], the tanh method [3], the tanh-coth method [4], the modified tanh-coth method [5–7], the inverse scattering method [8], Hirota’s bilinear method [9], the sine-cosine method [10], the  $(G'/G)$ -expansion method [11], the  $(G'/G, 1/G)$ -expansion method [12], the Exp-function method [13], F-expansion method [14–16] and so on.

Consider a nonlinear partial differential equation (NPDE) in polynomial form

$$\Omega(\xi, s, u_s, u_\xi, u_{\xi\xi}, u_{\xi\xi\xi}, \dots) = 0, \quad (1.1)$$

where  $(\xi, s) \in \mathbb{R} \times \mathbb{R}_+$  is the independent variable and  $u$  is the dependent variable. Applying the variable traveling transformation

$$u(\xi, s) = u(\varpi), \quad \varpi = \xi - \epsilon s, \quad (1.2)$$

we change (1.1) to a nonlinear ordinary differential equation (NODE)

$$\Theta(\varpi, u, u', u'', u''', \dots) = 0, \quad (1.3)$$

where  $' := \frac{d}{d\varpi}$ . In 2012, Kudryashov [17] proposed his method for finding exact solutions of the NPDE (1.1). He looked for exact solutions taking into account the expression  $u(\varpi) = \sum_{i=0}^x \alpha_i \chi^i$  where  $\chi = \frac{1}{1+e^\varpi}$  which is a solution of the equation  $\frac{d\chi}{d\varpi} = \chi^2 - \chi$ . A modified Kudryashov method

<sup>5</sup> Author to whom any correspondence should be addressed.

was introduced by replacing the natural exponential function  $e^\varpi$  by the general exponential function  $a^\varpi$  in [18–21]. In these research articles, authors found exact solutions of the NPDE (1.1) by using the expansion  $u(\varpi) = \sum_{i=0}^x \alpha_i \chi^i$  where  $\chi = \frac{1}{1 \pm a^\varpi}$  which is a solution of the equation  $\frac{d\chi}{d\varpi} = \ln a(\chi^2 - \chi)$ . Thereafter, some authors [22–25] applied a general Kudryashov method to explore exact solutions of the NPDE (1.1). They used a rational type expansion  $u(\varpi) = \sum_{i=0}^x \alpha_i \chi^i / \sum_{j=0}^y \beta_j \chi^j$  where  $\chi = \frac{1}{1 + Ce^\varpi}$  which is a solution of the equation  $\frac{d\chi}{d\varpi} = \chi^2 - \chi$ . Recently, some authors have discussed the improvement of the Kudryashov

solved by expanding its general solution in the form

$$u(\varpi) = \frac{\sum_{i=0}^x \alpha_i \chi^i(\varpi)}{\sum_{j=0}^y \beta_j \chi^j(\varpi)}, \quad (2.1)$$

where  $\alpha_i, \beta_j (i = 0, 1, \dots, x, j = 0, 1, \dots, y)$  are constants to be determined and  $\chi$  solves the general auxiliary equation

$$\chi'(\varpi) = \sigma \chi^n(\varpi) - \chi(\varpi), \quad 1 < n \in \mathbb{N}, \quad 0 \neq \sigma \in \mathbb{R}. \quad (2.2)$$

Solving equation (2.2), gives a set of general solutions in the form

$$\chi(\varpi) = \begin{cases} \frac{1}{n - \sqrt[n]{\sigma + C \exp[(n-1)\varpi]}}, & n = 2, 4, 6, 8, \dots, \\ \frac{\pm 1}{n - \sqrt[n]{\sigma + C \exp[(n-1)\varpi]}}, & n = 3, 7, 11, 15, \dots, \\ \frac{\pm 1}{n - \sqrt[n]{\sigma + C \exp[(n-1)\varpi]}} + \frac{\pm i}{n - \sqrt[n]{\sigma + C \exp[(n-1)\varpi]}}, & n = 5, 9, 13, 17, \dots \end{cases} \quad (2.3)$$

method. S M Ege [26, 27] first extend and improved the Kudryashov method by new auxiliary equation. Abdus Salam and Habiba [28] improved the general Kudryashov method given in [22] by introducing the auxiliary equation  $\frac{d\chi}{d\varpi} = \sigma \chi^3 - \chi, 0 \neq \sigma \in \mathbb{R}$ . This equation has the general solution  $\chi = \frac{\pm 1}{\sqrt{\sigma + Ce^{2\varpi}}}$ .

In this paper, we improve the general Kudryashov method given in [22] by a general auxiliary equation  $\chi'(\varpi) = \sigma \chi^n(\varpi) - \chi(\varpi), 1 < n \in \mathbb{N}, 0 \neq \sigma \in \mathbb{R}$ . So, we present a very straightforward and effective method called general improved Kudryashov method for exact solutions of nonlinear evolution equations. The merits of the introduced method lie in two directions. Firstly, it leads to both the solitary and periodic solutions. Secondly, the solution strategy, by this method and the computing system Mathematica, is of utter simplicity, and can be easily extended to all types of nonlinear evolution equations. Moreover, If  $n = 2$  and  $n = 3$  our general auxiliary equation reduces to the auxiliary equations used in [17, 22] and [28], respectively. Hence, a novel improved Kudryashov method for finding exact solutions of nonlinear evolution equations in mathematical physics is introduced. As application examples, new sets of exact solutions for the combined KdV–mKdV equation and the (2+1) dimensional Zakharov–Kuznetsov equation are presented.

## 2. Demonstration of the general improved Kudryashov method

Consider the NPDE (1.1) together with the variable traveling transformation (1.2) and the transformed NODE (1.3). For simplicity, we integrate the NODE (1.3), provided that all terms include derivatives, and set the integration constants to be zero. Subsequently, the transformed NODE (1.3) can be

The positive numbers  $x$  and  $y$  can be specified by balancing the linear and nonlinear terms of highest order in equation (1.3). Inserting equations (2.1) and (2.2) into equation (1.3), yields an algebraic equation in  $\chi$  and its powers. Equating the coefficients of the terms that containing the same power for  $\chi$  to zero, gives an algebraic system of equations in  $\alpha_i, \beta_j$  and  $\epsilon$ . With the help of the computer symbolic system *Mathematica*, we can obtain  $\alpha_i, \beta_j$  and  $\epsilon$ . Eventually, by using these values and the solutions (2.3) of equation (2.2), we can produce exact traveling wave solutions of equation (1.1). In fact, equation (2.2) has many solutions unlike the solutions given in equation (2.3). But we only consider the solutions (2.3), because they have interesting and effective properties.

In the following section, our method is applied for  $n = 5$  to find exact traveling wave solutions of the combined KdV–mKdV equation and the (2+1)-dimensional Zakharov–Kuznetsov equation.


## 3. Applications

### 3.1. The combined KdV–MKdV equation

The KdV and mKdV are known as solitary equations which have been extensively researched. The nonlinear terms of the KdV and mKdV equations usually arise simultaneously in many physical problems such as fluid mechanics, quantum field theory and plasma physics. These nonlinear terms construct a nonlinear evolution equation called the combined KdV–mKdV equation

$$u_s + \delta u u_\xi + \rho u^2 u_\xi + u_{\xi\xi\xi} = 0, \quad (\xi, s) \in \mathbb{R} \times \mathbb{R}_+. \quad (3.1)$$

Here, we apply the general improved Kudryashov method to investigate the combined KdV–mKdV equation (3.1). By



**Figure 1.** (a) 3D plot of the solution  $u_1(\xi, s)$ . (b) 2D plot of the solution  $u_1(\xi, s)$  at  $s = 0$ . Here,  $\sigma = 0.0002$ ,  $\rho = C = 1$ ,  $\delta = 0.002$ ,  $\alpha_0 = 0.005$ ,  $\beta_0 = 0.05$  and  $\beta_1 = 0.5$ .

using the transformation (1.2), we can convert (3.1) into a solutions of equation (3.1) as follows  
NODE

$$-\epsilon u' + \delta uu' + \rho u^2 u' + u''' = 0. \quad (3.2)$$

$$u_1(\xi, s) = \frac{\sqrt[4]{\sigma + C \exp[4\phi(\xi, s)]}}{\beta_1 \pm \beta_0 \sqrt[4]{\sigma + C \exp[4\phi(\xi, s)]}} \times \left[ \frac{\alpha_0 + A}{102\beta_1^2 \sqrt{6\rho}} \pm \frac{155\beta_0^3}{(\sigma + C \exp[4\phi(\xi, s)])^{\frac{1}{2}}} + \frac{255\beta_1^3}{(\sigma + C \exp[4\phi(\xi, s)])^{\frac{3}{4}}} + \frac{2448\beta_1^3 \sigma}{(\sigma + C \exp[4\phi(\xi, s)])^{\frac{5}{4}}} \right], \quad (3.6)$$

Taking the homogeneous balance between  $u''$  and  $u^3$ , we get  $x = y + 4$ . Let  $y = 1$ , then  $x = 5$ . Hence, we can put the traveling wave solution of equation (3.1) in the form

$$u(\varpi) = \frac{\alpha_0 + \alpha_1 \chi + \alpha_2 \chi^2 + \alpha_3 \chi^3 + \alpha_4 \chi^4 + \alpha_5 \chi^5}{\beta_0 + \beta_1 \chi}. \quad (3.4)$$

Substituting equations (3.4) and (2.2) with  $n = 5$  into equation (3.3), gives an algebraic equation in  $\chi$  and its powers. Equating the coefficients of the terms that containing the same power for  $\chi$  to zero, yields an algebraic system of equations in  $\alpha_i$ ,  $\beta_j$  ( $i = 0, 1, 2, 3, 4, 5, j = 0, 1$ ) and  $\epsilon$  (see appendix A). Solving the obtained system by *Mathematica*, we have the following sets of solutions.

#### Case A.

$$\begin{cases} \alpha_0 = \alpha_0, & \alpha_1 = \frac{36\alpha_0\beta_0^2\beta_1 - 180\alpha_0^2\beta_0\beta_1\delta + 12\alpha_0^2\beta_1(3\beta_0\delta + 2\alpha_0\rho)}{6(\alpha_0^2\rho + \beta_0^2 + \alpha_0\beta_0\delta) - \alpha_0(3\beta_0\delta + 2\alpha_0\rho)}, \\ \alpha_2 = \pm \frac{155\beta_0^3}{102\beta_1^2\sqrt{6\rho}}, & \alpha_3 = \pm \frac{10\beta_0^2}{\beta_1\sqrt{96\rho}}, & \alpha_4 = 0, & \alpha_5 = \pm \frac{\sqrt{96}\beta_1\sigma}{\sqrt{\rho}}, \\ \beta_0 = \beta_0, & \beta_1 = \beta_1, & \epsilon = \frac{\alpha_0(3\beta_0\delta + 2\alpha_0\rho)}{6\beta_0^2}, \end{cases} \quad (3.5)$$

where  $\alpha_0$ ,  $\beta_0$  and  $\beta_1$  are free constants. Substituting the values (3.5) into (3.4) and using (2.3), yields traveling wave

$$u_2(\xi, s) = \frac{\sqrt[4]{\sigma + C \exp[4\phi(\xi, s)]}}{\beta_1 \pm i\beta_0 \sqrt[4]{\sigma + C \exp[4\phi(\xi, s)]}} \times \left[ \frac{\alpha_0 + A}{102\beta_1^2 \sqrt{6\rho}} \pm i \frac{155\beta_0^3}{(\sigma + C \exp[4\phi(\xi, s)])^{\frac{1}{2}}} \pm i \frac{255\beta_1^3}{(\sigma + C \exp[4\phi(\xi, s)])^{\frac{3}{4}}} \pm i \frac{2448\beta_1^3 \sigma}{(\sigma + C \exp[4\phi(\xi, s)])^{\frac{5}{4}}} \right], \quad (3.7)$$


where

$$A = \frac{36\alpha_0\beta_0^2\beta_1 - 180\alpha_0^2\beta_0\beta_1\delta + 12\alpha_0^2\beta_1(3\beta_0\delta + 2\alpha_0\rho)}{6(\alpha_0^2\rho + \beta_0^2 + \alpha_0\beta_0\delta) - \alpha_0(3\beta_0\delta + 2\alpha_0\rho)}$$

and

$$\phi(\xi, s) = \xi - \left[ \frac{\alpha_0(3\beta_0\delta + 2\alpha_0\rho)}{6\beta_0^2} \right] s.$$

Figure 1 shows the behaviour of the solution  $u_1(\xi, s)$  for  $\sigma = 0.0002$ ,  $\rho = C = 1$ ,  $\delta = 0.002$ ,  $\alpha_0 = 0.005$ ,  $\beta_0 = 0.05$  and  $\beta_1 = 0.5$ .



**Figure 2.** (a) 3D plot of the solution  $u_3(\xi, s)$ . (b) and (c) 2D plots of the solution  $u_3(\xi, s)$  at  $s = 0$  and  $s = 1$ , respectively. Here,  $\rho = 0.0001$ ,  $\sigma = C = 1$ ,  $\delta = 6.7$ ,  $\alpha_4 = 0.02$ ,  $\beta_0 = 1.5$  and  $\beta_1 = 0.002$ .

### Case B.

$$\begin{cases} \alpha_0 = \frac{\beta_0 \sqrt{9\delta^2 - 384\rho} - 3\beta_0\delta}{8\rho}, & \alpha_1 = \frac{\beta_1 \sqrt{3\delta^2 - 128\rho} - 3\beta_1\delta}{8\rho}, \\ \alpha_2 = \frac{\alpha_4(813\beta_0^2\sigma^2 - 8\alpha_4^2\rho)}{120\beta_1^2\sigma^2}, & \alpha_3 = \frac{-306\alpha_4\beta_0}{72\beta_1}, \quad \alpha_4 = \alpha_4, \quad \alpha_5 = 0, \\ \beta_0 = \beta_0, \quad \beta_1 = \beta_1, \quad \epsilon = \frac{\delta(3\delta - \sqrt{9\delta^2 - 384\rho}) - 64\rho}{32\rho}, \end{cases} \quad (3.8)$$

where  $\alpha_4$ ,  $\beta_0$  and  $\beta_1$  are free constants.

Substituting the values (3.8) into (3.4) and using (2.3), yields traveling wave solutions of equation (3.1) as follows

$$\begin{aligned} u_3(\xi, s) = & \frac{\sqrt[4]{\sigma + C \exp[4\phi(\xi, s)]}}{\beta_1 \pm \beta_0 \sqrt[4]{\sigma + C \exp[4\phi(\xi, s)]}} \\ & \times \left[ \frac{\beta_0 \sqrt{9\delta^2 - 384\rho} - 3\beta_0\delta}{8\rho} \right. \\ & \pm i \frac{\beta_1 \sqrt{3\delta^2 - 128\rho} - 3\beta_1\delta}{8\rho(\sigma + C \exp[4\phi(\xi, s)])^{\frac{1}{4}}} \\ & + \frac{\alpha_4(813\beta_0^2\sigma^2 - 8\alpha_4^2\rho)}{120\beta_1^2\sigma^2(\sigma + C \exp[4\phi(\xi, s)])^{\frac{1}{2}}} \\ & \pm i \frac{306\alpha_4\beta_0}{72\beta_1(\sigma + C \exp[4\phi(\xi, s)])^{\frac{3}{4}}} \\ & \left. + \frac{\alpha_4}{(\sigma + C \exp[4\phi(\xi, s)])} \right], \end{aligned} \quad (3.9)$$

$$\begin{aligned} u_4(\xi, s) = & \frac{\sqrt[4]{\sigma + C \exp[4\phi(\xi, s)]}}{\beta_1 \pm i \beta_0 \sqrt[4]{\sigma + C \exp[4\phi(\xi, s)]}} \\ & \times \left[ \frac{\beta_0 \sqrt{9\delta^2 - 384\rho} - 3\beta_0\delta}{8\rho} \right. \\ & \pm i \frac{\beta_1 \sqrt{3\delta^2 - 128\rho} - 3\beta_1\delta}{8\rho(\sigma + C \exp[4\phi(\xi, s)])^{\frac{1}{4}}} \\ & + \frac{\alpha_4(813\beta_0^2\sigma^2 - 8\alpha_4^2\rho)}{120\beta_1^2\sigma^2(\sigma + C \exp[4\phi(\xi, s)])^{\frac{1}{2}}} \\ & \pm i \frac{306\alpha_4\beta_0}{72\beta_1(\sigma + C \exp[4\phi(\xi, s)])^{\frac{3}{4}}} \\ & \left. + \frac{\alpha_4}{(\sigma + C \exp[4\phi(\xi, s)])} \right], \end{aligned} \quad (3.10)$$

where

$$\phi(\xi, s) = \xi - \left[ \frac{\delta(3\delta - \sqrt{9\delta^2 - 384\rho}) - 64\rho}{32\rho} \right] s.$$

Figure 2 shows the behaviour of the solution  $u_3(\xi, s)$  for  $\rho = 0.0001$ ,  $\sigma = C = 1$ ,  $\delta = 6.7$ ,  $\alpha_4 = 0.02$ ,  $\beta_0 = 1.5$  and  $\beta_1 = 0.002$ .

**Case C.**

$$\begin{cases} \alpha_0 = \alpha_0, \alpha_1 = \frac{\beta_1(-3\delta + \sqrt{9\delta^2 - 528\rho})}{8\rho}, \alpha_2 = \alpha_3 = \alpha_4 = \alpha_5 = 0, \\ \beta_0 = -\frac{\alpha_0}{66}(3\delta + \sqrt{9\delta^2 - 528\rho}), \beta_1 = \beta_1, \epsilon = \frac{-3\delta^2 + \delta\sqrt{9\delta^2 - 528\rho} - 88\rho}{32\rho}, \end{cases} \quad (3.11)$$

where  $\alpha_0$  and  $\beta_1$  are free constants.

Substituting the values (3.11) into (3.4) and using (2.3), yields traveling wave solutions of equation (3.1) as follows

$$u_5(\xi, s) = \frac{528\alpha_0\rho(\sigma + C \exp[4\phi(\xi, s)])^{\frac{1}{4}} \pm 66\beta_1(-3\delta + \sqrt{9\delta^2 - 528\rho})}{-8\alpha_0\rho(3\delta + \sqrt{9\delta^2 - 528\rho})(\sigma + C \exp[4\phi(\xi, s)])^{\frac{1}{4}} \pm 528\beta_1\rho}, \quad (3.12)$$

$$u_6(\xi, s) = \frac{528\alpha_0\rho(\sigma + C \exp[4\phi(\xi, s)])^{\frac{1}{4}} \pm 66i\beta_1(-3\delta + \sqrt{9\delta^2 - 528\rho})}{-8\alpha_0\rho(3\delta + \sqrt{9\delta^2 - 528\rho})(\sigma + C \exp[4\phi(\xi, s)])^{\frac{1}{4}} \pm 528i\beta_1\rho}, \quad (3.13)$$

where

$$\phi(\xi, s) = \xi - \left[ \frac{-3\delta^2 + \delta\sqrt{9\delta^2 - 528\rho} - 88\rho}{32\rho} \right] s.$$

Figure 3 shows the behaviour of the solution  $u_5(\xi, s)$  for  $\rho = 0.0001$ ,  $\sigma = C = 1$ ,  $\delta = 2.5$ ,  $\alpha_0 = -0.005$ , and  $\beta_1 = -0.002$ .

### 3.2. The (2+1)-dimensional modified Zakharov–Kuznetsov equation

In [29], Schamel constructed the (2+1)-dimensional modified Zakharov–Kuznetsov equation:

$$u_s + \mu u^2 u_\xi + \nu(u_{\xi\xi\xi} + u_{\xi\xi\xi}) = 0, \quad (\xi, \zeta, s) \in \mathbb{R}^2 \times \mathbb{R}_+, \quad (3.14)$$

which portrays the ion-acoustic waves inside a cold-ion plasma when the behavior of the electrons does not isothermal during their passway of the wave. According to the transformation:

$$u(\xi, \zeta, s) = u(\varpi), \quad \varpi = \xi + \zeta - \epsilon s, \quad (3.15)$$

Equation (3.14) can be converted into a NODE

$$-\epsilon u' + \mu u^2 u' + 2\nu u''' = 0. \quad (3.16)$$

Integrating the NODE (3.16) and putting the constants of integration to be zero, yields

$$-\epsilon u + \frac{\mu}{3} u^3 + 2\nu u'' = 0. \quad (3.17)$$

Balancing  $u^3$  with  $u''$  gives  $x = 5$  and  $y = 1$ . Therefore, we can set the traveling wave solution of equation (3.14) in

the form

$$u(\varpi) = \frac{\alpha_0 + \alpha_1\chi + \alpha_2\chi^2 + \alpha_3\chi^3 + \alpha_4\chi^4 + \alpha_5\chi^5}{\beta_0 + \beta_1\chi}, \quad (3.18)$$

substituting equations (3.18) and (2.2) with  $n = 5$  into equation (3.17), gets an algebraic equation in  $\chi$  and its powers. Equating the coefficients of the terms that containing the same power for  $\chi$  to zero, gives an algebraic system of equations in  $\alpha_i$ ,  $\beta_j$  ( $i = 0, 1, 2, 3, 4, 5$ ,  $j = 0, 1$ ) and  $\epsilon$  (see appendix B). Solving this system by *Mathematica*, we have the following sets of solutions.

**Case I.**


$$\begin{cases} \alpha_0 = 0, \alpha_1 = \alpha_1, \alpha_2 = \alpha_3 = \alpha_4 = 0, \alpha_5 = -2\alpha_1\sigma, \\ \beta_0 = 0, \beta_1 = \pm \frac{\alpha_1 i}{4} \sqrt{\frac{\mu}{\nu}}, \epsilon = -16\nu, \end{cases} \quad (3.19)$$

where  $\alpha_1$  is a nonzero free parameter.

Substituting the values (3.19) into (3.18) and using (2.3), gets traveling wave solution of equation (3.14) as follows

$$u_1(\xi, \zeta, s) = \pm \frac{4i(1 - 2\sigma)\sqrt{\frac{\nu}{\mu}}}{\sigma + C \exp[4(\xi + \zeta + 16\nu s)]}. \quad (3.20)$$

Figure 4 shows the behaviour of the solution  $u_1(\xi, \zeta, s)$  for  $\mu = 0.1$ ,  $\nu = -1$ ,  $\sigma = 0.25$  and  $C = 1$ .



**Figure 3.** (a) 3D plot of the solution  $u_5(\xi, s)$ . (b)–(d) 2D plots of the solution  $u_5(\xi, s)$  at  $s = 0$ ,  $s = 0.5$  and  $s = 1$ , respectively. Here,  $\rho = 0.0001$ ,  $\sigma = C = 1$ ,  $\delta = 2.5$ ,  $\alpha_0 = -0.005$ , and  $\beta_l = -0.002$ .

### Case II.

$$\begin{cases} \alpha_0 = \pm 4i \beta_0 \sqrt{\frac{3\nu}{\mu}}, \alpha_1 = \pm 4i \beta_1 \sqrt{\frac{3\nu}{\mu}}, \alpha_2 = \alpha_3 = 0, \alpha_4 = \pm 8i \beta_0 \sigma \sqrt{\frac{3\nu}{\mu}}, \\ \alpha_5 = \pm 8i \beta_1 \sigma \sqrt{\frac{3\nu}{\mu}}, \beta_0 = \beta_0, \beta_1 = \beta_1, \epsilon = -16\nu, \end{cases} \quad (3.21)$$

where  $\beta_1$  and  $\beta_2$  are free parameters.

Substituting the values (3.21) into (3.18) and using (2.3), gets traveling wave solution of equation (3.14) as follows

$$u_2(\xi, \zeta, s) = \pm 4i \sqrt{\frac{3\nu}{\mu}} \times \left( 1 + \frac{4\sigma}{\sigma + C \exp[4(\xi + \zeta + 16\nu s)]} \right). \quad (3.22)$$

Figure 5 shows the behaviour of the solution  $u_1(\xi, \zeta, s)$  for  $\mu = C = 1$  and  $\nu = \sigma = -1$ .

### Case III.


$$\begin{cases} \alpha_0 = \frac{\beta_1(9\mu^2 - 23\nu)}{\nu\sqrt{2\nu^2 - 1}}, \alpha_1 = 0, \alpha_2 = \pm \frac{3\mu\nu}{5\beta_1\sqrt{2\nu^2 - 1} + 37\nu}, \alpha_3 = 0, \\ \alpha_4 = \pm \frac{3\mu(\nu^2 + \sigma^2)}{2\beta_1\sqrt{2\nu^2 - 1}}, \alpha_5 = \beta_0 = 0, \beta_1 = \beta_1, \epsilon = \frac{-4\mu^2 + \mu(27\mu^2 - 74\nu) - 33\nu}{\sqrt{2\nu^2 - 1}}, \end{cases} \quad (3.23)$$

where  $\beta_1$  is free parameters.

Substituting the values (3.23) into (3.18) and using (2.3), gets traveling wave solution of equation (3.14) as follows

$$u_3(\xi, \zeta, s) = \pm \frac{(\sigma + C \exp[4\psi(\xi, \zeta, s)])^{\frac{1}{4}}}{\beta_1} \left[ \frac{\beta_1(9\mu^2 - 23\nu)}{\nu\sqrt{2\nu^2 - 1}} \pm \frac{3\mu\nu}{(5\beta_1\sqrt{2\nu^2 - 1} + 37\nu)(\sigma + C \exp[4\psi(\xi, \zeta, s)])^{\frac{1}{2}}} \pm \frac{3\mu(\nu^2 + \sigma^2)}{(5\beta_1\sqrt{2\nu^2 - 1})(\sigma + C \exp[4\psi(\xi, \zeta, s)])} \right], \quad (3.24)$$





**Figure 4.** (a)–(c) 3D plots of the solution  $u_1(\xi, \zeta, s)$  at  $s = 0, 0.2$  and  $0.4$ , respectively. Here,  $\mu = 0.1$ ,  $\nu = -1$ ,  $\sigma = 0.25$  and  $C = 1$ .

where

$$\psi(\xi, \zeta, s) = \xi + \zeta - \left( \frac{-4\mu^2 + \mu(27\mu^2 - 74\nu) - 33\nu}{\sqrt{2\nu^2 - 1}} \right) s.$$

Figure 6 shows the behaviour of the solution  $u_3(\xi, \zeta, s)$  for  $\mu = \nu = C = 1$ ,  $\sigma = 2$  and  $\beta_1 = -0.0002$ .

#### 4. Conclusive remarks


This section is devoted to some conclusive remarks.

**Remark 1.** According to the identity  $e^\theta = \cosh \theta + \sinh \theta$ , our traveling wave solutions, obtained by the general improved Kudryashov method, can be handily converted to solitary wave solutions. For example, the solution  $u_1(\xi, s)$  of the combined KdV–mKdV equation can be converted to a

solitary wave solution as follows

$$\begin{aligned} u_1^*(\xi, s) &= \frac{\sqrt[4]{\sigma + C(\cosh[4\phi(\xi, s)] + \sinh[4\phi(\xi, s)])}}{\beta_1 \pm \beta_0 \sqrt[4]{\sigma + C(\cosh[4\phi(\xi, s)] + \sinh[4\phi(\xi, s)])}} \\ &\times \left[ \frac{\alpha_0 + A}{102\beta_1^2 \sqrt[4]{6\rho}} \right. \\ &\pm \frac{155\beta_0^3}{(\sigma + C(\cosh[4\phi(\xi, s)] + \sinh[4\phi(\xi, s)])^{\frac{1}{2}}} \\ &+ \frac{255\beta_1^3}{(\sigma + C(\cosh[4\phi(\xi, s)] + \sinh[4\phi(\xi, s)])^{\frac{3}{4}}} \\ &\left. + \frac{2448\beta_1^3 \sigma}{(\sigma + C(\cosh[4\phi(\xi, s)] + \sinh[4\phi(\xi, s)])^{\frac{5}{4}}} \right], \end{aligned} \quad (4.1)$$

where  $A$  and  $\phi(\xi, s)$  are given in Case A. Also, the solution  $u_1(\xi, \zeta, s)$  of the (2+1)-dimensional modified Zakharov–Kuznetsov equation can be converted to a solitary wave



**Figure 5.** (a)–(c) 3D plots of the solution  $u_2(\xi, \zeta, s)$  at  $s = 0, 0.2$  and  $0.4$ , respectively. Here,  $\mu = C = 1$  and  $\nu = \sigma = -1$ .

solution as follows

$$u_1^*(\xi, \zeta, s) = \pm \frac{4i(1-2\sigma)\sqrt{\frac{\nu}{\mu}}}{\sigma + C(\cosh[4(\xi + \zeta + 16\nu s)] + \sinh[4(\xi + \zeta + 16\nu s)])}. \quad (4.2)$$

**Remark 2.** Using the identity  $e^{i\theta} = \cos \theta + i \sin \theta$ , our traveling wave solutions, obtained by the general improved Kudryashov method, can be easily converted to periodic wave solutions. For example, the solution  $u_1(\xi, s)$  of the combined KdV–mKdV equation can be converted to a periodic wave solution as follows

where  $A$  and  $\phi(\xi, s)$  are given in Case A. Also, the solution  $u_1(\xi, \zeta, s)$  of the (2+1)-dimensional modified Zakharov–Kuznetsov equation can be converted to a periodic wave solution as follows


$$u_1^{**}(\xi, \zeta, s) = \pm \frac{4i(1-2\sigma)\sqrt{\frac{\nu}{\mu}}}{\sigma + C(\cos[4i(\xi + \zeta + 16\nu s)] - i \sin[4i(\xi + \zeta + 16\nu s)])}. \quad (4.4)$$

**Remark 3.** This paper improved the general Kudryashov method [22] by a general auxiliary equation

$$\chi'(\varpi) = \sigma \chi^n(\varpi) - \chi(\varpi), \quad 1 < n \in \mathbb{N}, \quad 0 \neq \sigma \in \mathbb{R}, \quad (4.5)$$

$$u_1^{**}(\xi, s) = \frac{\sqrt[4]{\sigma + C(\cos[4i\phi(\xi, s)] - i \sin[4i\phi(\xi, s)])}}{\beta_1 \pm \beta_0 \sqrt[4]{\sigma + C(\cos[4i\phi(\xi, s)] - i \sin[4i\phi(\xi, s)])}} \left[ \frac{\alpha_0 + A}{102\beta_1^2 \sqrt[4]{6\rho}} \right. \\ \left. \pm \frac{155\beta_0^3}{(\sigma + C(\cos[4i\phi(\xi, s)] - i \sin[4i\phi(\xi, s)])^{\frac{1}{2}}} + \frac{255\beta_1^3}{(\sigma + C(\cos[4i\phi(\xi, s)] - i \sin[4i\phi(\xi, s)])^{\frac{3}{4}}} \right. \\ \left. + \frac{2448\beta_1^3 \sigma}{(\sigma + C(\cos[4i\phi(\xi, s)] - i \sin[4i\phi(\xi, s)])^{\frac{5}{4}}} \right], \quad (4.3)$$





**Figure 6.** (a)–(c) 3D plots of the solution  $u_3(\xi, \zeta, s)$  at  $s = 0, 0.5$  and  $1$ , respectively. Here,  $\mu = \nu = C = 1$ ,  $\sigma = 2$  and  $\beta_1 = -0.0002$ .

which has numerous general solutions depend on the natural number  $n$ , see (2.3). If  $n = 2$ , equation (4.5) reduces to the auxiliary equation  $\chi'(\varpi) = \sigma\chi^2(\varpi) - \chi(\varpi)$  which used in [17, 22] together with the expansion (2.1) to produce a families of exact solutions for some nonlinear evolution equations. Also, for  $n = 3$ , equation (4.5) reduces to the auxiliary equation  $\chi'(\varpi) = \sigma\chi^3(\varpi) - \chi(\varpi)$  which used in [28] together with the expansion (2.1) to produce a family of exact solutions for some nonlinear evolution equations. In our work, we just used equation (4.5) for  $n = 5$ . But one can use it for many values of the natural number  $n$  and obtain new pairwise disjoint sets of exact solutions. Moreover, we apply our method for the combined KdV–mKdV equation and the (2+1)-dimensional Zakharov–Kuznetsov equation as application examples. But, in fact, one can use it to solve many nonlinear evolution equations arising in mathematical physics, such as Hirota–Satsuma coupled KdV, Sawada–Kotera, Zhiber–Shabat and KdV–Burgers equations.

## 5. Conclusion

In this paper, the general Kudryashov method is improved by a novel auxiliary equation. So, a new method to constructing exact solutions for nonlinear evolution equations is introduced. This method is called the general improved Kudryashov method. The main merit of the general improved Kudryashov method over the

others lies in the fact that it uses a particularly straightforward and effective algorithm to obtain exact solutions for a large number of nonlinear evolution equations. Also, a great variety of exact solution can be derived easily on choosing the parameters that appeared. We apply this method to explore exact traveling wave solution for the combined KdV–mKdV equation and the (2+1)-dimensional Zakharov–Kuznetsov equation. These solutions include solitary and periodic wave solutions. Also, some of the obtained solutions are graphically sketched.

Besides that, our approach generalizes some previous approaches. It is based on improving the general Kudryashov method [22] by the general auxiliary equation (2.2) which has various general solutions depend on the natural number  $n$ . If  $n = 2$  and 3 our approach reduces to the approaches presented in [17, 22] and [28], respectively. In our work, we just used our general auxiliary equation for  $n = 5$ . But one can use it for many values of the natural number  $n$  and obtain new pairwise disjoint sets of exact solutions. In our point of view, there is no any demerit in applying our approach to finding exact solutions for nonlinear evolution equations.

## Acknowledgments

The authors would like to extend their appreciations to the Deanship of Scientific Research at King Khalid University for

funding their work through research groups program under grant number (R. G. P. 1/ 128/40).

## Appendix A

The system of algebraic equations in  $\alpha_i, \beta_j$  ( $i = 0, 1, 2, 3, 4, 5, j = 0, 1$ ) for the combined KdV–mKdV equation

$$\begin{aligned}\chi^0: & -2\alpha_1\beta_0^2 + \alpha_0(2\beta_0\beta_1 + \alpha_0\beta_0\delta - 2\beta_0^2\epsilon + \alpha_0^2\rho) = 0, \\ \chi^1: & -8\alpha_2\beta_0^2 + \alpha_1(2\beta_0\beta_1 + 2\alpha_0\beta_0\delta - 2\beta_0^2\epsilon + 3\alpha_0^2\rho) \\ & + \alpha_0\beta_1(2\beta_1 + \alpha_0\beta_1\delta + 4\beta_0\epsilon) = 0, \\ \chi^2: & \alpha_0(3\alpha_1^2\rho - 2\beta_1^2\epsilon) + \alpha_1(\alpha_1\beta_0\delta + 2\alpha_0\beta_1\delta + 4\beta_0\beta_1\epsilon) \\ & + \alpha_2(2\alpha_0\beta_0\delta - 6\beta_0\beta_1 - 2\beta_0^2\epsilon + 3\alpha_0^2\rho) - 18\alpha_3\beta_0^2 = 0, \\ \chi^3: & -32\alpha_4\beta_0^2 + \alpha_3(2\alpha_0\beta_0\delta - 22\beta_0\beta_1 - 2\beta_0^2\epsilon + 3\alpha_0^2\rho) \\ & + \alpha_2(2\beta_1^2 + 2\alpha_1\beta_0\delta + 2\alpha_0\beta_1\delta - 4\beta_0\beta_1\epsilon \\ & + 6\alpha_0\alpha_1\rho) + \alpha_1(\alpha_1\beta_1\delta - 2\beta_1^2\epsilon + \alpha_1^2\rho) = 0, \\ \chi^4: & -50\alpha_5\beta_0^2 + \alpha_4(46\beta_0\beta_1 + 2\alpha_0\beta_0\delta - 2\beta_0^2\epsilon \\ & + 3\alpha_0^2\rho) + \alpha_3(2\alpha_1\beta_0\delta - 8\beta_1^2 + 2\alpha_0\beta_1\delta + 4\beta_0\beta_1\epsilon \\ & + 6\alpha_0\alpha_1\rho) + \alpha_2(\alpha_2\beta_0\delta + 2\alpha_1\beta_1\delta - 2\beta_1^2\epsilon \\ & + 3\alpha_0\alpha_2\rho + 3\alpha_1^2\rho) + 10\beta_0\sigma(\alpha_1\beta_0 - \alpha_0\beta_1) = 0, \\ \chi^5: & \alpha_5(2\alpha_0\beta_0\delta - 78\beta_0\beta_1 - 2\beta_0^2\epsilon + 3\alpha_0^2\rho) \\ & + \alpha_4(2\alpha_1\beta_0\delta - 18\beta_1^2 + 2\alpha_0\beta_1\delta + 4\beta_0\beta_1\epsilon + 6\alpha_0\alpha_1\rho) \\ & + \alpha_3(2\alpha_2\beta_0\delta + 2\alpha_1\beta_1\delta + 3\alpha_1^2\rho - 2\beta_1^2\epsilon + 6\alpha_0\alpha_2\rho) \\ & + \alpha_2(\alpha_2\beta_1\delta + 3\alpha_1\alpha_2\rho + 24\beta_0^2\sigma) \\ & + 6\beta_1\sigma(\alpha_1\beta_0 - \alpha_0\beta_1) = 0, \\ \chi^6: & \alpha_5(-32\beta_1^2 + 2\alpha_1\beta_0\delta - 4\beta_0\beta_1\delta + 6\alpha_0\alpha_1\rho + 2\alpha_0\beta_1\delta) \\ & + \alpha_4(2\alpha_2\beta_0\delta + 2\alpha_1\beta_1\delta - 2\beta_1^2\epsilon + 3\alpha_1^2\rho \\ & + 6\alpha_0\alpha_2\rho) + \alpha_3(\alpha_3\beta_0\delta + 2\alpha_2\beta_1\delta + 6\alpha_1\alpha_2\rho \\ & + 3\alpha_0\alpha_3\rho + 42\beta_0^2\sigma) + \alpha_2(30\beta_0\beta_1\sigma + \alpha_2^2\rho) = 0, \\ \chi^7: & \alpha_5(2\alpha_2\beta_0\delta + 2\alpha_1\beta_1\delta - 2\beta_1^2\epsilon + 3\alpha_1^2\rho + 6\alpha_0\alpha_2\rho) \\ & + \alpha_4(2\alpha_3\beta_0\delta + 2\alpha_2\beta_1\delta + 6\alpha_1\alpha_2\rho + 6\alpha_0\alpha_3\rho \\ & + 64\beta_0^2\sigma) + \alpha_3(\alpha_3\beta_1\delta + 3\alpha_2^2\rho \\ & + 3\alpha_1\alpha_3\rho + 62\beta_0\beta_1\sigma) + 10\alpha_2\beta_1^2\sigma = 0, \\ \chi^8: & \alpha_5(2\alpha_3\beta_0\delta + 2\alpha_2\beta_1\delta + 6\alpha_1\alpha_2\rho + 6\alpha_0\alpha_3\rho + 90\beta_0^2\sigma) \\ & + \alpha_4(\alpha_4\beta_0\delta + 2\alpha_3\beta_1\delta + 3\alpha_2^2\rho + 3\alpha_0\alpha_4\rho \\ & + 6\alpha_1\alpha_3\rho + 102\beta_0\beta_1\sigma) + \alpha_3(3\alpha_2\alpha_3\rho + 24\beta_1^2\sigma) = 0, \\ \chi^9: & \alpha_5(2\alpha_4\beta_0\delta + 2\alpha_3\beta_1\delta + 3\alpha_2^2\rho + 6\alpha_0\alpha_4\rho + 6\alpha_1\alpha_3\rho \\ & + 150\beta_0\beta_1\sigma) + \alpha_4(\alpha_4\beta_1\delta + 6\alpha_2\alpha_3\rho \\ & + 3\alpha_1\alpha_4\rho + 42\beta_1^2\sigma) + \alpha_3^3\rho = 0, \\ \chi^{10}: & \alpha_5(\alpha_5\beta_0\delta + 2\alpha_4\beta_1\delta + 6\alpha_2\alpha_3\rho + 6\alpha_1\alpha_4\rho \\ & + 3\alpha_0\alpha_5\rho + 64\beta_1^2\sigma) + 3\alpha_4\rho(\alpha_3^3 + \alpha_2\alpha_4) = 0, \\ \chi^{11}: & \alpha_5(\alpha_5\beta_1\delta + 3\alpha_3^3\rho + 6\alpha_2\alpha_4\rho + 3\alpha_1\alpha_5\rho) + 3\alpha_3\alpha_4^2\rho = 0, \\ \chi^{12}: & \alpha_5(6\alpha_3\alpha_4\rho + 3\alpha_2\alpha_5\rho) + \alpha_4^3\rho = 0, \\ \chi^{13}: & 3\alpha_5\rho(\alpha_4^2 + \alpha_3\alpha_5) = 0, \\ \chi^{14}: & 3\alpha_4\alpha_5^2\rho = 0, \\ \chi^{15}: & \alpha_5^3\rho = 0.\end{aligned}$$

## Appendix B

The system of algebraic equations in  $\alpha_i, \beta_j$  ( $i = 0, 1, 2, 3, 4, 5, j = 0, 1$ ) for the (2+1)-dimensional Zakharov–Kuznetsov equation

$$\begin{aligned}\chi^0: & -3\alpha_0\beta_0^2\epsilon + \alpha_0^3\mu = 0, \\ \chi^1: & \alpha_1(3\alpha_0^2\mu - 3\beta_0^2\epsilon + 6\beta_0^2\nu) - 6\alpha_0\beta_0\beta_1(\epsilon + \nu) = 0, \\ \chi^2: & \alpha_2(24\beta_0^2\nu - 3\beta_0^2\epsilon + 3\alpha_0^2\mu) + \alpha_1(6\beta_0\beta_1\nu - 6\beta_0\beta_1\epsilon \\ & + 3\alpha_0\alpha_1\mu) + 3\alpha_0\beta_1^2(6\nu - \epsilon) = 0, \\ \chi^3: & \alpha_3(3\alpha_0^2\mu + 54\beta_0^2\nu - 3\beta_0^2\epsilon) + \alpha_2(18\beta_0\beta_1\nu \\ & + 6\alpha_0\alpha_1\mu - 6\beta_0\beta_1\epsilon) + \alpha_1(\alpha_1^2\mu - 3\beta_1^2\epsilon) = 0, \\ \chi^4: & \alpha_4(3\alpha_0^2\mu - 3\beta_0^2\epsilon + 96\beta_0^2\nu) + \alpha_2(6\alpha_0\alpha_1\mu \\ & - 6\beta_0\beta_1\epsilon + 66\beta_0\beta_1\nu) + \alpha_2(3\alpha_1^2\mu - 3\beta_1^2\epsilon \\ & + 3\alpha_0\alpha_2\mu + 6\beta_1^2\nu) = 0, \\ \chi^5: & \alpha_5(3\alpha_0^2\mu - 3\beta_0^2\epsilon + 150\beta_0\nu) + \alpha_4(138\beta_0\beta_1\nu \\ & + 6\alpha_0\alpha_1\mu - 6\beta_0\beta_1\epsilon) + \alpha_3(3\alpha_1^2\mu - 3\beta_1^2\epsilon \\ & + 6\alpha_0\alpha_2\mu + 24\beta_1^2\nu) + \alpha_1(3\alpha_2^2\mu - 36\beta_0^2\nu\sigma) \\ & + 36\alpha_0\beta_0\beta_1\nu\sigma = 0, \\ \chi^6: & \alpha_5(234\beta_0\beta_1\nu - 6\beta_0\beta_1\epsilon + 6\alpha_0\alpha_1\mu) \\ & + \alpha_4(54\beta_1^2\nu - 3\beta_1^2\epsilon + 6\alpha_0\alpha_2\mu + 3\alpha_1^2\mu) \\ & + \alpha_3(6\alpha_1\alpha_2\mu + 3\alpha_0\alpha_3\mu) + \alpha_2(\alpha_2\mu + 96\beta_0^2\nu\sigma) \\ & + 12\beta_1\nu\sigma(\alpha_0\beta_1 - \alpha_1\beta_0) = 0, \\ \chi^7: & \alpha_2(6\alpha_0\alpha_2\mu + 3\alpha_1^2\mu - 3\beta_1^2\epsilon + 96\beta_1^2\nu) \\ & + 6\alpha_4\mu(\alpha_0\alpha_3 + \alpha_1\alpha_2) + \alpha_3(3\alpha_2^2\mu + 3\alpha_1\alpha_3\mu \\ & - 180\beta_0^2\nu\sigma) - 108\alpha_2\beta_0\beta_1\nu\sigma = 0, \\ \chi^8: & 6\alpha_5\mu(\alpha_0\alpha_3 + \alpha_1\alpha_2) + \alpha_4(3\alpha_2^2\mu + 6\alpha_1\alpha_3\mu \\ & + 3\alpha_0\alpha_4\mu - 288\beta_0^2\nu\sigma) + \alpha_3(3\alpha_2\alpha_3\mu - 252\beta_0\beta_1\nu\sigma) \\ & - 36\alpha_2\beta_1^2\nu\sigma = 0, \\ \chi^9: & \alpha_5(6\alpha_0\alpha_4\mu - 420\beta_0^2\nu\sigma + 3\alpha_2^2\mu + 6\alpha_1\alpha_3\mu) \\ & + \alpha_4(6\alpha_2\alpha_3\mu + 3\alpha_1\alpha_4\mu - 444\beta_0\beta_1\nu\sigma) + \alpha_3(\alpha_3\mu \\ & - 96\beta_1^2\nu\sigma) + 30\beta_0\nu\sigma^2(\alpha_1\beta_0 - \alpha_0\beta_1) = 0, \\ \chi^{10}: & \alpha_5(6\alpha_2\alpha_3\mu + 3\alpha_0\alpha_5\mu - 684\beta_0\beta_1\nu\sigma + 6\alpha_1\alpha_4\mu) \\ & + \alpha_4(3\alpha_3^2\mu + 3\alpha_2\alpha_4\mu + 180\beta_1^2\nu\sigma) \\ & + \nu\sigma^2(72\alpha_2\beta_0^2 + 18\alpha_1\beta_0\beta_1 - 18\alpha_0\beta_1^2) = 0, \\ \chi^{11}: & \alpha_5(3\alpha_1\alpha_5\mu - 288\beta_1^2\nu\sigma + 3\alpha_3^2\mu + 6\alpha_2\alpha_4\mu) \\ & + \alpha_3(3\alpha_4^2\mu + 126\beta_0^2\nu\sigma^2) + 90\alpha_2\beta_0\beta_1\nu\sigma^2 = 0, \\ \chi^{12}: & \alpha_5(6\alpha_3\alpha_4\mu + 3\alpha_2\alpha_5\mu) + \alpha_4(192\beta_0^2\nu\sigma^2 + \alpha_4^2\mu) \\ & + 3\beta_1\nu\sigma^2(62\alpha_3\beta_0 + 10\alpha_2\beta_1) = 0, \\ \chi^{13}: & 3\alpha_5(\alpha_4^2\mu + \alpha_3\alpha_5\mu + 90\beta_0^2\nu\sigma^2) \\ & + 3\beta_1\nu\sigma^2(102\alpha_4\beta_0 + 24\alpha_3\beta_1) = 0, \\ \chi^{14}: & 3\alpha_5(\alpha_4\mu + 150\beta_0\beta_1\nu\sigma^2) + 126\alpha_4\beta_1^2\nu\sigma^2 = 0, \\ \chi^{15}: & \alpha_5(\alpha_5^2 + 192\beta_1^2\nu\sigma^2) = 0.\end{aligned}$$

## ORCID iDs

Abd-Allah Hyder  <https://orcid.org/0000-0001-9273-9512>

## References

- [1] Ghany H A and Hyder A 2012 White noise functional solutions for the Wick-type two-dimensional stochastic Zakharov–Kuznetsov equations *Int. Rev. Phys.* **6** 153–7
- [2] Wang M L 1995 Solitary wave solutions for variant Boussinesq equations *Phys. Lett. A* **199** 169–72
- [3] Wazwaz A M 2004 The tanh method for travelling wave solutions of nonlinear equations *Appl. Math. Comput.* **154** 713–23
- [4] Wazwaz A M 2007 The tanh-coth method for solitons and kink solutions for nonlinear parabolic equations *Appl. Math. Comput.* **188** 1467–75
- [5] El-Wakil S A, El-Labany S K, Zahran M A and Sabry R 2005 Modified extended tanh- function method and its applications to nonlinear equations *Appl. Math. Comput.* **161** 403–12
- [6] Ghany H A, Okb El Babb A S, Zabel A M and Hyder A 2013 The fractional coupled KdV equations: exact solutions and white noise functional approach *Chin. Phys. B* **22** 0805011
- [7] Ghany H A and Hyder A 2014 Abundant solutions of Wick-type stochastic fractional 2D KdV equations *Chin. Phys. B* **23** 0605031
- [8] Liu X Q, Jiang S, Fan W B and Liu W M 2004 Soliton solutions in linear magnetic field and time-dependent laser field *Commun. Nonlinear Sci. Numer. Simul.* **9** 361–5
- [9] Hirota R 1971 Exact solution of Korteweg–de Vries equation for multiple collisions of solitons *Phys. Rev. Lett.* **27** 1192–4
- [10] Wazwaz A M 2004 A sine-cosine method for handling nonlinear wave equations *Math. Comput. Modelling* **40** 499–508
- [11] Bekir A 2008 Application of the  $(G'/G)$ -expansion method for nonlinear evolution equations *Phys. Lett. A* **372** 3400–6
- [12] Uddin M H, Akbar M A, Khan Md A and Haque Md Abdul 2017 Close form solutions of the fractional generalized reaction duffing model and the density dependent fractional diffusion reaction equation *Appl. Comput. Math.* **6** 177–84
- [13] He J H and Wu X H 2006 Exp-function method for nonlinear wave equations *Chaos Solitons Fractals* **30** 700–8
- [14] Ghany H A, Hyder A and Zakarya M 2017 Non-Gaussian white noise functional solutions of  $\chi$ -Wick-type stochastic KdV equations *Appl. Math. Inf. Sci.* **11** 915–24
- [15] Hyder A 2017 Wick-type stochastic KdV equation based on Lévy white noise *J. Math. Sci.: Adv. Appl.* **45** 1–20
- [16] Zhao Y M 2013 F-expansion method and its application for finding new exact solutions to the Kudryashov–Sinelschchikov equation *J. Appl. Math.* **2013** 895760
- [17] Kudryashov N A 2012 One method for finding exact solutions of nonlinear differential equations *Commun. Nonlinear Sci. Numer. Simul.* **17** 2248–53
- [18] Ege S M and Misirli E 2014 The modified Kudryashov method for solving some fractional-order nonlinear equations *Adv. Difference Equ.* **2014** 135
- [19] Kilicman A and Silambarasan R 2018 Modified Kudryashov method to solve generalized Kuramoto–Sivashinsky equation *Symmetry* **10** 527
- [20] Kumar D, Seadawy A R and Joardar A K 2018 Modified Kudryashov method via new exact solutions for some conformable fractional differential equations arising in mathematical biology *Chin. J. Phys.* **56** 75–85
- [21] Zayed E M E and Alurfi K A E 2015 The modified Kudryashov method for solving some seventh order nonlinear PDEs in mathematical physics *World J. Modelling Simul.* **11** 308–19
- [22] Islam Md S, Khan K and Arnous A H 2015 Generalized Kudryashov method for solving some  $(3+1)$ -dimensional nonlinear evolution equations *New Trends Math. Sci.* **3** 46–57
- [23] Islam N, Khan K and Islam Md H 2019 Travelling wave solution of Dodd–Bullough–Mikhailov equation: a comparative study between Generalized Kudryashov and improved F-expansion methods *J. Phys. Commun.* **3** 055004
- [24] Mahmud F, Samsuzzoha Md and Akbar M A 2017 The generalized Kudryashov method to obtain exact traveling wave solutions of the PHI-four equation and the Fisher equation *Results Phys.* **7** 4296–302
- [25] Rahman M M, Habib M A, Ali H M S and Miah M M 2019 The generalized Kudryashov method: a renewed mechanism for performing exact solitary wave solutions of some NLEEs *J. Mech. Cont. Math. Sci.* **14** 323–39
- [26] Ege S M 2016 Extended traveling wave solutions for some nonlinear equations *Adv. Math. Sci. J.* **5** 179–89
- [27] Ege S M 2018 Extended traveling wave solutions for some integro partial differential equations *Int. J. Maps Math.* **1** 249–57
- [28] Salam Md A and Habiba U 2019 Application of the improved Kudryashov method to solve the fractional nonlinear partial differential equations *J. Appl. Math. Phys.* **7** 912–20
- [29] Schamel H 1973 A modified Korteweg–de Vries equation for ion acoustic waves due to resonant electrons *J. Plasma Phys.* **9** 377–87