

Corrigendum

Corrigendum: Inspiral into Gargantua (2016 *Class. Quantum Grav.* **33** 155002)

Samuel E Gralla¹ , Scott A Hughes^{2,3} and
Niels Warburton⁴

¹ Department of Physics, University of Arizona, Tucson, AZ 85721, United States of America

² Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, United States of America

³ MIT Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, United States of America

⁴ School of Mathematical Sciences and Complex & Adaptive Systems Laboratory, University College Dublin, Belfield, Dublin 4, Ireland

Received 30 January 2020, revised 17 February 2020

Accepted for publication 25 February 2020

Published 16 April 2020



CrossMark

Our analysis of the change in black hole spin during near-horizon inspiral (section 2.3) was incorrect due to a computational error. The affected equations are (23)–(25), which we repeat here:

$$\frac{d\epsilon}{dt} = \frac{0.102}{M} \frac{\mu^2}{M^2} x_0 \quad (23\text{--old})$$

$$\Delta\epsilon_{\text{total}} < 0.046 \frac{\mu}{M} (X_0 - 2^{1/3} \epsilon^{2/3}) \quad (24\text{--old})$$

$$\Delta \left[\frac{a}{M} \right]_{\text{total}} < 10^{-3} X_0^2 \frac{\mu^2}{M^2}. \quad (25\text{--old})$$

It is more convenient to present the corrected results in terms of $\epsilon^2 = 1 - (a/M)^2 \approx 2(1 - a/M)$, where $\epsilon \ll 1$. Using the same method described in the text and fixing the computational error, we instead find

$$\frac{d\epsilon^2}{dt} = \frac{0.399}{M} \frac{\mu^2}{M^2} x_0^2 \quad (23\text{--new})$$

$$\Delta[\epsilon^2]_{\text{total}} < 0.090 \frac{\mu}{M} X_0^2 \left(1 - \frac{2^{2/3} \epsilon^{4/3}}{X_0^2} \right) \quad (24\text{--new})$$

$$\Delta \left| \frac{a}{M} \right|_{\text{total}} < 0.045 \frac{\mu}{M} X_0^2. \quad (25\text{--new})$$

The most important difference is that the change in spin is now linear in the mass ratio, as opposed to quadratic. Note also that the new left-hand-side of (25) now correctly refers to the absolute value of a/M , which is necessary since a/M decreases during near-horizon inspiral. We thank Geoffrey Compère for pointing out these errors—see also reference [1].

The change in spin over the near-horizon inspiral is significantly larger than originally calculated, meaning that there is a smaller range of mass-ratios over which the near-horizon inspiral can occur (without spinning down the black hole such that the near-horizon region no longer exists). We may estimate this range based on the observation (section 3.1) that near-horizon inspiral is visible in the waveform for $a \gtrsim 0.9999M$ with a starting radius of $X_0 \approx 0.3$. Demanding $\Delta|a/M|_{\text{total}} < 0.0001$ using $X_0 = 0.3$, the new version of (25) gives the bound $\mu/M < 0.025$. Thus near-horizon inspiral remains consistent for both the LIGO and LISA sources discussed in the appendix (with mass ratios of order 10^{-3} and smaller), and our main conclusions about observability are unmodified.

ORCID iDs

Samuel E Gralla  <https://orcid.org/0000-0002-7468-9731>

Reference

- [1] Compère G, Fransen K and Jonas C 2019 arXiv:1909.12848 [gr-qc]