

Periodic wave solutions and stability analysis for the KP-BBM equation with abundant novel interaction solutions

Jalil Manafian^{1,5} , Onur Alp Ilhan²  and As'ad Alizadeh^{3,4} 

¹ Department of Applied Mathematics, Faculty of Mathematical Sciences, University of Tabriz, Tabriz, Iran

² Department of Mathematics, Faculty of Education, Erciyes University, 38039-Melikgazi-Kayseri, Turkey

³ Department of Mechanical Engineering, Urmia University of Technology, Urmia, Iran

⁴ Department of Mechanical Engineering, College of Engineering, University of Zakho, Zakho, Iraq

E-mail: j_manafianheris@tabrizu.ac.ir, oailhan@erciyes.edu.tr and asad.alizadeh2010@gmail.com

Received 13 October 2019, revised 31 December 2019

Accepted for publication 7 January 2020

Published 20 March 2020



Abstract

This paper aims at investigating periodic wave solutions for the (2+1)-dimensional KP-BBM equation, from its bilinear form, obtained using the Hirota operator. Two major cases were studied from two different ansatzes. The 3D, 2D and density representation illustrating some cases of solutions obtained have been represented from a selection of the appropriate parameters. The modulation instability is employed to discuss the stability of got solutions. That will be extensively used to report many attractive physical phenomena in the fields of acoustics, heat transfer, fluid dynamics, classical mechanics and so on.

Keywords: KP-BBM equation, Hirota bilinear operator method, periodic wave solution, modulation instability

(Some figures may appear in colour only in the online journal)

1. Introduction

Nonlinear phenomena play a very important role in the fields of mathematics, chemistry, biology, and physics sciences. These phenomena are for the most part modeled by nonlinear mathematical equations. For instance, in nonlinear physics, physical mechanisms of natural phenomena in the field of applied sciences and engineering, especially in plasma physics, elastic media, optical fibers, fluid dynamics, quantum mechanics, chimerical physics, biotechnology, signal processing, solid state physics, shallow water wave theory, are modeled by nonlinear partial differential equations (NLPDEs). However, the quest for the exact explicit solutions of these equations remains a hot topic. Moreover, looking for localized solutions and more specifically the solitary wave solutions [1–9], the lumps-type solutions [10–36], the interaction soliton-soliton, soliton-kink and kink-kink [37, 38], the interactions between solitary wave solutions and lump solutions [39–41], as well as the periodic wave solutions [42–44]

remains a very interesting subject for researchers. Several mathematical methods are used in the search for these solutions. For example, the Exp-function method [45, 46], the Homotopy perturbation technique [47], the inverse scattering method [48] and so on. For getting the lump solutions and their interactions authors have conjugated sufficient time to search the exact rational soliton solutions, for example, the nonlinear evolution equations [14], the Kadomtsev-Petviashvili (KP) equation [15], the reduced pgKP and pgbKP equations [17], the (2+1)-dimensional KdV equation [18], the (2+1)-dimensional Sawada-Kotera equation [34], the (2+1)-dimensional bSK equation [35, 36], the (2 + 1)-dimensional generalized fifth order KdV equation [37], the (2+1)-dimensional Burger equation [38], the generalized (3+1)-dimensional Shallow water-like equation [40], and the B-Kadomtsev-Petviashvili equation [46]. Various types of work for finding the periodic solitary wave solutions on the interaction between lump and other kinds of solitary, periodic and kink solitons for the (2+1)-dimensional Breaking Soliton equation [19], lump and interaction between different types of those on the variable-coefficient Kadomtsev-Petviashvili equation [20], and periodic

⁵ Author to whom any correspondence should be addressed.

type and periodic cross-kink wave solutions [21], and the (2+1)-dimensional extended Jimbo-Miwa equations [42] are achieved through the Hirota bilinear operator. Baskonus and co-authors constructed a family of wave solutions for some of nonlinear equations such as doubly dispersive equation [22], the Gilson-Pickering model [23], a (2+1)-dimensional coupling system with KdV equation [24], the Lonngren-Wave equation [25], the Ablowitz-Kaup-Newell-Segur equation [26], the Wu-Zhang system [27], the longitudinal wave equation in a magneto-electro-elastic circular rod [28], the generalized double combined Sinh-Cosh-Gordon equation [29], the Sharma-Tasso-Olver equation [30], by utilizing the extended sinh-Gordon equation expansion method, the modified $\exp(-\varphi(\zeta))$ -expansion function method and the improved Bernoulli sub-equation function method. In [31], Yukus and co-authors used the modified $\exp(-\varphi(\zeta))$ -expansion function method to construct some new analytical solutions with novel structure such as the trigonometric and hyperbolic function solutions. Shafiq, Rashidi, Hammouch and Khan [32] investigated the mixed convective stagnation point flow of Williamson liquid over a vertical stretched plate and convergent solutions for the temperature and velocity were constructed and analyzed. The bilinear method was employed to investigate the rogue wave solutions and the rogue type multiple lump wave solutions of the (2+1)-dimensional Benjamin-Ono equation by Zhao, He, and Gao [33].

The aim of this study is to construct the invariant solutions of the (2+1)-dimensional Kadomtsev-Petviashvili-Benjamin-Bona-Mahony (KP-BBM) equation of the form

$$u_{xt} + u_{xx} + \alpha(u^2)_{xx} + \beta u_{xxx} + \gamma u_{yy} = 0, \quad (1.1)$$

where α , β and γ the arbitrary constants. Equation (1.1) is formulated using the Kadomtsev-Petviashvili (KP) equation in which reads as [15]

$$(u_t + 6uu_x + u_{xxx})_x - u_{yy} = 0, \quad (1.2)$$

and also derived from the standard BBM equation [49] with below form

$$u_t + u_x + uu_x - u_{xx} = 0. \quad (1.3)$$

For equation (1.2), Ma [15] obtained the lump solutions by using the following transformation

$$u = 2(\ln f)_{xx}, \quad (1.4)$$

and also Manakov *et al* [50] acquired subclass of the lump solutions in which involving two free parameters. Besides, Zhao and Ma [51] used the Hirota bilinear form of the KP equation and gained twelve classes of the lump-kink solutions. Sufficient literature for KP-BBM equation (1.1) is provided here notably Wazwaz [52] constructed the new compact and noncompact of the KP-BBM and the ZK-BBM equations with the aid of extended tanh method in which constitutes the analytical method containing the expansion of tanh series. Saut and Tzvetkov [53] obtained localized solitary solutions via global well-posedness structures from the conservation

law and generated some soliton solutions depending upon the appropriate choice of the arbitrary functions. Continuing, Alam and Akbar [54] proposed the generalized (G'/G) -expansion method of Riccati equation and used it to establish the soliton-like solutions with the aid of symbolic computation. Later on, Tang *et al* [55] employed bifurcations of the travelling wave solutions to the KP-BBM equation and obtained the exact solutions. Some the exact lump solution, lump-kink solutions, lump-soliton solutions, and breather-wave solutions were examined in [56] by means of symbolic computation and the Hirota bilinear method. As remarkable work, the Landau-Lifshitz-Gilbert equation had been considered to study the propagation and interaction of magnetic solitons in a ferromagnetic thin film with the interfacial Dzyaloshinskii-Moriya interaction [57] and found the analytical forms for magnetic breathers and the first- to third-order rogue wave solutions [58]. Likewise, Deng and coauthors [59] studied the interaction phenomenon between the lump waves and stripe solitons to the (2+1)-dimensional Caudrey-Dodd-Gibbon-Kotera-Sawada equation by making use of the Hirota bilinear method. In [60], Dai and co-authors proved the existence of multi-wave solutions and obtained three-wave solution, periodic two-solitary-wave solutions, for the (1+2)-D Kadomtsev-Petviashvili equation via the extended three-soliton method.

Consider the Bell polynomial of Kadomtsev-Petviashvili-Benjamin-Bona-Mahony equation as follows

$$\mathbb{P}_{\text{KP-BBM}}(u) := u_{xt} + u_{xx} + \alpha(u^2)_{xx} + \beta u_{xxx} + \gamma u_{yy} = 0. \quad (1.5)$$

Let the Hirota derivatives in terms of the functions f and g can be written as

$$\prod_{i=1}^3 D_{j_i}^{\beta_i} f \cdot g = \prod_{i=1}^3 \left(\frac{\partial}{\partial j_i} - \frac{\partial}{\partial j'_i} \right)^{\beta_i} f(j) g(j') \Big|_{j'=j}, \quad (1.6)$$

where the vectors $J = (j_1, j_2, j_3) = (x, y, t)$, $J' = (j'_1, j'_2, j'_3) = (x', y', t')$ and $\beta_1, \beta_2, \beta_3$ are the arbitrary nonnegative integers, and its corresponding bilinear formalism equals. It is known that the KP-BBM equation possesses a Hirota bilinear form:

$$\begin{aligned} \mathbb{B}_{\text{KP-BBM}}(f) &:= (D_x D_t + D_x^2 + \beta D_x^3 D_t \\ &+ \gamma D_y^2 + \alpha D_x^2 - 3\beta D_x D_t) f \cdot f \\ &= 2[(1 - 3\beta)(ff_{xt} - f_x f_t) \\ &+ (1 + \alpha)(ff_{xx} - f_x^2) + \gamma(ff_{yy} - f_y^2) \\ &+ \beta(ff_{xxx} - 3f_x f_{xt} + 3f_{xx} f_t - f_t f_{xxx})] = 0. \end{aligned} \quad (1.7)$$

Use the following transformation between the function f and u :

$$u = 2(\ln f)_{xx}. \quad (1.8)$$

Employing the Bell polynomial theories of soliton equations, we achieve to the following relation as

$$\mathbb{P}_{\text{KP-BBM}}(u) = \left[\frac{\mathbb{B}_{\text{KP-BBM}}(f)}{f^2} \right]_{xx}. \quad (1.9)$$

Theorem 1.1. f solves (1.8) if and only if $u = 2(\ln f)_{xx}$ demonstrate a solution to the KP-BBM equation (1.5)

$$\begin{aligned} & (D_x D_t + D_x^2 + \beta D_x^3 D_t + \gamma D_y^2 + \alpha D_x^2 \\ & - 3\beta D_x D_t) f \cdot f = 2[(1 - 3\beta)(ff_{xt} - f_x f_t) \\ & + (1 + \alpha)(ff_{xx} - f_x^2) + \gamma(ff_{yy} - f_y^2) \\ & + \beta(ff_{xxx} - 3f_x f_{xxt} + 3f_{xx} f_{xt} - f_x f_{xxx})] = 0. \end{aligned} \quad (1.10)$$

Our purpose here is to discover the exact solutions of the KP-BBM equation under consideration the Hirota bilinear

$$H_5 = \sin(\Omega_4 x + \Omega_5 y + \Omega_6 t),$$

$$H_6 = \sinh(\Omega_7 x + \Omega_8 y + \Omega_9 t),$$

$$\begin{aligned} u &= 2 \frac{\partial^2}{\partial x^2} \ln(f) \\ &= 2 \frac{a_1 \Omega_1^2 H_1 + a_2 \Omega_1^2 H_2 - a_3 H_3 \Omega_4^2 + a_4 H_4 \Omega_7^2}{f} \\ &\quad - 2 \frac{(a_1 \Omega_1 H_1 - a_2 \Omega_1 H_2 - a_3 H_5 \Omega_4 + a_4 H_6 \Omega_7)^2}{f^2}, \end{aligned} \quad (2.3)$$

where $\Omega_i, i = 1, \dots, 9, a_i, j = 1, \dots, 4$, are free parameters in which are to find later. Plugging (2.3) into equation (1.10) and then collecting the coefficients, we get to the following results:

$$\begin{cases} \omega \Omega_4^4 - 6 \omega \Omega_4^2 \Omega_7^2 + \omega \Omega_7^4 + 4 \Omega_4 \Omega_6 - 3 \Omega_5^2 - 4 \Omega_7 \Omega_9 + 3 \Omega_8^2 = 0 \\ \omega \Omega_1^4 + 6 \omega \Omega_1^2 \Omega_7^2 + \omega \Omega_7^4 - 4 \Omega_1 \Omega_3 + 3 \Omega_2^2 - 4 \Omega_7 \Omega_9 + 3 \Omega_8^2 = 0 \\ 2 \omega \Omega_4^3 \Omega_7 - 2 \omega \Omega_4 \Omega_7^3 + 2 \Omega_4 \Omega_9 - 3 \Omega_5 \Omega_8 + 2 \Omega_6 \Omega_7 = 0, \\ 2 \omega \Omega_1^3 \Omega_7 + 2 \omega \Omega_1 \Omega_7^3 - 2 \Omega_1 \Omega_9 + 3 \Omega_2 \Omega_8 - 2 \Omega_3 \Omega_7 = 0 \\ \omega \Omega_1^4 - 6 \omega \Omega_1^2 \Omega_4^2 + \omega \Omega_4^4 - 4 \Omega_1 \Omega_3 + 3 \Omega_2^2 + 4 \Omega_4 \Omega_6 - 3 \Omega_5^2 = 0 \\ 2 \omega \Omega_1^3 \Omega_4 - 2 \omega \Omega_1 \Omega_4^3 - 2 \Omega_1 \Omega_6 + 3 \Omega_2 \Omega_5 - 2 \Omega_3 \Omega_4 = 0 \\ 16 \omega \Omega_1^4 a_1 a_2 + 4 \omega \Omega_4^4 a_3^2 + 4 \omega \Omega_7^4 a_4^2 - 16 \Omega_1 \Omega_3 a_1 a_2 + 12 \Omega_2^2 a_1 a_2 + 4 \Omega_4 \Omega_6 a_3^2 - 3 \Omega_5^2 a_3^2 - 4 \Omega_7 \Omega_9 a_4^2 + 3 \Omega_8^2 a_4^2 = 0. \end{cases} \quad (2.4)$$

method for getting the novel periodic solutions in which can be arisen with twenty one classes. Discussion about the nonlinear KP-BBM equation and the Hirota bilinear method are given. Also, the modulation instability of the KP-BBM equation is offered. In the continuation, we will offer the graphical illustrations of some solutions of the considered model along with the obtained solutions. After that, we will deal with the probe of solutions and we will finish by a conclusion.

2. New periodic-waves solutions for KP-BBM equation

Here, we will compose the periodic-waves solutions of equation (1.1), next three waves hypothesis can be discovered through employing the Hirota operator [47]. The solution can be expressed in the below as:

$$f = a_1 H_1 + a_2 H_2 + a_3 H_3 + a_4 H_4, \quad (2.1)$$

$$\begin{aligned} H_1 &= \exp(\Omega_1 x + \Omega_2 y + \Omega_3 t), \\ H_2 &= \exp(-\Omega_1 x - \Omega_2 y - \Omega_3 t), \\ H_3 &= \cos(\Omega_4 x + \Omega_5 y + \Omega_6 t), \\ H_4 &= \cosh(\Omega_7 x + \Omega_8 y + \Omega_9 t), \end{aligned} \quad (2.2)$$

Solving the equations we get:

Case I:

$$\begin{aligned} u_1 &= 2 \frac{a_1 \Omega_1^2 H_1 + a_4 H_4 \Omega_7^2}{f} - 2 \frac{(a_1 \Omega_1 H_1 + a_4 H_6 \Omega_7)^2}{f^2}, \\ f &= a_1 H_1 + a_4 H_4, \end{aligned} \quad (2.5)$$

$$\begin{aligned} H_1 &= e^{\Omega_1 x + \frac{\sqrt{-\beta \gamma (3 \alpha \beta - \alpha + 3 \beta - 1)}}{\beta \gamma} y - \frac{3 \Omega_1 (\alpha + 1)}{4 \beta \Omega_1^2 - 3 \beta + 1} t}, \\ H_4 &= \cosh \left(\frac{\sqrt{-3 \beta (\beta \Omega_1^2 - 3 \beta + 1)}}{3 \beta} x \right. \\ &\quad \left. + \frac{3 \Omega_7 (\alpha + 1)}{4 \beta \Omega_1^2 - 3 \beta + 1} t \right). \end{aligned} \quad (2.6)$$

Also, the existence condition of the solution is that

$$\begin{aligned} \beta (\beta \Omega_1^2 - 3 \beta + 1) &< 0, \\ \beta \gamma (3 \alpha \beta - \alpha + 3 \beta - 1) &< 0, \\ 4 \beta \Omega_1^2 - 3 \beta + 1 &\neq 0. \end{aligned}$$

Case II:

$$\begin{aligned} u_2 &= 2 \frac{-a_3 H_3 \Omega_4^2 + a_4 H_4 \Omega_7^2}{f} \\ &\quad - 2 \frac{(-a_3 H_5 \Omega_4 + a_4 H_6 \Omega_7)^2}{f^2}, \quad f = a_3 H_3 + a_4 H_4, \end{aligned} \quad (2.7)$$

$$H_3 = \cos \left(\sqrt{-\frac{3\beta(\alpha+1)}{\gamma(\beta\Omega_7^2 a_3^2 - 4\beta\Omega_7^2 a_4^2 - 3\beta a_3^2 + 3\beta a_4^2 + a_3^2 - a_4^2)}} a_4 \Omega_7^2 y \right),$$

$$H_4 = \cosh \left(\Omega_7 x - \frac{(\alpha a_3^2 - \alpha a_4^2 + a_3^2 - a_4^2) \Omega_7}{\beta\Omega_7^2 a_3^2 - 4\beta\Omega_7^2 a_4^2 - 3\beta a_3^2 + 3\beta a_4^2 + a_3^2 - a_4^2} t \right).$$

Also, the existence condition of the solution is that

$$\gamma(\beta\Omega_7^2 a_3^2 - 4\beta\Omega_7^2 a_4^2 - 3\beta a_3^2 + 3\beta a_4^2 + a_3^2 - a_4^2)\beta(\alpha+1) < 0.$$

Case III:

$$u_3 = 2 \frac{a_4 H_4 \Omega_7^2}{f} - 2 \frac{(a_4 H_6 \Omega_7)^2}{f^2}, \quad f = a_3 H_3 + a_4 H_4, \quad (2.8)$$

$$H_3 = \cos \left(\frac{\sqrt{\beta\gamma(3\alpha\beta - \alpha + 3\beta - 1)}}{\beta\gamma} y + \Omega_6 t \right),$$

$$H_4 = \cosh \left(\frac{\sqrt{\beta(3\beta - 1)}}{\beta} x + \frac{\alpha a_3^2 - \alpha a_4^2 + a_3^2 - a_4^2}{\sqrt{\beta(3\beta - 1)} a_4^2} t \right).$$

Also, the existence condition of the solution is that

$$\beta\gamma(3\alpha\beta - \alpha + 3\beta - 1) > 0, \quad \beta(3\beta - 1) > 0.$$

Case IV:

$$u_4 = 2 \frac{a_1 \Omega_1^2 H_1 - a_3 H_3 \Omega_4^2}{f} - 2 \frac{(a_1 \Omega_1 H_1 - a_3 H_5 \Omega_4)^2}{f^2},$$

$$f = a_1 H_1 + a_3 H_3, \quad (2.9)$$

$$H_1 = e^{\Omega_1 x + \frac{\sqrt{-\beta\gamma(3\alpha\beta - \alpha + 3\beta - 1)}}{\beta\gamma} y - \frac{3\Omega_1(\alpha+1)}{4\beta\Omega_1^2 - 3\beta + 1} t},$$

$$H_3 = \cos \left(\frac{\sqrt{3\beta(\beta\Omega_1^2 - 3\beta + 1)}}{3\beta} x + \frac{\sqrt{3\beta(\beta\Omega_1^2 - 3\beta + 1)(\alpha+1)}}{\beta(4\beta\Omega_1^2 - 3\beta + 1)} t \right).$$

Moreover, the existence condition of the solution is that

$$\beta(\beta\Omega_1^2 - 3\beta + 1) > 0,$$

$$\beta\gamma(3\alpha\beta - \alpha + 3\beta - 1) < 0.$$

Case V:

$$u_5 = 2 \frac{a_4 H_4 \Omega_7^2}{f} - 2 \frac{(a_4 H_6 \Omega_7)^2}{f^2}, \quad f = a_3 H_3 + a_4 H_4, \quad (2.10)$$

$$H_3 = \cos(\Omega_6 t),$$

$$H_4 = \cosh \left(\frac{\sqrt{\beta(3\beta - 1)}}{\beta} x + \frac{\sqrt{-\beta\gamma(3\alpha\beta - \alpha + 3\beta - 1)}}{\beta\gamma} y \right).$$

Moreover, the existence condition of the solution is that

$$\beta(3\beta - 1) > 0, \quad \beta\gamma(3\alpha\beta - \alpha + 3\beta - 1) < 0.$$

Case VI:

$$u_5 = 2 \frac{a_1 \Omega_1^2 H_1 + a_2 \Omega_1^2 H_2 - a_3 H_3 \Omega_4^2 + a_4 \Omega_7^2}{f} - 2 \frac{(a_1 \Omega_1 H_1 - a_2 \Omega_1 H_2 - a_3 H_5 \Omega_4)^2}{f^2}, \quad (2.11)$$

$$f = a_1 H_1 + a_2 H_2 + a_3 H_3 + a_4,$$

$$H_1 = e^{\Omega_1 x - \frac{\Omega_1(\alpha+1)}{\beta\Omega_1^2 - 3\beta + 1} t},$$

$$H_2 = e^{-\Omega_1 x + \frac{\Omega_1(\alpha+1)}{\beta\Omega_1^2 - 3\beta + 1} t},$$

$$H_3 = \cos \left(\frac{\sqrt{-\beta(3\beta - 1)}}{\beta} x + \frac{\sqrt{\beta\gamma(3\alpha\beta - \alpha + 3\beta - 1)}}{\beta\gamma} y + \frac{(\alpha+1)\sqrt{-\beta(3\beta - 1)}}{\beta(\beta\Omega_1^2 - 3\beta + 1)} t \right), \quad H_4 = 1.$$

Moreover, the existence condition of the solution is that

$$\beta(3\beta - 1) < 0, \quad \beta\gamma(3\alpha\beta - \alpha + 3\beta - 1) > 0.$$

Case VII:

$$u_7 = 2 \frac{a_1 \Omega_1^2 H_1 + a_2 \Omega_1^2 H_2 + a_4 H_4 \Omega_7^2}{f} - 2 \frac{(a_1 \Omega_1 H_1 - a_2 \Omega_1 H_2 + a_4 H_6 \Omega_7)^2}{f^2}, \quad (2.12)$$

$$\begin{aligned} \mathbf{f} &= -\frac{3\beta a_3^2 - 3\beta a_4^2 - a_3^2 + a_4^2}{4(2\beta\Omega_1^2 - 3\beta + 1)a_2}H_1 \\ &\quad + a_2H_2 + a_3H_3 + a_4H_4, \\ H_1 &= e^{\Omega_1 x + \frac{\sqrt{-3\beta\gamma(\alpha\beta\Omega_1^2 + \beta\Omega_1^2 - 3\alpha\beta + \alpha - 3\beta + 1)}}{3\beta\gamma}y - \frac{2(\alpha+1)}{3\beta\Omega_1}t}, \\ H_2 &= e^{-\Omega_1 x - \frac{\sqrt{-3\beta\gamma(\alpha\beta\Omega_1^2 + \beta\Omega_1^2 - 3\alpha\beta + \alpha - 3\beta + 1)}}{3\beta\gamma}y + \frac{2(\alpha+1)}{3\beta\Omega_1}t}, \\ H_4 &= \cosh\left(\frac{\sqrt{\beta(3\beta - 1)}}{\beta}x\right), \end{aligned}$$

Moreover, the existence condition of the solution is that

$$\begin{aligned} \beta(3\beta - 1) &> 0, \\ (3\beta\Omega_1^2 - 9\beta + 3)(3\beta - 1) &< 0, \\ (\beta\Omega_1^2 - 3\beta + 1)\Omega_1 &\neq 0. \end{aligned}$$

Case IX:

$$\begin{aligned} H_3 &= \cos\left(\frac{\sqrt{\beta\gamma(3\alpha\beta - \alpha + 3\beta - 1)}}{\beta\gamma}y\right. \\ &\quad \left.- \frac{2}{3}\frac{\sqrt{\beta\gamma(3\alpha\beta - \alpha + 3\beta - 1)}\sqrt{-3\beta\gamma(\alpha\beta\Omega_1^2 + \beta\Omega_1^2 - 3\alpha\beta + \alpha - 3\beta + 1)}}{\beta^2\gamma(\beta\Omega_1^2 - 3\beta + 1)\Omega_1}t\right). \end{aligned}$$

Moreover, the existence condition of the solution is that

$$\begin{aligned} (2\beta\Omega_1^2 - 3\beta + 1)a_2 &\neq 0, \\ \beta\gamma(3\alpha\beta - \alpha + 3\beta - 1) &> 0, \\ \beta\gamma(\alpha\beta\Omega_1^2 + \beta\Omega_1^2 - 3\alpha\beta + \alpha - 3\beta + 1) &< 0. \end{aligned}$$

Case VIII:

$$\begin{aligned} \mathbf{u}_8 &= 2\frac{a_2\Omega_1^2H_2 + a_4H_4\Omega_7^2}{f} \\ &\quad - 2\frac{(-a_2\Omega_1H_2 + a_4H_6\Omega_7)^2}{f^2}, \\ \mathbf{f} &= a_2H_2 - a_4H_3 + a_4H_4, \end{aligned} \quad (2.13)$$

$$H_2 = e^{-\Omega_1 x + \frac{\sqrt{-(3\beta\Omega_1^2 - 9\beta + 3)(3\beta - 1)(\alpha + 1)}}{3\sqrt{\beta\gamma(3\alpha\beta - \alpha + 3\beta - 1)}}y + \frac{2}{3}\frac{\alpha + 1}{\beta\Omega_1}t},$$

$$\begin{aligned} H_3 &= \cos\left(\frac{\sqrt{\beta\gamma(3\alpha\beta - \alpha + 3\beta - 1)}}{\beta\gamma}y\right. \\ &\quad \left.+ \frac{2}{3}\frac{\sqrt{-(3\beta\Omega_1^2 - 9\beta + 3)(3\beta - 1)(\alpha + 1)}}{(\beta\Omega_1^2 - 3\beta + 1)\Omega_1\beta}t\right), \\ H_4 &= \cosh\left(\frac{\sqrt{\beta(3\beta - 1)}}{\beta}x\right). \end{aligned}$$

$$\begin{aligned} \mathbf{u}_9 &= 2\frac{\left(\frac{a_4^2}{4a_2}H_1 + a_2H_2 + a_4H_4\right)\Omega_1^2}{f} \\ &\quad - 2\frac{\left(\frac{a_4^2}{4a_2}\Omega_1H_1 - a_2\Omega_1H_2 + a_4H_6\Omega_1\right)^2}{f^2}, \\ f &= \frac{a_4^2}{4a_2}H_1 + a_2H_2 + a_4H_4, \end{aligned} \quad (2.14)$$

$$\begin{aligned} H_1 &= e^{\Omega_1 x + \Omega_8 y - \frac{4\beta\Omega_1^3\Omega_9 + 2\alpha\Omega_1^2 - 3\beta\Omega_1\Omega_9 + 2\gamma\Omega_8^2 + 2\Omega_1^2 + \Omega_1\Omega_9}{\Omega_1(4\beta\Omega_1^2 - 3\beta + 1)}t}, \\ H_2 &= e^{-\Omega_1 x - \Omega_8 y + \frac{4\beta\Omega_1^3\Omega_9 + 2\alpha\Omega_1^2 - 3\beta\Omega_1\Omega_9 + 2\gamma\Omega_8^2 + 2\Omega_1^2 + \Omega_1\Omega_9}{\Omega_1(4\beta\Omega_1^2 - 3\beta + 1)}t}, \end{aligned}$$

$$\begin{aligned} H_4 &= \cosh(\Omega_1 x + \Omega_8 y + \Omega_9 t), \\ H_6 &= \sinh(\Omega_1 x + \Omega_8 y + \Omega_9 t), \\ \Omega_1(4\beta\Omega_1^2 - 3\beta + 1) &\neq 0. \end{aligned}$$

Case X:

$$\begin{aligned} \mathbf{u}_{10} &= 2\frac{a_2\Omega_1^2H_2 - a_3H_3\Omega_4^2 + a_4H_4\Omega_7^2}{f} \\ &\quad - 2\frac{(-a_2\Omega_1H_2 - a_3H_5\Omega_4 + a_4H_6\Omega_7)^2}{f^2}, \\ f &= a_2H_2 + a_3H_3 + a_4H_4, \end{aligned} \quad (2.15)$$

$$H_2 = e^{-\Omega_1 x - \frac{\Omega_1(6\alpha\beta\Omega_1^2 + \beta\gamma\Omega_8^2 + 6\beta\Omega_1^2 - 6\alpha\beta + 2\alpha - 6\beta + 2)}{3\Omega_8\gamma\beta\Omega_4}y - \frac{-36\beta^2\Omega_1^6(\alpha+1)^2 + \Xi_1 + \Xi_2 + \Xi_3 + \Xi_4}{18\beta(4\beta\Omega_1^2 - 3\beta + 1)\gamma\Omega_1\Omega_8^2(3\beta\Omega_1^2 - 3\beta + 1)}t},$$




Figure 1. Diagram of periodic-waves (2.5) using values $a_1 = .5$, $a_4 = 1$, $\Omega_1 = 1.5$, $\alpha = 1$, $\beta = 2$, $\gamma = -1$, $y = -4$, and (a) 3D plot, (b) density plot, and (c) 2D plot with (red $x = -10$, blue $x = 0$, and green $x = 10$).




Figure 2. Diagram of periodic-waves (2.8) using values $a_3 = 2.5$, $a_4 = 1$, $\Omega_7 = 1.3$, $\alpha = -2$, $\beta = 1$, $\gamma = 1$, $y = -4$, and (a) 3D plot, (b) density plot, and (c) 2D plot with (red $x = -2$, blue $x = 0$, and green $x = 2$).

$$\begin{aligned}\Xi_1 &= -6 \Omega_1^4 \beta (\alpha + 1) \\ &\times (11 \beta \gamma \Omega_8^2 - 12 \alpha \beta + 4 \alpha - 12 \beta + 4), \\ \Xi_2 &= \gamma \Omega_1^2 \Omega_8^2 \beta (26 \beta \gamma \Omega_8^2 + 147 \alpha \beta - 49 \alpha + 147 \beta),\end{aligned}$$

$$\begin{aligned}\Xi_3 &= -\Omega_1^2 (36 \alpha^2 \beta^2 + 49 \beta \gamma \Omega_8^2 - 24 \alpha^2 \beta \\ &+ 72 \alpha \beta^2 + 4 \alpha^2 - 48 \alpha \beta + 36 \beta^2 + 8 \alpha - 24 \beta + 4), \\ \Xi_4 &= -9 \gamma \Omega_8^2 (3 \beta - 1) (\beta \gamma \Omega_8^2 + 3 \alpha \beta - \alpha + 3 \beta - 1),\end{aligned}$$

$$\begin{aligned}H_3 &= \cos \left(\frac{\sqrt{-\beta(3\beta\Omega_1^2 - 3\beta + 1)}}{\beta} x + \frac{\Omega_8 \Omega_4}{\Omega_7} y \right. \\ &\left. - \frac{(-3\alpha\beta\Omega_1^2 + \beta\gamma\Omega_8^2 - 3\beta\Omega_1^2 + 3\alpha\beta - \alpha + 3\beta - 1)}{3\sqrt{\beta(3\beta\Omega_1^2 - 3\beta + 1)(4\beta\Omega_1^2 - 3\beta + 1)}} t \right),\end{aligned}$$

$$\begin{aligned}H_4 &= \cosh \left(\frac{\sqrt{-\beta(3\beta\Omega_1^2 - 3\beta + 1)}}{\beta} x + \Omega_8 y \right. \\ &\left. + \frac{-3\alpha\beta\Omega_1^2 + \beta\gamma\Omega_8^2 - 3\beta\Omega_1^2 + 3\alpha\beta - \alpha + 3\beta - 1}{3\sqrt{-\beta(3\beta\Omega_1^2 - 3\beta + 1)(4\beta\Omega_1^2 - 3\beta + 1)}} t \right).\end{aligned}$$

Case XI:

$$\begin{aligned}u_{11} &= 2 \frac{a_1 \Omega_1^2 H_1 + a_4 H_4 \Omega_7^2}{f} \\ &- 2 \frac{(a_1 \Omega_1 H_1 + a_4 H_6 \Omega_7)^2}{f^2}, \\ f &= a_1 H_1 + \frac{\sqrt{5(2\beta\Omega_1^2 + 3\beta - 1)(4\beta\Omega_1^2 - 3\beta + 1)}}{2\beta\Omega_1^2 + 3\beta - 1} H_3 \\ &+ a_4 H_4,\end{aligned}\tag{2.16}$$

$$\begin{aligned}H_2 &= e^{\Omega_1 x + \Omega_2 y + \Omega_3 t}, \\ H_3 &= \cos(\Omega_5 y + \Omega_6 t), \\ H_4 &= \cosh(\Omega_7 x + \Omega_8 y + \Omega_9 t),\end{aligned}$$




Figure 3. Diagram of periodic-waves (2.8) using values $a_3 = 1.1$, $a_4 = 1$, $\Omega_7 = 1.3$, $\alpha = -1$, $\beta = -2$, $\gamma = -1$, $y = -4$, and (a) 3D plot, (b) density plot, and (c) 2D plot with (red $x = -1.3$, blue $x = 0$, and green $x = 1.3$).




Figure 4. Diagram of periodic-waves (2.9) using values $a_1 = 0.5$, $a_3 = 1$, $\Omega_1 = 1.5$, $\alpha = -2$, $\beta = 2$, $\gamma = 1$, $y = -4$, and (a) 3D plot, (b) density plot, and (c) 2D plot with (red $x = -1$, blue $x = 0$, and green $x = 1$).




Figure 5. Diagram of periodic-waves (2.10) using values $a_3 = 0.2\gamma$, $a_4 = 1$, $\Omega_6 = 1.5$, $\alpha = -2$, $\beta = 2$, $\gamma = 1$, $y = -4$, and (a) 3D plot, (b) density plot, and (c) 2D plot with (red $x = -1$, blue $x = 0$, and green $x = 1$).

$$\Omega_2 = -\frac{(3\alpha\beta^2\Omega_1^4 + 3\beta^2\Omega_1^4 - 12\alpha\beta^2\Omega_1^2 + 4\alpha\beta\Omega_1^2 - 12\beta^2\Omega_1^2 + 9\alpha\beta^2 + 4\beta\Omega_1^2 - 6\alpha\beta + 9\beta^2 + \alpha - 6\beta + 1)\Omega_1}{\Omega_7(7\beta\Omega_1^2 - 3\beta + 1)\beta\gamma\Omega_8},$$




Figure 6. Diagram of periodic-waves (2.11) using values $a_2 = 0.5$, $a_3 = 1.5$, $a_4 = 1$, $\Omega_1 = 1.5$, $\alpha = 1$, $\beta = \frac{1}{4}$, $\gamma = -1$, $y = -4$, and (a) 3D plot, (b) density plot, and (c) 2D plot with (red $x = -1$, blue $x = 0$, and green $x = 1$).




Figure 7. Diagram of periodic-waves (2.12) using values $a_2 = 0.5$, $a_3 = 1.5$, $a_4 = 1$, $\Omega_1 = 1.5$, $\alpha = -2$, $\beta = \frac{1}{4}$, $\gamma = 1$, $y = -4$, and (a) 3D plot, (b) density plot, and (c) 2D plot with (red $x = -1$, blue $x = 0$, and green $x = 1$).




Figure 8. Diagram of periodic-waves (2.13) using values $a_2 = 0.5$, $a_4 = 1.4$, $\Omega_1 = 1.5$, $\alpha = -2$, $\beta = 2$, $\gamma = -1$, $y = -4$, and (a) 3D plot, (b) density plot, and (c) 2D plot with (red $x = -1$, blue $x = 0$, and green $x = 1$).




Figure 9. The dispersion relation between frequency $W(M, N)$ and wave number M under the various values $\alpha = 1, \beta = 2, \gamma = 3, q = 1$.




Figure 10. The dispersion relation between frequency $W(M, N)$ and wave number M under the various values (left) $\alpha = -3, \beta = 1, \gamma = 3, q = 1$ (right) $\alpha = 1, \beta = 2, \gamma = -3, q = 1$.




Figure 11. The dispersion relation between frequency $W(M, N)$ and wave number M under the various values (left) $\alpha = -1, \beta = 2, \gamma = -3, q = -1$ (right) $\alpha = 1, \beta = 0.2, \gamma = -3, q = -1$.

$$\begin{aligned}\Omega_3 &= -\frac{2(\alpha+1)(\beta\Omega_1^2-3\beta+1)}{3\Omega_1\beta(7\beta\Omega_1^2-3\beta+1)}, & \Omega_7 &= \frac{\sqrt{-\beta(3\beta\Omega_1^2-3\beta+1)}}{\beta}, \\ \Omega_5 &= 3\frac{\Xi_1(\alpha+1)\Omega_7\Omega_1}{(7\beta\Omega_1^2-3\beta+1)\gamma\Omega_8}, & \Omega_8 &= \frac{\sqrt{2\beta\gamma(3\alpha\beta\Omega_1^2+3\beta\Omega_1^2-3\alpha\beta+\alpha-3\beta+1)}}{\beta\gamma}, \\ \Omega_6 &= 2\frac{\Xi_1(\alpha+1)}{\Omega_1\beta(7\beta\Omega_1^2-3\beta+1)}, & \Omega_9 &= \frac{3\alpha\beta\Omega_1^2+3\beta\Omega_1^2-3\alpha\beta+\alpha-3\beta+1}{3\Omega_1\beta(4\beta\Omega_1^2-3\beta+1)}.\end{aligned}$$

$$\begin{aligned}\Omega_7 &= \frac{\sqrt{-\beta(3\beta\Omega_1^2-3\beta+1)}}{\beta}, \\ \Omega_8 &= \frac{\sqrt{3\gamma(7\beta\Omega_1^2-3\beta+1)(3\alpha\beta\Omega_1^2+3\beta\Omega_1^2-3\alpha\beta+\alpha-3\beta+1)}\Omega_1}{\gamma(7\beta\Omega_1^2-3\beta+1)}, \\ \Omega_9 &= -2\frac{3\alpha\beta\Omega_1^2+3\beta\Omega_1^2-3\alpha\beta+\alpha-3\beta+1}{\beta\Omega_1(7\beta\Omega_1^2-3\beta+1)}, \\ \Xi[1] &= \frac{1}{3}\sqrt{-18\beta^2\Omega_1^4-9\beta^2\Omega_1^2+3\beta\Omega_1^2+27\beta^2-18\beta+3}.\end{aligned}$$

Case XII:

$$\begin{aligned}\mathbf{u}_{12} &= 2\frac{a_2\Omega_1^2H_2-a_3H_3\Omega_4^2+a_4H_4\Omega_7^2}{f} \\ &\quad - 2\frac{(-a_2\Omega_1H_2-a_3H_5\Omega_4+a_4H_6\Omega_7)^2}{f^2}, \\ f &= a_2H_2+a_3H_3+a_4H_4,\end{aligned}\quad (2.17)$$

$$\begin{aligned}H_2 &= e^{\Omega_1x+\Omega_2y+\Omega_3t}, \\ H_3 &= \cos(\Omega_4x+\Omega_5y+\Omega_6t), \\ H_4 &= \cosh(\Omega_7x+\Omega_8y+\Omega_9t),\end{aligned}$$

$$\begin{aligned}\Omega_2 &= \frac{4}{3}\frac{\Omega_1(3\alpha\beta\Omega_1^2+3\beta\Omega_1^2-3\alpha\beta+\alpha-3\beta+1)}{\Omega_8\gamma\Omega_7\beta}, \\ \Omega_3 &= \frac{(28\beta\Omega_1^2-27\beta+9)(\alpha+1)}{18\Omega_1(4\beta\Omega_1^2-3\beta+1)\beta}, \\ \Omega_4 &= \frac{\sqrt{\beta(3\beta\Omega_1^2-3\beta+1)}}{\beta}, \\ \Omega_6 &= -\frac{3\alpha\beta\Omega_1^2+3\beta\Omega_1^2-3\alpha\beta+\alpha-3\beta+1}{3\Omega_4\beta(4\beta\Omega_1^2-3\beta+1)},\end{aligned}$$

$$\Omega_5 = 2\frac{9\alpha\beta^2\Omega_1^4+9\beta^2\Omega_1^4-18\alpha\beta^2\Omega_1^2+6\alpha\beta\Omega_1^2-18\beta^2\Omega_1^2+9\alpha\beta^2+6\beta\Omega_1^2-6\alpha\beta+9\beta^2+\alpha-6\beta+1}{\gamma\beta^2\Omega_4\Omega_7\Omega_8},$$

We obtained forty sets of solutions as mentioned above, we neglect to bring those category of solutions. The three-dimensional dynamic graphs of the wave and corresponding density plots, and two-dimensional plots were successfully depicted in figures 1–8 with the help of the Maple. We can see that the exponential function, the cosine function, and the hyperbolic cosine function react with each other and move forward. Due to analyze the dynamics properties briefly, we would like to discuss the evolution characteristic. By selecting the suitable values of parameters, the analytical treatment of periodic wave solution is presented in 1 and 3 including 3D plot, density plot, and 2D plot, when three spaces arise at spaces $x = -10$, $x = 0$, and $x = 10$. In 1, in origin two lines interact together. But in 3, the periodic wave only move in the direction of the negative t -axis to positive t -axis in the (x, t) -plane. And also, by choosing the given values for 1 and 3, respectively

$$\begin{aligned}a_1 &= 0.5, \quad a_4 = 1, \\ \Omega_1 &= 1.5, \quad \alpha = 1, \\ \beta &= 2, \quad \gamma = -1, \quad y = -4, \\ a_3 &= 1.1, \quad a_4 = 1, \\ \Omega_7 &= 1.3, \quad \alpha = -1, \\ \beta &= -2, \quad \gamma = -1, \quad y = -4,\end{aligned}$$

it is worth mentioning that this periodic wave solutions have the following features as

$$\lim_{x \rightarrow \pm\infty} u(x, t) = 0, \quad \lim_{t \rightarrow \pm\infty} u(x, t) = 0. \quad (2.18)$$

By increasing x and t to large sufficient value, then the value of $u(x, t)$ will vanish. By considering $a_3 = 2.5$, $a_4 = 1$, $\Omega_7 = 1.3$, $\alpha = -2$, $\beta = 1$, $\gamma = 1$, $y = -4$, the solution $u_2(x, y, t)$ given by (2.8) expresses the move two parallel (x, t) -periodic breathers as line $x = t$. By plugging $a_1 = 0.5$, $a_3 = 1$, $\Omega_1 = 1.5$, $\alpha = -2$, $\beta = 2$, $\gamma = 1$, $y = -4$, in 4, we can see that in origin two lines interact together. In 5, the x -periodic breather is presented. In 6 b, by inserting $a_2 = 0.5$, $a_3 = 1.5$, $a_4 = 1$, $\Omega_1 = 1.5$, $\alpha = 1$, $\beta = \frac{1}{4}$, $\gamma = -1$, $y = -4$, two lump solutions there are in direction of (x, t) -periodic breathers. Finally in 7 and 8 the periodic breather move in directions of (x, t) -periodic breathers and t -periodic breathers, respectively.

3. Stability analysis of KP-BBM equation

In this section, the concept of linear stability analysis will be applied to study the stability analysis for the giving equation (1.5). The perturbed solution of the KP-BBM equation given by

$$u(x, y, t) = q + \lambda U(x, y, t), \quad (3.1)$$

in above the relation constant q is a steady state solution of equation (3.1). Substituting (3.1) into equation, one can obtain

$$\begin{aligned} & 2 \left(\frac{\partial^2}{\partial x^2} U(x, y, t) \right) U(x, y, t) \alpha \lambda \\ & + 2 \left(\frac{\partial}{\partial x} U(x, y, t) \right)^2 \alpha \lambda + 2 \left(\frac{\partial^2}{\partial x^2} U(x, y, t) \right) \alpha q \\ & + \left(\frac{\partial^4}{\partial x^3 \partial t} U(x, y, t) \right) \beta \\ & + \gamma \frac{\partial^2}{\partial y^2} U(x, y, t) + \frac{\partial^2}{\partial x \partial t} U(x, y, t) \\ & + \frac{\partial^2}{\partial x^2} U(x, y, t) = 0, \end{aligned} \quad (3.2)$$

by linerization equation (3.2), we get

$$\begin{aligned} & 2 \left(\frac{\partial^2}{\partial x^2} U(x, y, t) \right) \alpha q \\ & + \left(\frac{\partial^4}{\partial x^3 \partial t} U(x, y, t) \right) \beta \\ & + \gamma \frac{\partial^2}{\partial y^2} U(x, y, t) + \frac{\partial^2}{\partial x \partial t} U(x, y, t) \\ & + \frac{\partial^2}{\partial x^2} U(x, y, t) = 0. \end{aligned} \quad (3.3)$$

Theorem 3.1. Presume that the solution of equation (3.3) has the bellow form

$$U(x, y, z, t) = \rho_1 e^{i(Mx + Ny + Wt)}, \quad (3.4)$$

where M, N is the normalized wave numbers, by inserting (3.4) into equation (3.3), then by solving for W , we can get

the following relation

$$W(M, N) = \frac{2 M^2 \alpha q + N^2 \gamma + M^2}{M(M^2 \beta - 1)}. \quad (3.5)$$

Proof. By substituting the relation (3.4) in the linear PDE (3.3), we get

$$\begin{aligned} & 2 \left(\frac{\partial^2}{\partial x^2} U(x, y, t) \right) \alpha q + \left(\frac{\partial^4}{\partial x^3 \partial t} U(x, y, t) \right) \beta \\ & + \gamma \frac{\partial^2}{\partial y^2} U(x, y, t) + \frac{\partial^2}{\partial x \partial t} U(x, y, t) + \frac{\partial^2}{\partial x^2} U(x, y, t) \\ & = e^{i(Mx + Ny + Wt)} \rho_1 \\ & \times (M^3 W \beta - 2 M^2 \alpha q - N^2 \gamma - M^2 - MW) = 0. \end{aligned} \quad (3.6)$$

By solving and simplifying we can find the value of $W(M, N, P)$ as form

$$W(M, N, P) = \frac{2 M^2 \alpha q + N^2 \gamma + M^2}{M(M^2 \beta - 1)}. \quad (3.7)$$

Therefore, we get to the required solution. Then the proof of the theorem is complete. \square

The relations for the propagation in equation (3.5) is investigated. The sign of $W(M, N)$ proposes either the solution will become larger or decay in a given period of time. When the sign of $W(M, N)$ is negative for all value of M , then any superposition of solutions of the form $e^{i(Mx + Ny + Wt)}$ will come to decay and the steady state is stable. In other hand, if the $W(M, N)$ is positive for some values of M , then with time some components of a superposition will become bigger rapidly, the steady state is unstable. If the maximum $W(M, N)$ of the is exactly 0, the dispersion is called marginally stable.

4. Conclusion

In this paper, the periodic wave solutions of the KP-BBM equation have been constructed. From the bilinear form of this equation, two test functions or ansatzes have been chosen. At first, we chose the ansatz as a combination of the exponential, cosine and hyperbolic cosine functions. The 3D, 2D, and density graphs illustrating some solutions were represented, showing different periodic waveforms. As a consequence, some new solutions, which include the new multi wave, breather wave, periodic, cross-kink wave solutions were caught. Through of Maple, the evolution phenomenon of these waves is seen in figures 1–11, respectively. Mainly, by choosing specific parameter constraints all cases the two-dimension, and three-dimension in solitons can be captured from the multi wave and periodic wave solutions. The obtained solutions are extended with numerical simulation to analyze graphically, which results into multi wave, breather wave, periodic, cross-kink wave solutions. Moreover, we studied the linear stability analysis of the KP-BBM equation in the previous section. That will be extensively used

to report many attractive physical phenomena in the fields of acoustics, heat transfer, fluid dynamics, classical mechanics and so on.

Conflict of Interest

The authors declare that they have no conflict of interest.

Funding

There is no fund for this manuscript.

ORCID iDs

Jalil Manafian  <https://orcid.org/0000-0001-7201-6667>
Onur Alp Ilhan  <https://orcid.org/0000-0003-1618-6439>
As'ad Alizadeh  <https://orcid.org/0000-0003-4678-7470>

References

- [1] Manafian J and Lakestani M 2016 Abundant soliton solutions for the Kundu-Eckhaus equation via $\tan(\phi/2)$ -expansion method *Optik* **127** 5543–51
- [2] Manafian J 2015 On the complex structures of the Biswas-Milovic equation for power, parabolic and dual parabolic law nonlinearities *Eur. Phys. J. Plus* **130** 1–20
- [3] Baskonus H M and Bulut H 2016 Exponential prototype structures for (2+1)-dimensional Boiti–Leon–Pempinelli systems in mathematical physics *Waves Random Complex Medium* **26** 201–8
- [4] Manafian J, Foroutan M R and Guzali A 2017 Applications of the ETEM for obtaining optical soliton solutions for the Lakshmanan–Porsezian–Daniel model *Eur. Phys. J. Plus* **132** 494
- [5] Manafian J and Aghdaei M F 2016 Abundant soliton solutions for the coupled Schrödinger–Boussinesq system via an analytical method *Eur. Phys. J. Plus* **131** 97
- [6] Zhou Q, Ekici M, Sonmezoglu A, Manafian J, Khaleghizadeh S and Mirzazadeh M 2016 Exact solitary wave solutions to the generalized Fisher equation *Optik* **127** 12085–92
- [7] Manafian J and Heidari S 2019 Periodic and singular kink solutions of the Hamiltonian amplitude equation *Adv. Math. Models Appl.* **4** 134–49
- [8] Manafian J and Lakestani M 2016 Dispersive dark optical soliton with Tzitzéica type nonlinear evolution equations arising in nonlinear optics *Opt. Quantum Electron.* **48** 1–32
- [9] Sindi C T and Manafian J 2016 Wave solutions for variants of the KdV–Burger and the K(n, n)–Burger equations by the generalized G'/G -expansion method *Math. Methods Appl. Sci.* **87** 114
- [10] Satsuma J and Ablowitz M J 1979 Two-dimensional lumps in nonlinear dispersive systems *J. Math. Phys.* **20** 1496
- [11] Ma W X, Zhou Y and Dougherty R 2016 Lump-type solutions to nonlinear differential equations derived from generalized bilinear equations *Int. J. Mod. Phys. B* **30** 1640018
- [12] Lü J, Bilge S, Gao X, Bai Y and Zhang R 2018 Abundant lump solution and interaction phenomenon to Kadomtsev–Petviashvili–Benjamin–Bona–Mahony equation *J. Appl. Math. Phys.* **6** 1733–47
- [13] Foroutan M R, Manafian J and Ranjbaran A 2018 Lump solution and its interaction to (3+1)-D potential-YTSF equation *Nonlinear Dyn.* **92** 2077–92
- [14] Tang Y N, Tao S Q and Guan Q 2016 Lump solitons and the interaction phenomena of them for two classes of nonlinear evolution equations *Comput. Math. Appl.* **72** 2334–42
- [15] Ma W X 2015 Lump solutions to the kadomtsev-petviashvili equation *Phys. Lett. A* **379** 1975–8
- [16] Yang J Y and Ma W X 2016 Lump solutions to the bKP equation by symbolic computation *Int. J. Modern Phys. B* **30** 1640028
- [17] Ma W X, Qin Z Y and Lv X 2016 Lump solutions to dimensionally reduced p-gKP and p-gbKP equations *Nonlinear Dyn.* **84** 923–31
- [18] Wang C J 2016 Spatiotemporal deformation of lump solution to (2+1)-dimensional KdV equation *Nonlinear Dyn.* **84** 697–702
- [19] Manafian J, Mohammadi-Ivatlo B and Abapour M 2019 Lump-type solutions and interaction phenomenon to the (2+1)-dimensional Breaking Soliton equation *Appl. Math. Comput.* **13** 13–41
- [20] Ilhan O A, Manafian J and Shahriari M 2019 Lump wave solutions and the interaction phenomenon for a variable-coefficient Kadomtsev–Petviashvili equation *Comput. Math. Appl.* **78** 2429–48
- [21] Ilhan O A and Manafian J 2019 Periodic type and periodic cross-kink wave solutions to the (2+1)-dimensional breaking soliton equation arising in fluid dynamics *Mod. Phys. Lett. B* **1950277** 1–26
- [22] Cattani C, Sulaiman T A, Baskonus H M and Bulut H 2018 Solitons in an inhomogeneous Murnaghan's rod *Eur. Phys. J. Plus* **133** 228
- [23] Baskonus H M 2019 Complex soliton solutions to the Gilson–Pickering model *Axioms* **8** 18
- [24] Khalique C M and Mhlanga I E 2018 Travelling waves and conservation laws of a (2+1)-dimensional coupling system with Korteweg–de Vries equation *Appl. Math. Nonlinear Sci.* **3** 241–54
- [25] Baskonus H M, Bulut H and Sulaiman T A 2019 New Complex Hyperbolic Structures to the Lonngren–Wave equation by Using Sine-Gordon Expansion Method *Appl. Math. Nonlinear Sci.* **4** 129–38
- [26] Eskitaş E I, Aktaş M B and Baskonus H M 2019 New Complex and Hyperbolic Forms for Ablowitz–Kaup–Newell–Segur Wave equation with Fourth Order *Appl. Math. Nonlinear Sci.* **4** 93–100
- [27] Yel G, Baskonus H M and Bulut H 2019 Regarding some novel exponential travelling wave solutions to the Wu–Zhang system arising in nonlinear water wave model *Indian J. Phys.* **93** 1031–9
- [28] Baskonus H M, Bulut H and Atangana A 2016 Smart Materials and Structures Paper On the complex and hyperbolic structures of the longitudinal wave equation in a magneto-electro-elastic circular rod *Smart Mater. Struct.* **25** 035022
- [29] Baskonus H M 2017 New complex and hyperbolic function solutions to the generalized double combined Sinh–Cosh–Gordon equation *AIP Conf. Proc.* vol 1798020018
- [30] Sulaiman T A, Yokus A, Gulluoglu N and Baskonus H M 2018 Regarding the numerical solutions of the Sharma–Tasso–Olver equation *ITM Web Conf.* **22** 01036
- [31] Yokus A, Baskonus H M, Sulaiman T A and Bulut H 2017 Numerical simulation and solutions of the two-component second order KdV evolutionary system *Numer. Methods Partial Diff. Equ.* **34** 211–27
- [32] Shafiq A, Rashidi M M, Hammouch Z and Khan I 2019 Analytical investigation of stagnation point flow of Williamson liquid with melting phenomenon *Phys. Scr.* **94** 035204

- [33] Zhao Z L, He L and Gao Y 2019 Rogue wave and multiple lump solutions of the (2+1)-dimensional benjamin-ono equation in fluid mechanics *Complexity* **2019** 249635
- [34] Huang L L and Chen Y 2017 Lump solutions and interaction phenomenon for (2+1)-dimensional Sawada-Kotera equation *Commun. Theor. Phys.* **67** 473–8
- [35] Lu J Q and Bilige S D 2017 Lump solutions of a (2+1)-dimensional bSK equation *Nonlinear Dyn.* **90** 2119–24
- [36] Manafian J and Lakestani M 2019 Lump-type solutions and interaction phenomenon to the bidirectional sawada-kotera equation *Pramana* **92** 41
- [37] Lü J, Bilige S and Chaolu T 2018 The study of lump solution and interaction phenomenon to (2+1)-dimensional generalized fifth-order KdV equation *Nonlinear Dyn.* **91** 1669–76
- [38] Wang C J, Dai Z D and Liu C F 2016 Interaction between kink solitary wave and rogue wave for (2+1)-dimensional burgers equation *Mediterr. J. Math.* **13** 1087–98
- [39] He B and Meng Q 2019 Bilinear form and new interaction solutions for the sixth-order Ramani equation *Appl. Math. Lett.* **98** 411–8
- [40] Zhang Y, Dong H H, Zhang X E *et al* 2017 Rational solutions and lump solutions to the generalized (3+1)-dimensional Shallow Water-like equation *Comput. Math. Appl.* **73** 246–52
- [41] Wang J, An H L and Li B 2019 Non-traveling lump solutions and mixed lump kink solutions to (2+1)-dimensional variable-coefficient caudrey dodd gibbon kotera sawada equation *Mod. Phys. Lett. B* **1950262**
- [42] Manafian J 2018 Novel solitary wave solutions for the (3+1)-dimensional extended Jimbo-Miwa equations *Comput. Math. Appl.* **76** 1246–60
- [43] Dai Z D, Liu J, Zeng X P and Liu Z J 2008 Periodic kink-wave and kinky periodic-wave solutions for the jimbo-miwa equation *Phys. Lett. A* **372** 5984–6
- [44] Geng X G and Ma Y L 2007 N-soliton solution and its wronskian form of a (3+1)-dimensional nonlinear evolution equation *Phys. Lett. A* **369** 285–9
- [45] Dehghan M, Manafian J and Saadatmandi A 2011 Application of the Exp-function method for solving a partial differential equation arising in biology and population genetics *Int. J. Numer. Methods Heat Fluid Flow* **21** 736–53
- [46] Ma W X and Zhu Z 2012 Solving the (3+1)-dimensional generalized KP and BKP equations by the multiple exp-function algorithm *Appl. Math. Comput.* **218** 11871–9
- [47] Dehghan M and Manafian J 2009 The solution of the variable coefficients fourth-order parabolic partial differential equations by homotopy perturbation method *Z. Nat.forsch. A* **64a** 420–30
- [48] Ramani A 1981 Inverse scattering, ordinary differential equations of Painlevé type and Hirota's bilinear formalism *Ann. New York Acad. Sci.* **373** 54–67
- [49] Wazwaz A M 2005 Exact solutions of compact and noncompact structures for the KP-BBM equation *Appl. Math. Comput.* **169** 700–12
- [50] Manakov S V, Zakharov V E, Bordag L A and Matveev V B 1977 *Phys. Lett. A* **63** 205–6
- [51] Zhao H Q and Ma W X 2017 Mixed lump-kink solutions to the KP equation *Comput. Math. Appl.* **74** 1399–405
- [52] Wazwaz A M 2008 The extended tanh method for new compact and noncompact solutions for the KP-BBM and the ZK-BBM equations *Chaos Solitons Fractals* **38** 1505–16
- [53] Saut J C and Tzvetkov N 2004 Global well-posedness for the KP-BBM equations *Appl. Math. Res. Express* **1** 1–16
- [54] Alam M N and Akbar M A 2014 Exact traveling wave solutions of the KP-BBM equation by using the new approach of generalized (G'/G)-expansion method *Springerplus* **3** 43
- [55] Tang S, Huang X and Huang W 2010 Bifurcations of travelling wave solutions for the generalized KP-BBM equation *Appl. Math. Comput.* **216** 2881–90
- [56] Lü J, Bilige S, Gao X, Bai Y and Zhang R 2018 Abundant lump solutions and interaction phenomena to the Kadomtsev–Petviashvili–Benjamin–Bona–Mahony equation *J. Appl. Math. Phys.* **6** 1733–47
- [57] Tan J, Deng Z H, Wu T and Tang B 2019 Propagation and interaction of magnetic solitons in a ferromagnetic thin film with the interfacial Dzyaloshinskii–Moriya interaction *J. Magn. Magn. Mater.* **475** 445–52
- [58] Deng Z H, Wu T, Tanga B, Wang X Y, Zhao H P and Deng K 2018 Breathers and rogue waves in a ferromagnetic thin film with the Dzyaloshinskii–Moriya interaction *Eur. Phys. J. Plus* **133** 450
- [59] Deng Z H, Chang X, Tan J N, Tang B and Deng K 2019 Characteristics of the lumps and stripe solitons with interaction phenomena in the (2+1)-dimensional Caudrey–Dodd–Gibbon–Kotera–Sawada equation *Int. J. Theor. Phys.* **58** 92–102
- [60] Daia Z, Lin S, Fu H and Zeng X 2010 Exact three-wave solutions for the KP equation *Appl. Math. Comput.* **216** 1599–604