

## On conditions for an operator to be in the class $\mathcal{S}_p$

A. S. Holevo

The Schatten classes  $\mathcal{S}_p$  of operators on a Hilbert space  $\mathcal{H}$  play a significant role in recent studies in quantum information theory (for instance, see [1]–[3] and references therein). In the present paper we give sufficient conditions for an operator to belong to  $\mathcal{S}_p$  in the case when the Hilbert space is the space of an irreducible representation of the canonical commutation relations (CCR) (for example,  $\mathcal{H} = L^2(\mathbb{R}^s)$  – the case often occurring in applications). The problem is closely related to the asymptotics of the singular values, and in this question we rely on the classical treatise [4] (see also the papers [5] and [6], devoted to the case  $\mathcal{H} = L^2(G)$ , where  $G$  is a bounded domain in  $\mathbb{R}^s$ ).

Let  $\mathcal{H}$  be a separable Hilbert space, and  $\mathcal{S}_p$  ( $p > 0$ ) the Schatten class consisting of all compact operators  $\tau$  on  $\mathcal{H}$  whose singular values  $s_n(\tau)$  satisfy the condition  $\sum_{n=1}^\infty |s_n(\tau)|^p < \infty$ . In what follows  $\mathcal{H}$  will be the space of an irreducible representation of the CCR:

$$a_j a_k^* - a_k^* a_j = \delta_{jk} I, \quad a_j a_k - a_k a_j = 0.$$

The annihilation operators  $a_j$  and creation operators  $a_j^*$  are necessarily unbounded, and, strictly speaking, these relations hold on a common invariant dense domain  $\mathcal{D}$  (for instance, see [7] and [8] for details and rigorous treatments). It is well known that the operators

$$N_j = a_j^* a_j = a_j a_j^* - I \tag{1}$$

are essentially selfadjoint and have pure point spectrum with eigenvalues  $n_j = 0, 1, 2, \dots$ . The operators  $N_j$ ,  $j = 1, \dots, s$ , commute, so that the eigenvalues of the total number operator  $N = \sum_{j=1}^s N_j$  have the form  $\sum_{j=1}^s n_j$ .

**Theorem.** *Let  $\tau \in \mathcal{S}_2$  and assume that for some  $k = 1, 2, \dots$  the range of  $\tau$  is contained in the domains  $\text{dom } a_j^k$  for all  $j = 1, \dots, s$  and that  $a_j^k \tau \in \mathcal{S}_2$ . Then*

$$s_n(\tau) = o(n^{-[k/(2s)+1/2]}), \tag{2}$$

which implies that  $\tau \in \mathcal{S}_p$  for all  $p > 2s/(k + s)$ .

The idea of the proof is to use the decomposition  $\tau = (I + N)^{-k/2} \sigma$ , where the operator  $\sigma = (I + N)^{k/2} \tau$  is well defined and can be shown to be Hilbert–Schmidt under the conditions of the theorem, so that by equation (III.7.12) in [4]

$$s_n(\sigma) = o(n^{-1/2}). \tag{3}$$

To prove that  $\sigma \in \mathcal{S}_2$  we use the inequality

$$(I + N)^k \leq (s + 1)^{k-1} \left( I + \sum_{j=1}^s N_j^k \right) \tag{4}$$

---

AMS 2010 Mathematics Subject Classification. Primary 47B10.

and the identity

$$N_j^k = \sum_{r=1}^k c_r (a_j^*)^r a_j^r \quad (c_r > 0, c_k = 1) \tag{5}$$

(following from the second equality in (1)), both holding on an appropriate domain containing  $\mathcal{D}$ .

The operator  $(I + N)^{k/2}$  has a compact inverse  $(I + N)^{-k/2}$ , so that by the properties of the singular values (see [4], Corollary II.2.2)

$$s_{2n}(\tau) \leq s_{2n-1}(\tau) \leq s_n((I + N)^{-k/2}) s_n(\sigma). \tag{6}$$

Arguing along the lines in [6], we obtain

$$s_n((I + N)^{-k/2}) = O(n^{-k/(2s)}). \tag{7}$$

Indeed, the eigenvalues of  $(I + N)^{-k/2}$  have the form  $(1 + \sum_{j=1}^s n_j)^{-k/2}$ ,  $n_j \geq 0$ . The number  $\#(R)$  of integer points  $(n_1, \dots, n_s)$  in the simplex  $\sum_{j=1}^s x_j \leq R$ ,  $x_j \geq 0$ , is estimated by its volume  $R^s/s!$ . Given  $n$ , take  $R$  such that  $s_n((I + N)^{-k/2}) = (1 + R)^{-k/2}$ ; then  $n \leq \#(R) = O(R^s) = O(s_n((I + N)^{-k/2})^{-2s/k})$ , hence (7) follows. Combining (7), (3), and (6), we get (2).

In the Schrödinger representation,  $\mathcal{H} = L^2(\mathbb{R}^s)$  and

$$a_j \psi = \frac{1}{\sqrt{2}} \left( x_j + \frac{\partial}{\partial x_j} \right) \psi = \frac{1}{\sqrt{2}} e^{-x_j^2/2} \frac{\partial}{\partial x_j} e^{x_j^2/2} \psi,$$

hence  $a_j^k \psi = 2^{-k/2} e^{-x_j^2/2} \partial^k e^{x_j^2/2} \psi / \partial x_j^k$  (all the derivatives are generalized). Then the condition of the theorem takes the following form: the kernel  $\tau(x, y)$  of the operator  $\tau$ , as well as all the functions

$$e^{-x_j^2/2} \frac{\partial^k}{\partial x_j^k} e^{x_j^2/2} \tau(x, y), \quad j = 1, \dots, s,$$

are square integrable with respect to  $(x, y)$ . Similar formulations are obtained for the Bargmann–Berezin representation in the space of holomorphic functions and for the ‘noncommutative Fourier transform’ of the operator  $\tau$  [8].

The action of the operator  $a$  is especially simple when  $\tau$  has the  $P$ -representation with respect to the coherent states:  $\tau = \int_{\mathbb{C}^s} P(z)|z\rangle\langle z| d^s z$  (here  $P(z)$  is a complex-valued function; see [8], for instance). Then the condition of the theorem takes the form

$$\int_{\mathbb{C}^s} \int_{\mathbb{C}^s} \overline{P(z)} P(w) \left[ 1 + \sum_{j=1}^s (z_j^* w_j)^k \right] \exp(-|z-w|^2) d^s z d^s w < \infty.$$

### Bibliography

- [1] R. L. Frank and E. H. Lieb, *J. Math. Phys.* **58**:6 (2017), 062204, 7 pp.
- [2] A. C. Холево, *УМН* **70**:2(422) (2015), 141–180; English transl., A. S. Holevo, *Russian Math. Surveys* **70**:2 (2015), 331–367.

- [3] G. De Palma, D. Trevisan, V. Giovannetti, and L. Ambrosio, *J. Math. Phys.* **59**:8 (2018), 081101, 25 pp.
- [4] И. Ц. Гохберг, М. Г. Крейн, *Введение в теорию линейных несамосопряженных операторов*, Наука, М. 1965, 448 с.; English transl., I. C. Gohberg and M. G. Kreĭn, *Introduction to the theory of linear nonselfadjoint operators*, Transl. Math. Monogr., vol. 18, Amer. Math. Soc., Providence, R.I. 1969, xv+378 pp.
- [5] S. Agmon, *Comm. Pure Appl. Math.* **15**:2 (1962), 119–147.
- [6] В. И. Параска, *Матем. сб.* **68(110)**:4 (1965), 623–631; English transl., V. I. Paraska, Amer. Math. Soc. Transl. Ser. 2, vol. 79, Amer. Math. Soc., Providence, R.I. 1969, pp. 87–95.
- [7] M. Reed and B. Simon, *Methods of modern mathematical physics*, vol. II: *Fourier analysis, self-adjointness*, Academic Press, New York–London 1975, xv+361 pp.
- [8] А. С. Холево, *Вероятностные и статистические аспекты квантовой теории*, 2-е изд., ИКИ, М.–Ижевск 2003, 404 с.; English transl., A. Holevo, *Probabilistic and statistical aspects of quantum theory*, 2nd ed., Quad./Monogr., vol. 1, Edizioni della Normale, Pisa 2011, xvi+323 pp.

**Alexander S. Holevo**

Steklov Mathematical Institute  
of Russian Academy of Sciences, Moscow  
E-mail: [holevo@mi-ras.ru](mailto:holevo@mi-ras.ru)

Presented by A. V. Bulinski

Accepted 05/DEC/19

Translated by THE AUTHOR