

Null second order corrections to Casimir energy in weak gravitational field

A.P.C.M Lima,^{a,1} G. Alencar,^a C.R. Muniz^b and R.R. Landim^a

^aUniversidade Federal do Ceará

Caixa Postal 6030, Campus do Pici, 60455-760, Fortaleza, Ceará, Brazil

^bGrupo de Física Teórica, Faculdade de Educação, Ciências e Letras de Iguatu, Universidade Estadual do Ceará,

Av. Dário Rabêlo, 63502-253 Iguatu, Ceará, Brazil

E-mail: augustopcml@gmail.com, geovamaci@gmail.com,
celiomuniz131@gmail.com, renan@fisica.ufc.br

Received May 1, 2019

Revised May 29, 2019

Accepted June 8, 2019

Published July 4, 2019

Abstract. The discussion of vacuum energy is currently a subject of great theoretical importance, specially concerning the cosmological constant problem in General Relativity. From Quantum Field Theory, it is stated that vacuum states subject to boundary conditions may generate tensions on these boundaries related to a measurable non-zero renormalized vacuum energy: the Casimir effect. As such, investigating how these vacuum states and energy behave in curved backgrounds is just natural and might provide important results in the near future. In this paper we revisit a model of the Casimir effect for a massless scalar field in a weak gravitational field background, with rectangular Dirichlet boundaries, which has been proposed and further generalized in the literature. A trick originally used to simplify calculations is shown to lead to a inconsistent value for the energy shift, and by performing explicit mode expansion we arrive at an unexpected result: null gravitational correction even at order $(M/R)^2$, in opposition to earlier results.

Keywords: gravity, quantum field theory on curved space

ArXiv ePrint: [1903.00512](https://arxiv.org/abs/1903.00512)

¹Corresponding author.

Contents

1	Introduction	1
2	Casimir effect in a weak gravitational field — brief review	2
2.1	Real massless scalar field in curved background	2
2.2	Casimir energy for the flat space-time case	3
2.3	Sorge’s result	3
2.4	Generalizations	5
3	Revisiting	6
3.1	Explicit calculation of second order corrections to the Casimir energy	7
3.2	Generalization to spherically symmetrical space-time	8
4	Conclusions	9

1 Introduction

One rather odd aspect of quantum field theory lies in the connection between quantum vacuum and the Casimir effect [1]: fluctuations of a confined field can generate measurable interactions even at vacuum states. Mathematically, this comes from the renormalization of the vacuum expected value of the stress-energy-momentum tensor in the presence of boundaries(or analogously topological identification conditions) [2]. This quite general character makes it possible to extend the original model of electromagnetic fluctuations trapped between conducting surfaces for various fields, boundary conditions, surface geometries and space-time topologies.

The first convincing experimental validations came about in 1997 [3] and by [4] in 1998. Although the model might seem theoretically simple, a number of practical difficulties arise when measuring casimir forces, such as deformation of the surfaces, corrections from finite conductivity, temperature, material boundary rugosity, among others. A few recent experiments are listed in references [5–9]. Unfortunately, as with many such measures expected at very high energy scales, present technology is very far from detecting tiny corrections such as those from extended standard-model theories and gravitational effects in the Casimir effect, though, it is possible to use current experimental data to impose bounds on parameters of the respective theories (more on this in the next paragraph).

For both theoretical and phenomenological interests, the study of Casimir effect in non-trivial space-time topologies has gotten attention in the past decades. For being intrinsically related to quantum vacuum fluctuations (some authors say however that the effect is not actually related to zero point fluctuations, see for example [10]), the investigation of the Casimir effect in curved background might be of cosmological value(look [11] for an general view at the cosmological problem). The subject of casimir interaction in curved backgrounds has already been extensively worked on, as can be exemplified from [12–44]. As recent and potentially important phenomenological applications, we can highlight a few examples such as the bounds found on a Lorentz violation parameter [36] (which is lower than previously found ones), on one from Kehagias-Sfetsos solution for Horava-Lifshitz gravity [43], and a proposed test for the Heisenberg-Coulomb theory based on gravitonic casimir effect [44].

In this work we revisit the model of [32] where an expression for Casimir energy for a massless real scalar field in a weak static gravitational field is derived. Through a perturbative method, first order (in M/R) corrections are shown to be zero and a second order energy shift is found. Works from other authors generalizing the space-time geometry are also exemplified [33–37]. Although the original approach is simple and elegant, in circumventing more lengthy calculations, an equivocated simplification is applied to obtain second order corrections in [32], leading to a inconsistent value for the Casimir energy. In this paper we rederive the results by performing explicit expansion of the mode solutions and carrying out the calculations directly, we show that the gravitational correction in the mean Casimir energy density obtained by the model is null even at second order. The result is also extended for a more general space-time equivalent to the one proposed at [33]. As a number of works followed from Sorge’s paper, the present authors hope that the results presented here might be helpful in revising earlier ones, including those that might be of phenomenological importance like [36] as well as fueling pertinent questions to the general study of Casimir effect and quantum field theory on curved space-time.

2 Casimir effect in a weak gravitational field — brief review

In this section we present a review of the fundamentals for Casimir effect in curved space-time and a few results from the literature, we follow heavily [32], which is the main reference of this paper.

2.1 Real massless scalar field in curved background

Consider a real massless scalar field is subjected to Dirichlet boundary conditions on rectangular parallel plates with a background static weak gravitational field. Measurements are taken by a static observer with four-velocity:

$$u^\mu = (-g_{00})^{-1/2} \delta_0^\mu. \quad (2.1)$$

The action of the scalar field in curved background is given by:

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi + \frac{1}{2} \epsilon R \phi^2 \right], \quad (2.2)$$

so that the field equation in curved space-time is:

$$\frac{1}{\sqrt{-g}} \partial_\mu [\sqrt{-g} g^{\mu\nu} \partial_\nu \phi] + \epsilon R \phi = 0. \quad (2.3)$$

To find a simple expression for the energy density, first we work out orthonormal mode solutions in the sense of the scalar product:

$$\langle \phi_m, \phi_n \rangle = i \int_\Sigma \sqrt{g_\Sigma} n^\mu [\phi_n^* \partial_\mu \phi_m - (\partial_\mu \phi_n^*) \phi_m] d\Sigma, \quad (2.4)$$

where Σ stands for the boundaries, and

$$\langle \phi(n, \vec{k}_n), \phi(m, \vec{k}_m) \rangle = \delta(\vec{k}_n - \vec{k}_m) \delta_{nm}. \quad (2.5)$$

Here the \vec{k} terms refers to the transverse wave numbers and the n, m are discrete modes generated by the boundary condition impositions on the field. In terms of these orthonormal modes the mean vacuum energy can be expressed as:

$$\bar{\epsilon} = \frac{1}{V_p} \sum_n \int d^2k \sqrt{g_\Sigma} (-g_{00})^{-1} T_{00}[\psi_n^*, \psi_n], \quad (2.6)$$

where V_p is the proper volume of the cavity and $T_{00}[\psi_n^*, \psi_n]$ is defined after the energy-momentum component:

$$T_{00}[\psi_n^*, \psi_n] = \partial_0 \psi_n^* \partial_0 \psi_n - g_{00} g^{\mu\nu} \partial_\mu \psi_n^* \partial_\nu \psi_n. \quad (2.7)$$

2.2 Casimir energy for the flat space-time case

In the case $g_{\mu\nu} = \eta_{\mu\nu}$, the field is governed by the usual Klein-Gordon equation. The boundaries are planes of coordinate separation L such that $\phi(z=0) = \phi(z=L) = 0$, orthonormal solutions are given by:

$$\phi(x) = \frac{1}{2\pi\sqrt{\omega_{nF}L}} \sin\left(\frac{n\pi z}{L}\right) \exp[-i(\omega_{nF}t - k_\perp x_\perp)], \quad (2.8)$$

where:

$$\omega_{nF} = \sqrt{k_\perp^2 + (n\pi/L)^2} \quad (2.9)$$

Is the mode frequency (F standing for flat). Applying these modes on equation (2.6) results in:

$$\bar{\epsilon} = \frac{1}{8\pi^2 L} \sum_n \int d^2k_\perp \omega_{nF}. \quad (2.10)$$

This integral diverges as expected. Using Schwinger's proper time representation and zeta function regularization we obtain the renormalized value for the Casimir energy in minkowski space-time:

$$\bar{\epsilon}_{cas} = -\frac{\pi^2}{1440L^4}. \quad (2.11)$$

2.3 Sorge's result

In reference [32], from F. Sorge, the Casimir energy is calculated perturbatively for the following weak gravitational field space-time metric:

$$ds^2 = -(1 + 2\Phi)dt^2 + (1 - 2\Phi)dl^2, \quad (2.12)$$

for rectangular Dirichlet boundaries of coordinate surface S and coordinate separation L . The factor Φ is expanded inside the cavity (to order $(M/R)^2$) as:

$$\Phi = -M/r \simeq -M/R + Mz/R^2 = \Phi_0 + \gamma z. \quad (2.13)$$

Introducing a rectangular coordinate system inside the cavity. The center of the inner plate has spatial coordinates $(0, 0, 0)$ while the center of the outer one has $(0, 0, L)$. We consider $L \ll \sqrt{S}, M$, so that the plates are finite but we can still approximate transverse modes by plane waves.

The line element then looks like

$$ds^2 = -(1 + 2\Phi_0 + 2\gamma z)dt^2 + (1 - 2\Phi_0 - 2\gamma z)dl^2, \quad (2.14)$$

with a coordinate change

$$dt \rightarrow (1 + 2\Phi_0)^{-1/2} dt \quad (2.15)$$

$$d\vec{x} \rightarrow (1 - 2\Phi_0)^{-1/2} d\vec{x}, \quad (2.16)$$

and we arrive at

$$ds^2 = -(1 + 2\gamma z)dt^2 + (1 - 2\gamma z)dl^2. \quad (2.17)$$

The line element now is the same as (2.14) with $\phi_0 = 0$. This is to tell us that up to order $[M/R]^2$ the parameter of interest is γ , that of the spatially varying field. It is shown in [38] that as long as there are no cross terms in the metric, the Casimir energy for a general constant perturbation measured by observer (2.1) is unchanged. This is physically expected, since those space-times are equivalent to the Minkowski one through a coordinate rescaling (actually any constant Φ_0 can be gauged away, weak or not).

The case $\gamma = 0$, or first order approximation is calculated explicitly and shown to be null in [32]. Next, for second order correction metric (2.17) is considered. For this space-time the field equation becomes

$$-(1 - 4\gamma z)\partial_t^2 \phi + \nabla^2 \phi = 0, \quad (2.18)$$

where $\nabla^2 \equiv \delta^{ij}\partial_i\partial_j$. With mode solutions of the form

$$\phi_{n,k} = \chi_n(z)e^{-i(\omega_n t - k_\perp x_\perp)}. \quad (2.19)$$

The asymptotic solutions are

$$\chi(u) = A_n u^{-1/4} \sin\left(\frac{2}{3}u^{3/2} + \varphi\right), \quad (2.20)$$

where:

$$\begin{aligned} u(z) &= -(z - b/a)a^{1/3} \\ a &= 4\gamma\omega_n^2 \\ b &= \omega_n^2 - k_\perp^2. \end{aligned} \quad (2.21)$$

And:

$$\omega_n = (1 + \gamma L)\omega_{nF} \quad (2.22)$$

Are the frequencies. Instead of working the modes explicitly the author considers an expansion in terms of the flat case solution (2.11)

$$\phi_n = \phi_n^{(0)} + \delta\phi_n. \quad (2.23)$$

Then, the energy-momentum tensor and Casimir energy are expanded as:

$$T_{00}[\phi_n^*, \phi_n] = T_{00}[\phi_n^{(0)*}, \phi_n^{(0)}] + \{T_{00}[\delta\phi_n^*, \phi_n^{(0)}] + c.c.\}, \quad (2.24)$$

and:

$$\bar{\epsilon} = \bar{\epsilon}^{(0)} + \delta\bar{\epsilon}. \quad (2.25)$$

By substituting the first term of the r.h.s. of (2.24) on (2.6) and renormalizing we obtain:

$$\bar{\epsilon}_{(\text{ren})}^{(0)} = -(1 - 2\gamma L_p) \frac{\pi^2}{1440 L_p^4}, \quad (2.26)$$

and the second term, in analogy to the flat space-time expression (2.10), is to be calculated by integrating the frequency change of the modes (other geometrical terms can be neglected since $\delta\phi$ itself is of order γ):

$$\delta\bar{\epsilon} = \frac{1}{8\pi^2 L} \sum_n \int d^2 k_\perp \delta\omega_n = \gamma L_p \left(-\frac{\pi^2}{1440 L_p^4} \right), \quad (2.27)$$

arriving at

$$\bar{\epsilon}_{\text{casimir}} = -(1 - \gamma L_p) \frac{\pi^2}{1440 L_p^4}, \quad (2.28)$$

which is the original result presented by the author.

2.4 Generalizations

In this subsection we briefly review a few generalizations proposed in the literature to the above result, calculations will not be shown explicitly, since accurate derivations of the Casimir energy will be presented in the next section. In [35] the authors consider the far field limit in the case of a rotating source, the metric used is

$$ds^2 = (1 + 2\phi)dt^2 - (1 - 2\phi)dl^2 - 4adtd\varphi, \quad (2.29)$$

where $b = 1 - 2a\Omega$, which is related to the rotation parameters. Through coordinate change it is cast into (first order terms are eliminated similarly to Sorge's case)

$$ds^2 = (1 + 2b\gamma z)dt^2 - (1 - 2\gamma z)dl^2. \quad (2.30)$$

The procedure from last subsection is followed tightly, the mean Casimir energy density is found to be

$$\bar{\epsilon}_{\text{cas}} = -[1 - \gamma L_p(1 + 3a\Omega)] \frac{\pi^2}{1440 L_p^4}. \quad (2.31)$$

Thermal corrections are also calculated but we will omit them for conciseness.

In [33] the space-time metric is proposed

$$ds^2 = (1 + 2\gamma_0 + 2\lambda_0 z)dt^2 - (1 + 2\gamma_1 + 2\lambda_1 z)dl^2, \quad (2.32)$$

which is then simplified to Fermi-Coordinates assuming the form

$$ds^2 = (1 + 2\lambda_0 z')dt^2 - dl'^2. \quad (2.33)$$

Notice that the parameters from the spacial part of the metric are eliminated from the metric, though they will be present in the boundary conditions. A general analysis is provided for scalar and vector fields with both Neumann and Dirichlet conditions, for our case (scalar field and Dirichlet boundaries) the result presented for the mean Casimir energy is

$$\bar{\epsilon}_{\text{cas}} = -\left(1 + \gamma_0 + \lambda_0 \frac{L_p}{2}\right) \frac{\pi^2}{1440 L^4}. \quad (2.34)$$

Note that only the denominator is expressed in terms of the proper length.

In [34] the background space-time is that of extended theories of gravity (see the reference for more detailed discussion and explanation on the parameters used). The metric is put in a familiar form:

$$ds^2 = (1 + 2\Phi_0 + 2\Lambda z)dt^2 - (1 - 2\Psi_0 - 2\Sigma z)dl^2. \quad (2.35)$$

The authors explain that in this theory there is a vacuum curvature scalar $\mathcal{R} \simeq \mathcal{R}_1 + \mathcal{R}_2 z$, so geometric coupling is considered in equation (2.3). The result obtained is

$$\bar{\epsilon}_{\text{cas}} = -[1 - 3(\Phi_0 + \Psi_0) - (2\Sigma - \Lambda)L_p] \frac{\pi^2}{1440L_p^4} + \frac{\epsilon\mathcal{R}_2}{192L_p}. \quad (2.36)$$

The Casimir effect analysed in the recent paper [37] it is quite similar to another reference that approaches extended theories of gravity [34, 36]. We found to be difficult to reproduce their results because in the Casimir effect part the authors practically only exhibit both the solution in Sorge modes and the normalization constant, and then they already present the result of the energy without greater details.

Notice that the afore mentioned results do not coincide if the proper parameter equivalence is established, also first order corrections are found, what is not in concordance with the arguments for first order correction in [32]. In the next section we will present explicit calculations to properly review these results.

3 Revisiting

Let us first review Sorge's trick for second order calculations. The same arguments which led to it could be used to compute the first order correction as well, let us see what it gives. For $\Phi = \Phi_0$ in (2.12), substituting mode (2.8) on (2.7) and integrating in (2.6) we get

$$\bar{\epsilon}^0 = \frac{1}{V_p} \frac{1}{(2\pi)^2 2L} (1 - 3\Phi_0) \sum_n \int d^2k \omega_{nF}. \quad (3.1)$$

Renormalizing and expressing in terms of L_P it becomes

$$\bar{\epsilon}_{\text{ren}}^0 = -(1 - 4\Phi_0) \frac{\pi^2}{1440L_p^4}. \quad (3.2)$$

The frequencies are $\omega_n \simeq (1 + 2\Phi_0)\omega_{nF}$ so we should have

$$\delta\bar{\epsilon} = -2\Phi_0 \frac{\pi^2}{1440L_p^4}. \quad (3.3)$$

Summing both contributions gives a nonzero energy shift. The assumption that $\delta\bar{\epsilon}$ in (2.25) can be calculated by substituting ω_{nF} for $\delta\omega$ in the flat case expression is inconsistent with the explicit mode expansion method, which in the case of constant perturbation, recovers the physically expected null correction. We will show next that the correction in the above equation is instead connected to a geometrical term reminiscing of the normalization condition (2.4).

3.1 Explicit calculation of second order corrections to the Casimir energy

We begin by casting (2.20) into a more familiar expression by rewriting this mode axial solution in terms of the flat space-time mode solution plus a perturbative term

$$\chi(z) = \frac{1}{2\pi\sqrt{\omega_0 L}} \sin\left(\frac{n\pi z}{L}\right) + \gamma\chi^{(1)}(z) + O[M/R]^3. \quad (3.4)$$

For that we need to find φ, A_n and ω_n . The frequencies are obtained by requiring periodicity of the solution ($\psi(0) = \psi(L) = 0$), according to

$$\omega_n \simeq (1 + \gamma L)\omega_0, \quad (3.5)$$

where $\omega_0 = [k^2 + (n\pi/L)^2]^{1/2}$. Requiring that $\chi(z=0) = 0$ immediately gives

$$\phi = -\frac{2}{3}u^{3/2}(0). \quad (3.6)$$

For A_n we first express (2.4) taken on a hypersurface $t = 0$ in terms of χ

$$\begin{aligned} \langle \psi(n, k_1), \psi(n, k_2) \rangle &= (\omega_n + \omega_m) \int_V d^3x (1 - 4\gamma z) \chi_n \chi_m e^{i(k_{1\perp} - k_{2\perp})x_\perp} = \\ &= \delta^2(k_{1\perp} - k_{2\perp}) \delta_{nm}, \end{aligned} \quad (3.7)$$

which leads to

$$A_n = \left[(2\pi)^2 2\omega \int_V d^3x (1 - 4\gamma z) \Theta_n^2 \right]^{-1/2}, \quad (3.8)$$

where $\Theta_n = u^{-1/4} \sin(\frac{2}{3}u^{3/2} + \varphi)$. Plugging the result on the mode solution leads to the correction

$$\begin{aligned} \chi^{(1)}(z) &= [2n\pi\omega_0^2 L^2 (L - z) z \cos(n\pi z/L) \\ &\quad + L(2n^2\pi^2 z + 2k^2 L^2 z - k^2 L^3) \sin(n\pi z/L)] / 4Ln^2\pi^3 \sqrt{\omega_0 L}. \end{aligned} \quad (3.9)$$

With the full form of the mode solution (3.4) it can be checked that equation (2.3), as well as boundary and normalization conditions are satisfied to order $(M/R)^2$.

Now all that is left is to use the definition (2.6). For $\chi(z)$, equation (2.7) reads

$$T_{00}[\psi_n^*, \psi_n] = \frac{1}{2}\omega_n^2 \chi_n^2 + \frac{1}{2}(1 + 4\gamma z)[k_\perp^2 \chi_n^2 + (\partial_z \chi_n)^2]. \quad (3.10)$$

The nonrenormalized vacuum energy density is

$$\bar{\epsilon} = \frac{1}{V_p} \frac{S}{8\pi^2 L} (1 + \gamma L/2) \sum_n \int d^2 k_\perp \omega_0. \quad (3.11)$$

The standard proper-time representation for the k_\perp integration and Riemann zeta function regularization lead to the renormalized Casimir energy density

$$\bar{\epsilon}_{\text{cas}} = -\frac{\pi^2}{1440L^4} (1 + 2\gamma L) = -\frac{\pi^2}{1440L_p^4}. \quad (3.12)$$

So, in terms of the proper (measured by static observer) length of the cavity, the mean Casimir energy density (and hence total energy) is just the same as the flat space-time case for a rigid cavity.

Either the expression (3.12) tells us something important or not is tricky to say, since in generalized coordinates it becomes a lot more difficult to distinguish between actual physical effects and those simulated by choice of coordinates. The “measured” Casimir energy density looking the same as that of flat space-time could be a effect of not considering covariant boundary conditions (remember that conditions are imposed on the coordinate surface and separation for the plates) for example. Putting those arguments aside, from a heuristic point of view it is interesting that the shift in the field solution is found in this calculation to be just enough to compensate for the geometrical factors. As this computation of the Casimir energy is a highly approximate and simplified model (moreover, for a specific choice of field, boundary, background and observer), this result is by no means a definitive answer, yet it draws attention for going against our naive intuition. Also, it calls for an revision of other works in the literature.

It is important to remark that measuring the Casimir energy in a curved background is not an completely closed objective concept and that little is known about the quantum vacuum in the presence of gravity. More studies are indeed needed in order to give solid predictions about the behavior of vacuum energy in curved backgrounds.

Ignoring these factors, although the result presented here might seem disappointing from the phenomenological point of view, gravitational corrections are still expected on thermal effects, non static space-times such as Kerr’s [42] as well as higher order corrections (sadly, even corrections of order $(M/R)^2$ are far from what would be detectable with current technology).

3.2 Generalization to spherically symmetrical space-time

A general spherically symmetrical space-time line element looks like (in isotropic coordinates)

$$ds^2 = -e^{2\phi_t(r)} dt^2 + e^{-2\phi_s(r)} (dr^2 + r^2 d\Omega^2), \quad (3.13)$$

where ϕ_s e ϕ_t are general functions of r (not (r,t) , since by Birkhoff’s theorem any spherically symmetrical solution will also be static). In the same fashion as the last section we can treat this problem (in weak field regime) using the line element

$$ds^2 = -(1 + 2\gamma_t z) dt^2 + (1 - 2\gamma_s z) dl^2. \quad (3.14)$$

The EoM for the scalar field is

$$-(1 - 2\gamma_t z) \partial_t^2 \phi + (1 + 2\gamma_s z) \nabla^2 \phi + (\gamma_t - \gamma_s) \partial_z \phi = 0. \quad (3.15)$$

With solutions of the form (2.19) we get

$$\partial_z^2 \chi + 2\gamma_- \partial_z \chi + (\omega^2 - k_\perp^2) \chi - 4\gamma_+ \omega^2 z \chi = 0, \quad (3.16)$$

where $\gamma_+ = (\gamma_s + \gamma_t)/2$ and $\gamma_- = (\gamma_t - \gamma_s)/2$. The solutions are

$$\chi(z) = A_n (1 - \gamma_- z) \Theta(u(z)), \quad (3.17)$$

Where $\Theta(u)$ and frequency modes are defined in analogy to the last section with $\gamma \rightarrow \gamma_+$.

It will be instructive to keep a couple terms implicit in the calculation. Let (3.7) be rewritten as

$$\langle \phi(n, k_1), \phi(n, k_2) \rangle = (\omega_n + \omega_m) \int_V d^3x (1 - \beta z) \chi_n \chi_m e^{i(k_{1\perp} - k_{2\perp})x_\perp}, \quad (3.18)$$

and (2.6) as

$$\bar{\epsilon} = \frac{1}{V_p} \sum_n \int d^3x \int d^2k (1 - \lambda z) T_{00}[\psi_n^*, \psi_n]. \quad (3.19)$$

Also, γ is replaced by γ_+ in (3.10). Applying the solution (3.17) to the corresponding expressions leads to the renormalized energy in the form

$$\bar{\epsilon}_{\text{ren}} = -\frac{S\pi^2}{1440L^3} \frac{1}{V_p} \left[1 + \frac{1}{2}(\beta - \lambda + 2\gamma_+)L \right] = -\frac{\pi^2}{1440L_p^4}. \quad (3.20)$$

In this form it is easier to see where each term comes from. One can check that inserting the Minkowski solution in (3.19) leads to the γ_+ and λ term while the correction associated exclusively to the shift in the modes is

$$\delta\bar{\epsilon}_{\text{ren}} = \beta L \epsilon_0, \quad (3.21)$$

which for the original case was $\delta\bar{\epsilon} = 2\gamma\epsilon_0$. That leads to the zero correction of our calculation. So the multiplicative factor involved is not directly associated to the frequency correction as assumed in [32], but rather to the spatial metric from the normalization integral. So, also for this case no gravitational correction is expected. As discussed in the end of last subsection these calculations are very simplified, so the result could suffer drawbacks in contrast to an “exact” solution. But as far as internal consistency of the calculation goes, replacing (2.27) by (3.21) is strictly necessary.

Of course, this is for zero temperature, perfectly plane and conducting surfaces, among other idealizations. These are not the case in any real world experiments, so using this model to propose phenomenological tests to gravity theories as proposed in [34, 36, 37] might very well still be done by considering gravitational terms arising from thermal corrections originally considered in these references.

4 Conclusions

In this paper we have revisited the problem of a Casimir apparatus of parallel plates in the weak gravitational background described by (2.17). We followed the model from [32], now deriving the second order mode solution explicitly, finding a null correction for the mean Casimir energy density, in variance with the former author and, consequently, with all the works based in it, including the more recent ones which take into account extended theories of gravity [36, 37]. The method originally used was discussed and shown to fail when applied to the first order correction. We generalize this result to a static spherically symmetrical space-time (3.14). The resulting corrections are also zero, in variance with [33, 34].

From a phenomenological point of view, the results obtained here are not as interesting as the previous ones since there is no actual energy shift, although corrections may appear when considering thermal effects, finiteness and border effects of the boundaries, as well as higher corrections. An useful analysis that should to be made in the sense here discussed is with respect to the stationary (Kerr-like) space-times, which according to the literature

already presents first order gravitational corrections in the Casimir energy density [42], this is in accordance with [38], since those effects can show up at non-static space-times even if metric coefficients are constant.

Computation of the “vacuum” energy is not, by itself, a closed theoretical concept. Moreover, the one presented here and in the main aforementioned references is a very simplified and approximate one, so there is a number of factors not considered here that could affect the result. Putting those aside from the moment, this odd apparent absence of gravitational correction for the cases considered might be interesting though, from a theoretical point of view. It is curious that the action of the curved background up to second order is to exactly cast coordinate separation into proper distance as shown in equation (3.12). It also raises the question that, should the static gravitational field (3.14) generate no correction to the Casimir energy up to second order, then, on which order(or if) and from where should those effects actually emerge.

Although the obtained results appear to be partially frustrating in the sense of near future attempts of measuring gravitational changes in the Casimir effect, the interpretation of these results is to be tackled cautiously, and the question as to how vacuum energy fluctuations behave in curved space-times for simpler cases such as weak-field approximation might provide useful hints for the understanding of wider pictures. As such, the present authors hope to aid further discussions on the subject.

Acknowledgments

The authors would like to thank Alexandra Elbakyan and sci-hub, for removing all barriers in the way of science.

We acknowledge the financial support by Conselho Nacional de Desenvolvimento Científico e Tecnológico(CNPq) and Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico(FUNCAP) through PRONEM PNE0112-00085.01.00/16.

We also thank the referee for the insightful remarks provided.

References

- [1] H.B.G. Casimir, *On the attraction between two perfectly conducting plates*, *Indag. Math.* **10** (1948) 261 (*Kon. Ned. Akad. Wetensch. Proc.* **51** 793 (1948), *Front. Phys.* **65** 342 (1987), *Kon. Ned. Akad. Wetensch. Proc.* **100N3-4** 61 (1997)) [[INSPIRE](#)].
- [2] M. Visser, *Regularization versus renormalization: why are Casimir energy differences so often finite?*, *Particles* **2** (2018) 14 [[arXiv:1601.01374](#)] [[INSPIRE](#)].
- [3] S.K. Lamoreaux, *Demonstration of the Casimir force in the 0.6 to 6 micrometers range*, *Phys. Rev. Lett.* **78** (1997) 5 [*Erratum ibid.* **81** (1998) 5475] [[INSPIRE](#)].
- [4] U. Mohideen and A. Roy, *Precision measurement of the Casimir force from 0.1 to 0.9 micrometers*, *Phys. Rev. Lett.* **81** (1998) 4549 [[physics/9805038](#)] [[INSPIRE](#)].
- [5] C.-C. Chang, A.A. Banishev, G.L. Klimchitskaya, V.M. Mostepanenko and U. Mohideen, *Reduction of the Casimir force from indium tin oxide film by UV treatment*, *Phys. Rev. Lett.* **107** (2011) 090403.
- [6] A.A. Banishev, C.C. Chang, R. Castillo-Garza, G.L. Klimchitskaya, V.M. Mostepanenko and U. Mohideen, *Modifying the Casimir force between indium tin oxide film and Au sphere*, *Phys. Rev. B* **85** (2012) 045436 [[arXiv:1201.5585](#)] [[INSPIRE](#)].

- [7] C.C. Chang, A.A. Banishev, R. Castillo-Garza, G.L. Klimchitskaya, V.M. Mostepanenko and U. Mohideen, *Gradient of the Casimir force between Au surfaces of a sphere and a plate measured using an atomic force microscope in a frequency-shift technique*, *Phys. Rev. B* **85** (2012) 165443 [[arXiv:1204.3146](#)] [[INSPIRE](#)].
- [8] A.A. Banishev, C.C. Chang, G.L. Klimchitskaya, V.M. Mostepanenko and U. Mohideen, *Measurement of the gradient of the Casimir force between a nonmagnetic gold sphere and a magnetic nickel plate*, *Phys. Rev. B* **85** (2012) 195422 [[arXiv:1205.0872](#)] [[INSPIRE](#)].
- [9] A.A. Banishev, G.L. Klimchitskaya, V.M. Mostepanenko and U. Mohideen, *Demonstration of the Casimir force between ferromagnetic surfaces of a Ni-coated sphere and a Ni-coated plate*, *Phys. Rev. Lett.* **110** (2013) 137401 [[arXiv:1211.1470](#)] [[INSPIRE](#)].
- [10] H. Nikolic, *Proof that Casimir force does not originate from vacuum energy*, *Phys. Lett. B* **761** (2016) 197 [[arXiv:1605.04143](#)] [[INSPIRE](#)].
- [11] J. Martin, *Everything you always wanted to know about the cosmological constant problem (but were afraid to ask)*, *Comptes Rendus Physique* **13** (2012) 566 [[arXiv:1205.3365](#)] [[INSPIRE](#)].
- [12] G. Bimonte, G. Esposito and L. Rosa, *From Rindler space to the electromagnetic energy-momentum tensor of a Casimir apparatus in a weak gravitational field*, *Phys. Rev. D* **78** (2008) 024010 [[arXiv:0804.2839](#)] [[INSPIRE](#)].
- [13] J.S. Dowker and R. Critchley, *Covariant Casimir calculations*, *J. Phys. A* **9** (1976) 535.
- [14] L.H. Ford, *Quantum vacuum energy in a closed universe*, *Phys. Rev. D* **14** (1976) 3304 [[INSPIRE](#)].
- [15] M.B. Altaie and J.S. Dowker, *Spinor fields in an Einstein universe: finite temperature effects*, *Phys. Rev. D* **18** (1978) 3557 [[INSPIRE](#)].
- [16] S.S. Baym and M. Oezcan, *Casimir energy in a curved background with a spherical boundary: an exactly solvable case*, *Phys. Rev. D* **48** (1993) 2806 [[INSPIRE](#)].
- [17] M. Karim, A.H. Bokhari and B.J. Ahmedov, *The Casimir force in the Schwarzschild metric*, *Class. Quant. Grav.* **17** (2000) 2459 [[INSPIRE](#)].
- [18] M.R. Setare, *Casimir stress for concentric spheres in de Sitter space*, *Class. Quant. Grav.* **18** (2001) 4823 [[hep-th/0105236](#)] [[INSPIRE](#)].
- [19] E. Elizalde, S. Nojiri, S.D. Odintsov and S. Ogushi, *Casimir effect in de Sitter and anti-de Sitter brane worlds*, *Phys. Rev. D* **67** (2003) 063515 [[hep-th/0209242](#)] [[INSPIRE](#)].
- [20] A.N. Aliev, *Casimir effect in the space-time of multiple cosmic strings*, *Phys. Rev. D* **55** (1997) 3903 [[INSPIRE](#)].
- [21] E. Elizalde, *Uses of zeta regularization in QFT with boundary conditions: A Cosmo-topological Casimir effect*, *J. Phys. A* **39** (2006) 6299 [[hep-th/0607185](#)] [[INSPIRE](#)].
- [22] E.R. Bezerra de Mello, V.B. Bezerra and A.A. Saharian, *Electromagnetic Casimir densities induced by a conducting cylindrical shell in the cosmic string spacetime*, *Phys. Lett. B* **645** (2007) 245 [[hep-th/0611310](#)] [[INSPIRE](#)].
- [23] E. Elizalde, S.D. Odintsov and A.A. Saharian, *Repulsive Casimir effect from extra dimensions and Robin boundary conditions: from branes to pistons*, *Phys. Rev. D* **79** (2009) 065023 [[arXiv:0902.0717](#)] [[INSPIRE](#)].
- [24] M. Nouri-Zonoz and B. Nazari, *Vacuum energy and the spacetime index of refraction: A New synthesis*, *Phys. Rev. D* **82** (2010) 044047 [[arXiv:0904.2904](#)] [[INSPIRE](#)].
- [25] B. Nazari and M. Nouri-Zonoz, *Electromagnetic Casimir effect and the spacetime index of refraction*, *Phys. Rev. D* **85** (2012) 044060 [[arXiv:1110.4257](#)] [[INSPIRE](#)].
- [26] A.A. Saharian and M.R. Setare, *Casimir effect for curved boundaries in Robertson-Walker spacetime*, *Class. Quant. Grav.* **27** (2010) 225009 [[arXiv:1006.2709](#)] [[INSPIRE](#)].

- [27] V.B. Bezerra, G.L. Klimchitskaya, V.M. Mostepanenko and C. Romero, *Thermal Casimir effect in closed Friedmann universe revisited*, *Phys. Rev. D* **83** (2011) 104042 [[INSPIRE](#)].
- [28] V.B. Bezerra, V.M. Mostepanenko, H.F. Mota and C. Romero, *Thermal Casimir effect for neutrino and electromagnetic fields in closed Friedmann cosmological model*, *Phys. Rev. D* **84** (2011) 104025 [[arXiv:1110.4504](#)] [[INSPIRE](#)].
- [29] K. Milton and A.A. Saharian, *Casimir densities for a spherical boundary in de Sitter spacetime*, *Phys. Rev. D* **85** (2012) 064005 [[arXiv:1109.1497](#)] [[INSPIRE](#)].
- [30] E.R. Bezerra de Mello and A.A. Saharian, *Topological Casimir effect in compactified cosmic string spacetime*, *Class. Quant. Grav.* **29** (2012) 035006 [[arXiv:1107.2557](#)] [[INSPIRE](#)].
- [31] F. Sorge, *Casimir effect and free fall in a Schwarzschild black hole*, in *15th Marcel Grossmann meeting on recent developments in theoretical and experimental general relativity, astrophysics and relativistic field theories (MG15) Rome, Italy, 1–7 July 2018*, 2018, [arXiv:1807.03968](#) [[INSPIRE](#)].
- [32] F. Sorge, *Casimir effect in a weak gravitational field*, *Class. Quant. Grav.* **22** (2005) 5109 [[INSPIRE](#)].
- [33] B. Nazari, *Casimir effect of two conducting parallel plates in a general weak gravitational field*, *Eur. Phys. J. C* **75** (2015) 501 [[arXiv:1508.02150](#)] [[INSPIRE](#)].
- [34] G. Lambiase, A. Stabile and A. Stabile, *Casimir effect in extended theories of gravity*, *Phys. Rev. D* **95** (2017) 084019 [[arXiv:1611.06494](#)] [[INSPIRE](#)].
- [35] V.B. Bezerra, H.F. Mota and C.R. Muniz, *Casimir effect due to a slowly rotating source in the weak field approximation*, *Phys. Rev. D* **89** (2014) 044015 [[arXiv:1401.2084](#)] [[INSPIRE](#)].
- [36] M. Blasone, G. Lambiase, L. Petruzzello and A. Stabile, *Casimir effect in post-Newtonian gravity with Lorentz-violation*, *Eur. Phys. J. C* **78** (2018) 976 [[arXiv:1808.04425](#)] [[INSPIRE](#)].
- [37] L. Buoninfante, G. Lambiase, L. Petruzzello and A. Stabile, *Casimir effect in quadratic theories of gravity*, *Eur. Phys. J. C* **79** (2019) 41 [[arXiv:1811.12261](#)] [[INSPIRE](#)].
- [38] A. Zhang, *Theoretical analysis of Casimir and thermal Casimir effect in stationary space-time*, *Phys. Lett. B* **773** (2017) 125 [[arXiv:1710.03100](#)] [[INSPIRE](#)].
- [39] V.B. Bezerra, H.F. Mota and C.R. Muniz, *Remarks on a gravitational analogue of the Casimir effect*, *Int. J. Mod. Phys. D* **25** (2016) 1641018 [[INSPIRE](#)].
- [40] G.L. Klimchitskaya and V.M. Mostepanenko, *New constraints on Yukawa-type corrections to Newtonian gravity at short separations*, *Grav. Cosmol.* **20** (2014) 3 [[arXiv:1403.5778](#)] [[INSPIRE](#)].
- [41] G.L. Klimchitskaya and V.M. Mostepanenko, *Constraints on axion and corrections to Newtonian gravity from the Casimir effect*, *Grav. Cosmol.* **21** (2015) 1 [[arXiv:1502.07647](#)] [[INSPIRE](#)].
- [42] F. Sorge, *Casimir energy in Kerr space-time*, *Phys. Rev. D* **90** (2014) 084050 [[INSPIRE](#)].
- [43] C.R. Muniz, V.B. Bezerra and M.S. Cunha, *Hořava-Lifshitz gravity effects on Casimir energy in weak field approximation and infrared regime*, *Phys. Rev. D* **88** (2013) 104035 [[arXiv:1311.2570](#)] [[INSPIRE](#)].
- [44] J.Q. Quach, *Gravitational Casimir effect*, *Phys. Rev. Lett.* **114** (2015) 081104 [Erratum *ibid.* **118** (2017) 139901] [[arXiv:1502.07429](#)] [[INSPIRE](#)].